

## REFERENCES

- Acuña, J. G. (1938): Catálogo descriptivo de las orquídeas cubanas. *Est. Exp. Agron.* Santiago de las Vegas No. 60. 221 pp.
- Acuña, J. G. (1964): Flora de las ciénagas de Cuba. *Poeyana Ser. A.* No. 3. 10 pp.
- Acuña, J. G. and Borhidi, A. (1971): Catálogo preliminar de los helechos cubanos. Mscr. 30 pp.
- Acuña, J. G., Pinal, del S. Clara and Acuña, O. Gea (1967): Catálogo florístico de Isla de Pinos. *Acad. Sci. Cuba Ser. Isla de Pinos* No. 22. 54 pp.
- Acuña, J. G., Roig, T. M. and Sanchez, L. A. (1959): Informes sobre reconocimiento botánico de la península de Guanahacabibes. *Sub-proyecto No. 1. Mem. Est. Exp. Agron.* Santiago de las Vegas 137—140.
- Adams, C. D. (1972): *Flowering Plants of Jamaica*. Univ. West Indies Mona, Jamaica. 848 pp.
- Adams, D. A. (1963): Factors influencing vascular plant zonation in North Carolina Salt Marshes. *Ecology* **44**: 445—456.
- Aguerrevere, S. E. et al. (1939): Exploración de la Gran Sabana. *Rev. Fomento* **19**: 501—726.
- Aiello, A. (1979): A re-examination of Portlandia (Rubiaceae) and associated taxa. *J. Am. Arb.* **60**: 38—126.
- Airy Shaw, H. K. (1965): Diagnosis of new families, new names, etc. *Kew. Bull.* **18**: 249—273.
- Alain, H. Liogier (1946): Notas taxonomicas y ecológicas sobre la flora de Isla de Pinos. *Contr. Ocas. Mus. Hist. Nat. Coll. La Salle* No. 7. 11 pp.
- Alain, H. Liogier (1950): Notas sobre la vegetación de la Loma de Cajalbana, Pinar del Rio. *Rev. Soc. Bot. Cubana* **7** (1—2—3): 8—18.
- Alain, H. Liogier (1953): El endemismo en la flora de Cuba. *Mem. Soc. Hist. Nat. Cubana* **21** (2): 187—193.
- Alain, H. Liogier (1954): Segunda herborización por el Toa. Exploración del Pico Galano. *Rev. Soc. Bot. Cubana* **11** (4): 87—101.
- Alain, H. Liogier (1956): Exploraciones botánicas en la Sierra del Cristal. *Rev. Soc. Bot. Cubana* **13** (1—3): 24—39.
- Alain, H. Liogier (1958): La flora de Cuba: sus principales características, su origen probable. *Rev. Soc. Bot. Cubana* **15** (2—3): 36—59.
- Alain, H. Liogier (1960): Flora de Baracoa. *Rev. Soc. Geogr. Cubana* **30** (4): 13—22.
- Alain, H. Liogier (1962): *Flora de Cuba* Vol. V. Sucre, Caracas 362 pp.
- Alain, H. Liogier (1968): Novitates Antillanae III. *Brittonia* **20**: 148—161.
- Alain, H. Liogier (1969): Suplemento de la Flora de Cuba. Sucre, Caracas; 150 pp.
- Alain, H. Liogier (1978): La flora de Española: análisis, origen probable. *Colecc. Conf. Botánica. Acad. Cien. Rep. Domin.* **3**: 3—32.
- Alain, H. Liogier (1980): Novitates Antillanae, VIII. *Phytologia* **47**: 167—198.
- Alain, H. Liogier (1981): *Flora of Hispaniola: Part I. Phytol. Mem.* III. Moldenke, New Jersey, 218 pp.
- Alain, H. Liogier (1983): *Flora de la Española, II.* Univ. Centr. del Este Vol. 44. Ser. Cien 15. San Pedro de Macoris, R. D. 420 pp.
- Alain, H. Liogier (1985a): *Flora de la Española III.* Univ. Centr. del Este Vol. 45 Ser. Cien 15. San Pedro de Macoris, R. D. 431 pp.
- Alain, H. Liogier (1985b): *Descriptive Flora of Puerto Rico and Adjacent Islands. Spermatophyta* Vol. 1. Univ. Puerto Rico, Santo Domingo. R. D. 352 pp.

- Alain, H. Liogier and L. F. Martorell (1982): *Flora of Puerto Rico and Adjacent Islands: a Systematic Synopsis*. Univ. Puerto Rico. Corripio Santo Domingo R. D. 342 pp.
- Alissov, B. P. et al. (1954): A Course in Climatology. *Climates of the Earth* Vol. 3. Gidrometeor. Izd. 320 pp.
- Allen, P. E. (1952): Distribution and variation in Roystonea. *Ceiba* **3** (1): 1–18.
- Alvarez Condé, J. (1958): *Historia de la Botánica en Cuba*. Publ. Junta Nac. Habana 353 pp.
- Andréanszky, G. (1954): Essai sur une système phylogénétique des groupements végétaux. *Ann. Mus. Hist. Nat. Hung.* **5**: 175–189.
- Andréanszky, G. (1956): Les étapes et les conditions biologiques de l'évolution de la flore tertiaire en Hongrie. *Acta Bot. Acad. Sci. Hung.* **2**: 221–240.
- Andréanszky, G. (1959): *Die Flora der Sarmatischen Stufe in Ungarn*. Akadémiai Kiadó, Budapest 360 pp.
- Andréanszky, G. (1962a): Das Trockenelement in der alttertiären Flora Mitteleuropas auf Grund paläobotanischer Forschungen in Ungarn. *Vegetatio Acta Geobot.* **11**: 95–111.
- Andréanszky, G. (1962b): Das Trockenelement in der jungtertiären Flora Mitteleuropas. *Vegetatio Acta Geobot.* **11**: 155–172.
- Antonovics, J., Bradshaw, A. D. and Turner, R. G. (1971): Heavy metal tolerance in plants. *Adv. Ecol. Res.* **7**: 1–85.
- Are de Cuba, — *Inst. Geogr. Acad. Cien. de Cuba*. Havana; 1965.
- Arens, K. (1958): O cerrado como vegetação oligotrofica. *Bol. Fac. Fil. Cienc. Letr. Univ. Sao Paulo* **224** (159): 59–77.
- Ares, J. O. (1972): Equitability, competition and seasonal succession in a plant community. *J. Ecol.* **60**: 325–331.
- Ares, J. O. and León, R. J. C. (1972): An ecological assessment of the influence of grazing on plant community structure. *J. Ecol.* **60**: 333–342.
- Aristeguieta, L. and Medina, E. (1965): Protección y quema de la sabana llanera. *Bol. Soc. Venezol. Cien. Nat.* **26**: No. 109. 129–139.
- Arldt, W. (1917–1922): *Handbuch der Palaeogeographie I–II*. Bornträger, Leipzig.
- Ascanio, O., Perez, J. M. and Hernandez, A. (1968): *Mapa genético de los suelos de Cuba I: 1 000 000*. Inst. de Suelos Acad. Scien. De Cuba. Havana.
- Ashton, P. S. (1964a): A quantitative phytosociological technique applied to tropical mixed rainforest. *Malay. For.* **27**: 307–317.
- Aston, P. S. (1964b): Ecological studies in the mixed dipterocarp forests in the Brunei State. *Oxford Forestry Mem.* No. **25**: Clarendon Press, Oxford.
- Asprey, G. F. and Robbins, R. G. (1953): The Vegetation of Jamaica. *Ecol. Monogr.* **23**: 359–412.
- Atlas de Cuba. 1978. Havana.
- Atlas Nacional de Cuba. 1970. La Habana. 132 pp.
- Aubréville, A. M. (1938): La forêt coloniale: Les forêts de l'Afrique occidentale française. *Annal. Acad. Sci. Colon.* Paris **9**: 1–245.
- Aubréville, A. M. (1947): The disappearance of the tropical forests of Africa. *Unasylva* **1**: 5–11.
- Aubréville, A. M. (1949): *Climats, forêts et désertification de l'Afrique tropicale*. Soc. Édit. Géogr. Marit. et Colon. Paris 628 pp.
- Aubréville, A. M. (1950): Le concept d'association dans la forêt dense équatoriale de la basse Côte d'Ivoire. *Bull. Soc. Bot. France* **98**: 145–158.
- Aubréville, A. M. (1964): La théorie astronomique de E. Bernard sur le balancement de l'équateur calorifique et ses conséquences sur les déplacements de la forêt équatoriale africaine. *Adansonia* **4**: 216–227.
- Aubréville, A. M. (1965a): Les étranges savanes des llanos de l'Orénoque. *Adansonia* **5**: 1–13.
- Aubréville, A. M. (1965b): Principes d'une systématique des formations végétales tropicales. *Adansonia* **5**: 153–196.
- Aubréville, A. M. (1965c): Conceptions modernes en bioclimatologie et classification des formations végétales. *Adansonia* **5**: 297–306.
- Aubréville, A. M. (1966a): Les lisières forêt-savane des régions tropicales. *Adansonia* **6**: 177–187.
- Aubréville, A. M. (1966b): Le Costa Rica quelques aspects du pays, de son climat, de sa végétation et de sa flore. *Adansonia* **6**: 29–54.

- Austin, M. P. and Greig-Smith, P. (1968): The application of quantitative methods to vegetation survey II. Some methodological problems of data from rainforests. *J. Ecol.* **56**: 827–844.
- Austin, M. P. and Orlóci, L. (1966): Geometric models in ecology II. An evaluation of some ordination techniques. *J. Ecol.* **54**: 217–227.
- Axelrod, D. I. (1963): Fossil floras suggest stable not drifting continents. *Geophys. Res.* **68**: 3257–3263.
- Axelrod, D. I. (1975): Evolution and biogeography of Madrean-Tethyan sclerophyll vegetation. *Ann. Mo. Bot. Gard.* **62**: 280–334.
- Azzi, G. (1954): *Écologie agricole*. Ballière et fils. Paris 414 pp.
- Bader, F. J. W. (1960): Die Verbreitung borealer und subarktischer Holzgewächse im Gebirgen des Tropengürtels. *Nova Acta Leop. N. F.* **23**, No. 148. 3–544.
- Bagnouls, F. and Gaussen, H. (1952): Saison sèche et Index xérothermique. — *Doc. pour les Cart. des Prod. Végét. Sér.: Généralités*. **3**: 1–48.
- Bagnouls, F. and Gaussen, H. (1957): Les climats biologiques et leur classification. — *Ann. de Géogr.* **355**: 193–220.
- Bagnouls, F. and Gaussen, H. (1964): Les climats tropicaux et la végétation. *Adansonia* **4**: 262–268.
- Bailey, L. H. (1938): *Thrinax — The Peaberry Palms*. *Gent. Herb.* **4**: 129–149.
- Balárová-Tuláčková, E. (1984): Phytosociological characteristics of the *Sclerio interruptae-Rhynchos-poretum subimberbis* Yaguaramas savanna, (Cuba). *Folia Geobot. Phytotax.* **19**: 317–321.
- Balárová-Tuláčková, E. (1986): Sinopsis preliminar de las comunidades sabanas de las regiones occidental y central de Cuba e Isla de la Juventud, incluyendo algunas comunidades de contacto. *DOC. Phytosoc.* **10**: 215–222.
- Balárová-Tuláčková, E. and Capote, R. P. (1982): A new savanna-like community of the Sierra del Rosario Mountains, Cuba. *Folia Geobot. Phytotax.* **17**: 137–148.
- Balárová-Tuláčková, E. and Capote, R. P. (1985a): Contribution to the knowledge of some savanna and forb-rich communities on the Isla de la Juventud (Cuba). *Folia Geobot. Phytotax.* **20**: 17–39.
- Balárová-Tuláčková, E. and Capote, R. P. (1985b): The *Cassio lineatae-Aristidetum neglectae*, a new association of the serpentine in Cuba. *Folia Geobot. Phytotax.* **20**: 177–183.
- Balárová-Tuláčková, E. and García, E. E. (1987): Contribución acerca de las comunidades secundarias de gramíneas en Cuba. *Phytocoenología* **15**: 39–49.
- Balárová-Tuláčková, E. and Surli, M. (1983): Contribution to the phytosociological characteristics of the Yaguaramas savannas (Cinefuegos Province, Cuba). *Folia Geobot. Phytotax.* **18**: 1–12.
- Ball, M. M. and Harrison, C. G. A. (1969): Origin of the Gulf and Caribbean and implications regarding ocean ridge extension, migration, and shear. *Gulf Coast Assoc. Geol. Soc. Trans.*, **19**: 287–294.
- Bannister, P. (1966): The use of subjective estimates of coverabundance as the basis for ordination. *J. Ecol.* **54**: 665–674.
- Basilevich, R., Drozdov, A. V. and Rodin, L. E. (1969): *Basic Theoretical Problems of Biological Productivity*. Nauka Acad. Sci. Leningrad.
- Baumann-Bodenheim, M. G. (1955): Über die Beziehungen der neu-caledonischen Flora zwischen tropischen und den südhemisphärisch subtropischen... *Ber. Geobot. Inst. Rübel* (1956) 64–71.
- Beard, J. S. (1942): Montane vegetation in the Antilles. *Caribb. For.* **3**: 61–74.
- Beard, J. S. (1944): Climax vegetation in tropical America. *Ecol.* **25**: 127–158.
- Beard, J. S. (1946a): The progress of plant succession on the Soufrière of St. Vincent. *J. Ecol.* **33**: 1–9.
- Beard, J. S. (1946b): The Mora forests of Trinidad, British West Indies. *Ibidem* **33**: 173–192.
- Beard, J. S. (1949): The natural vegetation of the Windward and Leeward Islands. *Oxford Forest Mem.* No. **21**.
- Bard, J. S. (1953): The savanna vegetation of northern tropical America. *Ecol. Monogr.* **23**: 149–215.
- Beard, J. S. (1955): The classification of tropical American vegetation types. *Ecology* **36**: 89–100.
- Beard, J. S. (1959): The origin of African Proteaceae. *J. S. Afr. Bot.* **25**: 231–235.
- Beard, J. S. (1967): Some Vegetation Types of Tropical Australia in relation to those of Africa and America. *J. Ecol.* **55**: 271–290.
- Beckett, P. H. T. (1965): The cation exchange equilibria of calcium and magnesium. *Soil. Sci.* **100**: 118–123.
- Belousov, V. V. (1970): Against the hypothesis of ocean-floor spreading. *Tectonophysics*, **9**: 489–511.
- Bennett, H. H. and Allison, R. V. (1928): *The Soils of Cuba*. Vol. 24: Trop. Plant Res. Found., Washington 410 pp.

- Benett, H. H. and Allison, R. V. (1966): *Los Suelos de Cuba y Algunos Nuevos Suelos de Cuba*. Habana, Ed. Rev. 375 pp.
- Berazin, R. (1979): La vegetación serpentinícola de "Lomas de Galindo" Canasi. Abstract of thesis CSC. Havana, 21 pp.
- Berazain, R. (1981a): Sobre el endemismo de la flórula serpentinícola de "Lomas de Galindo" Canasi. Habana, *Rev. Jard. Bot. Nac. Cuba* **2**: 29–47.
- Berazain, R. (1981b): Reporte preliminar de plantas serpentinícolas acumuladoras e hiperacumuladoras de algunos elementos. *Rev. Jard. Bot. Nac. Cuba* **2**: 48–59.
- Berry, E. W. (1934): Pleistocene plants from Cuba. *Bull. Torr. Club* **61**: 237–240.
- Bews, J. W. (1929): *The World's Grasses*. Longmans Green, London.
- Bharucha, F. R. and Shanbag, G. Y. (1957): Precipitation effectiveness in relation to the vegetation of India, Pakistan and Burma. *Bot. Memoirs Univ. Bombay* No. 3.
- Bisse, J. (1974): Nuevos arboles de la Flora de Cuba I. *Feddes Report* **85**: 587–608.
- Bisse, J. (1975): Nuevos arboles de la flora de Cuba. *Ciencias, Univ. Habana Ser. 10. Bot.* **2**: 1–23.
- Bisse, J. (1975): Die floristische Stellung und Gliederung Kubas. *Wiss. Z. Friedrich Schiller Univ. Jena* **24**: 365–371.
- Bisse, J. (1980): La subdivisión florística de la región nororiental de Cuba. *Rev. Jard. Bot. Nac. Cuba* **1**: (1) 11–18.
- Bisse, J., Lippold, H. and Casper, S. J. (1975): Beiträge zur Kenntnis der westindischen Pinguicula. *Wiss. Z. Friedrich Schiller Univ. Jena* **24**: 377–385.
- Bizot, M. (1973): Contributions à la flora Bryologique de Cuba. *The Bryologist* **76**: 340–346.
- Black, C. C. (1971): Ecological implications of dividing plants into groups with distinct photosynthetic production capacities. *Adv. Ecol. Res.* **7**: 87–114.
- Black, G. A., Dobzhansky, T. and Pavan, C. (1950): Some attempts to estimate species diversity and population density of trees in Amazonian forests. *Bot. Gaz.* **111**: 413–425.
- Blackie, J. R. (1965): *Comparison of Methods of Estimating Evaporation in East-Africa*. Paper presented at the Third Specialist Meeting on Applied Meteorology. Mgunga, Kenya
- Blackshaw, G. N. (1920): Magnesia impregnated soils. *S. Afric. J. Sci.* **17**: 171–178.
- Blydenstein, J. (1962): La sabana de Trachypogon del Alto Llano. *Bol. Soc. Venezol. Cien. Nat.* **23**: No. 102, 139–206.
- Blydenstein, J. (1963): Cambios en la vegetación después del protección contra el fuego I–II. *Bol. Soc. Venezol. Cien. Nat.* Nos. 100, 103, 233–238, 239–244.
- Blydenstein, J. (1967): Tropical savanna vegetation of the llanos of Columbia. *Ecology* **48**: 1–15.
- Borhidi, A. (1961): Klimadiagramme und klimazonale Karte Ungarns. *Ann. Univ. Budapest, Ser. Biol.* **4**: 21–50.
- Borhidi, A. (1969): *A növény és környezete* (Plant and environment) In Hungarian. In: Kárpáti–Sárkány (eds): *A növények világa*. Vol. II, Gondolat Kiadó, Budapest, 351–416.
- Borhidi, A. (1972a): La taxonomía del género *Platygyne* Merc. — *Ann. Mus. Nat. Hist. Nat. Hung.* **64**: 89–94.
- Borhidi, A. (1972b): La taxonomía del género *Heptanthus* Griseb. *Ann. Univ. Budapest, Ser. Biol.* **14**: 85–90.
- Borhidi, A. (1973): *Fundamentals of Geobotany of Cuba*. Thesis for D. Sc. In Hungarian. Budapest, 292 pp.
- Borhidi, A. (1974): *Kubában térképeztünk*. (Mapping in Cuba) In Hungarian. Budapest, Gondolat pp. 245+116 Tables
- Borhidi, A. (1975): The effects of serpentine soils on tropical flora and vegetation in Cuba. *XII. Intern. Bot. Congr.* Leningrad. Abstracts I. 110.
- Borhidi, A. (1976): *Fundamentals of Geobotany of Cuba*. In Hungarian Summary of D. Sc. theses. Budapest, 16 pp.
- Borhidi, A. (1977): *Tetrazygiopsis*, género nuevo de las Antillas, y el género *Tetrazygia* L. C. Rich. (Melastomataceae) en Cuba. *Acta Bot. Acad. Sci. Hung.* **23**: 33–39.
- Borhidi, A. (1980): New names and new species in the flora of Cuba II. *Acta Bot. Acad. Sci. Hung.* **26**: 257–277.
- Borhidi, A. (1981): Die Entwicklung der Flora von den Antillen im Lichte der Kontinentalverschiebungstheorie. Paper abstract. *Ber. Bot. Mus. Linz* **13**: 41.

- Borhidi, A. (1982a): Entwicklung der Flora der Antillen im Licht der Kontinentalverschiebungstheorie. *Stapfia* (Linz) **10**: 235–241.
- Borhidi, A. (1982b): Studies in Rondeletieae (Rubiaceae) III. The genera *Rogiera* and *Arachnothryx*. *Acta Bot. Hung.* **28**: 65–72.
- Borhidi, A. (1985): Phytogeographic survey of Cuba I. The phytogeographic characteristics and evolution of the flora of Cuba. *Acta Bot. Hung.* **31**: 3–34.
- Borhidi, A. (1988a): Vegetation dynamics of savannization processes in Cuba. *Vegetatio* **75**: 177–183.
- Borhidi, A. (1988b): The main vegetation units of Cuba. *Acta Bot.* **33**: 151–185.
- Borhidi, A. (1988c): Efectos de la roca serpentina en la adaptación y evolución de la flora y vegetación tropical en Cuba. *Acta Bot. Hung.* **34**: 123–174.
- Borhidi, A., Acuña, J. G. and Muñiz, O. (1973): New plants in Cuba III. *Acta Bot. Acad. Sci. Hung.* **19**: 37–45.
- Borhidi, A. and Fernandez, M. Z. (1981a): Studies in Rondeletieae (Rubiaceae), I. A new genus: *Roigella*. *Acta Bot. Acad. Sci. Hung.* **27**: 309–312.
- Borhidi, A. and Fernandez, M. Z. (1981b): Studies in Rondeletieae (Rubiaceae), II. A new genus: *Suberanthus*. *Acta Bot. Acad. Sci. Hung.* **27**: 313–316.
- Borhidi, A. and Fernandez, M. Z. (1984): Revisión del género *Machaonia* H. et B en Cuba. (Rubiaceas Cubanás, IV). *Acta Bot. Sci. Hung.* **30**: 27–46.
- Borhidi, A. and Herrera, R. (1977): Génesis, características y clasificación de los ecosistemas de sabana en Cuba. *Cien. Biol. Acad. Cien. Cuba* **1**: 115–130.
- Borhidi, A., Imchanitzkaya, N. N. and Muñiz, O. (1978): Dendrological novelties in Cuba. *Acta Agron. Acad. Sci. Hung.* **27**: 428–437.
- Borhidi, A., Járai-Komlódi, M. and Moncada, M. (1980): *Acunaeanthus*, a new genus of Rubiaceae. *Acta Bot. Acad. Sci. Hung.* **26**: 279–289.
- Borhidi, A. and Kereszty, Z. (1979): New names and new species in the flora of Cuba Antillas. *Acta Bot. Acad. Sci. Hung.* **25**: 1–37.
- Borhidi, A. and Muñiz, O. (1971a): Combinationes novae florae Cubanae I. *Bot. Közlem.* **58**: 175–178.
- Borhidi, A. and Muñiz, O. (1971b): New plants in Cuba I. *Acta Bot. Acad. Sci. Hung.* **17**: 1–36.
- Borhidi, A. and Muñiz, O. (1973a): New plants in Cuba II. *Acta Bot. Acad. Sci. Hung.* **18**: 29–48.
- Borhidi, A. and Muñiz, O. (1973b): Novedades florísticas en la flora de Cuba I. *Bot. Közlem.* **59**: 187–192.
- Borhidi, A. and Muñiz, O. (1975): Combinationes novae florae Cubanae II. *Bot. Közlem.* **62**: 25–27.
- Borhidi, A. and Muñiz, O. (1976): New plants in Cuba IV. *Acta Bot. Acad. Sci. Hung.* **21**: 221–230.
- Borhidi, A. and Muñiz, O. (1977a): Myrtaceae novae Cubanae I. *Bot. Közlem.* **64**: 13–22.
- Borhidi, A. and Muñiz, O. (1977b): Plantas nuevas en la flora de Cuba V. *Acta Bot. Acad. Sci. Hung.* **22**:
- Borhidi, A. and Muñiz, O. (1977c): Adiciones al conocimiento de la flora cubana (I) *Harpalyce*, *Banara*, *Xylosma*. *Cien. Biol. Acad. Cien. Cuba* **1**: 131–145.
- Borhidi, A. and Muñiz, O. (1977d): Notas sobre Acatáceas Cubanás I. *Oplonia* y *Elytraria*. *Acta Bot. Acad. Sci. Hung.* **23**: 303–318.
- Borhidi, A. and Muñiz, O. (1977e): Myrtaeae novae Cubanae II. *Bot. Közlem.* **64**: 211–220.
- Borhidi, A. and Muñiz, O. (1977f): Revisión del género *Croton* L. (Euphorbiaceae) en Cuba. *Annal. Mus. Nat. Hist. – Hung.* Budapest **69**: 41–53.
- Borhidi, A. and Muñiz, O. (1978): El género *Jacquinia* L. (Theophrastaceae) en Cuba. *Plant Syst. Evol. (Wien)* **129**: 1–11.
- Borhidi, A. and Muñiz, O. (1983): *Catálogo de plantas cubanas amenazadas o extinguidas*. Academia de Cuba, Havana 85 pp.
- Borhidi, A. and Muñiz, O. (1984): *Mapa de la vegetación potencial de Cuba*. Imp. Juceplan, Havana 1–45 pp.
- Borhidi, A. and Muñiz, O. (1985): Adiciones al catálogo de las palmas de Cuba. *Acta Bot. Hung.* **31**: 225–230.
- Borhidi, A. and Muñiz, O. (1986): Phytogeographic survey of Cuba II. Floristic relations and phytogeographic subdivision. *Acta Bot. Hung.* **32**: 3–48.

- Borhidi, A., Muñiz, O. and Del-Risco, E. (1979): Clasificación fitocenológica de la vegetación de Cuba. *Acta Bot. Acad. Sci. Hung.* **25**: 263—301.
- Borhidi, A., Muñiz, O. and Del-Risco, E. (1983): Plant communities of Cuba I. Fresh- and salt water, swamp and coastal vegetation. *Acta Bot. Hung.* **29**: 337—376.
- Borhidi, A. and Pócs, T. (1985): Effects of serpentine on the distribution of cryptogams in Cuba. Abstr. IAB Conf. of Bryoecology Budapest—Vácrátót, 1985. *Abstr. Bot.* **9**: Suppl. 3: 14.
- Boletín de la Lluvia. Inst. Recurs. Hidraul. Havana. 1966—1970.
- Bouillettenne, R. (1926): Savanes équatoriales de l'Amérique de Sud. *Soc. Roy. Bot. Belg. Bull.* **58**: 217—223.
- Bouillettenne, R. (1930): Un voyage botanique dans le Bas-Amazone. *Mission Biol. Belg. au Brésil* **2**: 1—185.
- Britton, N. L. (1971): The natural vegetation of the Isle of Pines. *New York Bot. Gard.* **17**: 64—71.
- Brooks, R. R., Lee, J. and Jaffré, T. (1974): Some New Zealand and New Caledonia plant accumulators of nickel. *J. Ecol.* **62**: 493—499.
- Brown, W. H. (1919): Vegetation of the Philippine Islands. *Philippine Bureau of Science, Dept. Agric. Nat. Res. (Manila)* **13**: 1—434.
- Budowski, G. (1956a): Regeneration systems in tropical American lowlands. *Caribb. Forester* **17**: 75—90.
- Budowski, G. (1956b): Tropical savannas, a sequence of forest felling and repeated burnings. *Turrialba* **6**: 23—33.
- Budowski, G. (1965): Distribution of tropical American rainforest species in the light of successional processes. *Turrialba* **15**: 40—42.
- Budowski, G. (1970): The distinction between old secondary and climax species in tropical Central American lowland forests. *Trop. Ecol.* **11**: 44—48.
- Budyko, M. T. (1956): *The Heat Balance of the Surface*. Gidromet. Izd. 255 pp.
- Butt, B. D. (1942): Some East African vegetation communities. *J. Ecol.* **30**: 65—146.
- Busse, W. (1908): Die periodischen Grassbrände im tropischen Afrika, Ihr Einfluss auf die Vegetation und ihre Bedeutung für die Landeskultur. *Mitt. Deutsch. Schutzgebieten* **21**: (2) Berlin.
- Butterlin, J. (1956): *La constitution géologique et la structure des Antilles* Paris, Centre Nat. Réch. Scient. 453 pp.
- Cabrer-Mestre, P. (1964): Introducción al estudio de las tierras aridas en la América Latina. *Agroecn. Cuba* No. 1—2.
- Cain, S. A. (1939): The Climax and its complexities. *Amer. Midland. Nat.* **21**: 146—181.
- Cain, S. A. (1944): *Foundations of Plant Geography*. Harper and Boss. NY. 556 pp.
- Cain, S. A. and Castro, G. M. de (1959): *Manual of Vegetation Analysis*. Harper and Brothers New York 325 pp.
- Capote, R. P. and Berazain, R. (1984): Clasificación de las formaciones vegetales de Cuba. *Rev. Jard. Bot. Nac. Cuba* **5(2)**: 1—48.
- Capote, R. P., Menéndez, L., Vilamajó, D. and Borhidi, A. (1985): Nuevas asociaciones de bosque tropical en la Reserva de la biosfera, Sierra del Rosario, Pinar del Río, Cuba. — *Mem. Primer Simp. Bot. Cuba* **3**: 190—211.
- Capot-Rey, R. (1951): Une carte de l'indice d'aridité au Sahara Français. *Bull. de l'Assoc. Géogr. Franc.* (1951) 73—76.
- Carabia, J. P. (1937): Distribution of Cacti in Cuba. *Cactus and Succ. J.* **8**: 202—204.
- Carabia, J. P. (1940): Brief review of the Cuban flora. *Mem. Soc. Cub. Hist. Nat.* **14**: (1): 45—49.
- Carabia, J. P. (1941): Contribuciones al estudio de la flora Cubana. Gymnospermae. *Caribb. Forester* **2**: 83—99.
- Carabia, J. P. (1943): Endemism in the Cuban flora. *Suppl. Amer. J. Bot.* **29**(10): 4.
- Carabia, J. P. (1945a): A brief review of the Cuban Flora. In: Verdoorn, F.: *Plant and Plant Science in Latin America. Chronica Bot. Co.* 68—70.
- Carabia, J. P. (1945b): Vegetation of Sierra de Nipe, Cuba. *Ecol. Monogr.* **15**: 324—341.
- Carreras, R. and Vales, M. A. (1988): Estudio comparativo de la madera de Bucida L. (Combretaceae). *Acta Bot. Hung.* **32**: 247—253.
- Catasus, L. J. G. (1980): Nuevas especies de gramíneas para Cuba. *Acta Bot. Cubana* **4**: 1—11.

- Chapman, J. D. and White, F. (1970): *The Evergreen Forests of Malawi*. Common. For. Inst. Univ. of Oxford. 189 pp.
- Charter, C. G. (1941): *Reconnaissance Survey of the Soil of British Honduras*. Trinidad Govt Printer.
- Charter, J. R. and Keay, R. W. J. (1960): Assessment of the Olokemeji fire control experiment 28 years after institution. *Niger. For. Inf. Bull.* No. 3.
- Cheng-Chiang, Chen (1957): Climatic classification and agricultural regions in Taiwan. *Proc. IX. Pac. Sci. Congr.* Bangkok. 20 pp.
- Chrysler, M. A. (1948): Origin of the West Indian fern flora. *Amer. Fern J.* **38**: 176–186.
- Ciferri, R. (1936): Studio geobotanico dell'Isola Hispaniola (Antille). *Atti Ist. Bot. Univ. Pavia* **8**: 3–336.
- Clark, J. D. (1959): *The Prehistory of Southern Africa*. Penguin Books. London
- Cole, M. M. (1960): Cerrado, Caatinga and Pantanal: the distribution and origin of the savanna-vegetation of Brazil. *Geogr. J.* **126**(2): 168–179.
- Cole, M. M. (1963): Vegetation and geomorphology in Northern Rhodesia: An aspect of the distribution of the savanna of Central Africa. *Geogr. J.* **129**(3): 290–310.
- Cook, M. T. and Gleason, H. A. (1928): *Ecological Survey of the flora of Porto Rico*. *J. Dept. Agricult. Porto Rico*. 139 pp.
- Corner, E. J. H. (1919): The durian theory or the origin of the modern tree. *Ann. Bot. n.s.* **13**: 367–414.
- Corner, E. J. H. (1964): *The Life of Plants*. Weidenfeld and Nicholson London 319 pp.
- Corner, E. J. H. (1966): *The Natural History of Palms*. Weidenfeld and Nicholson London.
- Corral, J. I. (1939): La union de Cuba con el continente americano. *Rev. Soc. Cub. Ing.* **33**: 582–681.
- Corral, J. I. (1940): El geosinclinal cubano. *Rev. Soc. Cub. Ing.* **34**: 485–623.
- Crooke, W. M. (1956): Effect of soil reaction on uptake of nickel from a serpentine soil. *Soil Sci.* **81**: 269–276.
- Crooke, W. M. and Inkson, R. H. E. (1955): The relationship between nickel toxicity and major nutrient supply. *Plant and Soil*. **6**: 1–15.
- Curtis, J. T. (1947): The palo verde forest type near Gonaives, Haiti, and its relation to the surrounding vegetation. *Caribb. Forester* **8**: 1–26.
- Dahlgren, B. E. and Glassman, S. F. (1963): A revision of the genus *Copernicia*. *Gent Herb.* **9**(2): 1–232.
- Dansereau, P. (1951): Description and recording of vegetation upon a structural basis. *Ecology* **32**: 172–229.
- Dansereau, P. (1957): *Biogeography. An Ecological Perspective*. Ronald Press, New York. 902 pp.
- Dansereau, P. (1958): An universal system for recording vegetation. *Contr. Inst. Bot. Univ. Montreal*, No. 72. 1–52.
- Dansereau, P. (1966): Studies on the vegetation of Puerto Rico, I. *Univ. of Puerto Rico Spec. Publ.* No. 1, 1–45.
- Dansereau, P. and Buell, F. P. (1966): Studies on the vegetation of Puerto Rico II. *Univ. of Puerto Rico Spec. Publ.* No. 1. 46–287.
- Dansereau, P. and Lems, K. (1957): The grading of dispersal types in plant communities and their ecological significance. *Contr. Inst. Bot. Univ. Montreal* **71**: 52 pp.
- Daubenmire, R. (1968): Ecology of fire in grasslands. *Adv. Ecol. Res.* **5**: 209–266.
- Davies, J. A. (1965): Estimation of insolation for West Africa. *Q. Journ. Roy. Met. Soc.* **91**: 359–363.
- Davies, J. A. (1966): The assesment of evapotranspiration for Nigeria. *Geogr. Annual.* **48/A**: 139–156.
- Davies, J. A. and Robinson, P. J. (1969): A simple energy balance approach to the moisture balance climatology of Africa. In: Thomas, M. F. and Whittington, G. V. (eds.) *Environment and Land Use in Africa*. London, Methuen and Co. Ltd. 25–56.
- Davis, J. H. (1940): The ecology and geologic role of mangrove in Florida. *Carnegie Inst. Washingt. Publ.* **517**: 303–412.
- Davis, J. H. (1942): The ecology of the vegetation and topography of the sand keys of Florida. *Carnegie Inst. Washingt. Publ.* **524**: 115–195.
- Davis, T. A. W. (1941): On the island origin of the endemic trees of the British Guiana Peneplain. *J. Ecol.* **29**: 1–13.
- Davis, T. A. W. and Richards, P. W. (1933–1934): The Vegetation of Morabilli Creek, British Guiana: An ecological study of a limited area of tropical rain forest. Part I. *J. Ecol.* **21**: 350–381. Part II. *J. Ecol.* **22**: 106–155.

- Dawkins, H. C. (1961): New methods of improving stand composition in tropical forest. *Caribb. Forester* **22**: 12–20.
- Deacon, E. L., Priestly, C. H. B. and Swinbank, W. C. (1958): Evaporation and the water balance. In: Arid zone research. X. Climatology. *Rev. Res. UNESCO*. 9–34.
- Del-Risco, E. (1975): Situacion actual de la mapificación de la vegetación de Cuba. Abstr. XII. Intern. Bot. Congr. Leningrad. **I**: 165.
- Del-Risco, E. and Borhidi, A. (1975): Estructura del manto vegetal. In: *Informe sobre el estado de los conocimientos relativos a los ecosistemas forestales tropicales de Cuba; bosques perennifolios, semicaducífolios*. Informe al UNESCO, 23–43.
- Del-Risco, E. and Borhidi, A. (1976): Sovremennoe sostoyanie chartographirovania rastitelnosti Kubi. *Geobot. Kartogr.* (Leningrad) (1976) 58–59.
- Del-Risco, E. and Borhidi, A. (1977): El area de Zapata, Cuba I. Clasificación fitocenológica y descripción de la vegetación. — Mscr. 48 pp.
- Dengo, G. (1969): Problems of tectonic relations between Central America and the Caribbean. *Gulf Coast Assoc. Geol. Soc. Trans.*, **19**: 311–320.
- Dietz, R. and Holden, L. (1970): Reconstruction of Pangea: Breakup and dispersion of continents, Permian to present. *Geophys. Res.* **75**: No. 26., 4939–4956.
- Donselaar, J. van (1965): An ecological and phytogeographic study of northern Surinam savannas. *Wentia* **14**: 1–163.
- Drew, A. and Reilly, C. (1972): Observation on copper-tolerance in the vegetation of a Zambian copper clearing. *J. Ecol.* **60**: 439–444.
- Duarte, P. P. B. (1982): Distribución de los musgos en las regiones fitogeográficas de Cuba. *Acta. Bot. Cub.* No. 7. 1–20.
- Duek, J. J. (1975): Contribution of the flora of Cuba: Osmundaceae, Schizaeaceae and Gleicheniaceae (Pteridophyta). *Feddes Repert.* **87**: 325–360.
- Duvigneaud, P. (1959): Plantes "cobaltophytes" dans le Haut Katanga. Études sur la vegetations du Katanga et de ses sols metallifères, 2. *Bull. Soc. Roy. Bot. Belg.* **91**: 111–134.
- Duvigneaud, P. and Plancke, J. (1959): Les "Acrocephalus" arborescetes des plateaux Katangais. Études sur la végétation du Katanga et de ses sols métallifères, 5. *Biol. Jaarb. Gand.* **27**: 214–257.
- Duvigneaud, P. and Timperman, J. (1959): Études sur le genre *Crotalaria*. Études sur la vegetation du Katanga et de ses sols métallifères 3. *Bull. Soc. Roy. Bot. Belg.* **91**: 135–176.
- Eardley, A. J. (1954): Tectonic relations of North and South America. Am. Assoc. Petroleum South America. *Amer. Assoc. Petr. Geol. Bull.* **38**.
- Eden, M. J. (1964): The savanna ecosystem — Northern Rupununi, British Guiana, *McGill Univ. Savanna Res. Ser.* **1**: 1–216.
- Eggeling, W. J. (1947): Observations on the ecology of Budongo rainforest, Uganda *J. Ecol.* **34**: 20–87.
- Eiten, G. (1963): Habitat, Flora of Fazenda Campinha, Sao Paulo, Brasil. Univ. Sao Paulo. *Simpos. Sobre o Cerrado*. 181–231.
- Eiten, G. (1968): Vegetation forms. *Bol. de Inst. Bot.* **4**: 1–88.
- Eiten, G. (1972): The cerrado vegetation of Brasil. *Bot. Review* **38**: 201–341.
- Ellenberg, H. and Mueller-Dombois, D. (1966a): Tentative physiognomic ecological classification of plant formations of the Earth. *Ber. Geobot. Inst. Rübel* **37**: 21–56.
- Ellenberg, H. and Mueller-Dombois, D. (1966b): A key to Raunkiaer's plant life forms with revised subdivisions. *Ber. Geobot. Inst. Rübel* **37**: 56–73.
- Emberger, L. (1930): Sur une formule climatique applicable en géographie botanique. *C. R. Acad. Sci.* **191**: 389–391.
- Emberger, L. (1955): Une classification biogéographique des climats. *Rec. Trav. l'Univ. Montpellier, Sér. Bot.* **7**: 3–43.
- Franshave, D. B. (1954): Forest types of British Guiana. *Caribb. Forester* **15**: 73–111.
- Franshave, D. B. (1971): The vegetation of Zambia. *Forest. Res. Bull.* No. 7, 1–67. Lusaka. Governm. Printer.
- Fedorov, A. A. (1966): The structure of the tropical rainforest and speciation in the humid tropics. *J. Ecol.* **54**: 1–11.
- Fekete, G. and Szukló-Lacza, J. (1971): A survey of the plant lifeform systems and the respective research approaches III. *Ann. Mus. Nat. Hist.—Nat. Hung.* **63**: 37–50.

- Fekete, J. (1988): *Trópusi Talajok* (Tropical Soils). In Hungarian. Akadémiai Kiadó, Budapest. 510 pp.
- Finko, V. I., Korin, I. Z. and Formell, C. F. (1967): Sobre la edad de la corteza del intemperismo y las lateritas de Cuba. *Rev. Geol. Acad. Cien. Cuba* **1**(1): 29–47.
- Finol, U. H. (1971): Nuevos parámetros a considerarse en el análisis estructural de las selvas vírgenes tropicales. *Rev. For. Venez.* **21**: 29–42.
- Fors, J. A. (1941): Informe sobre plantaciones forestales en Cuba. *Caribb. Forester* **2**: 138–140.
- Fors, J. A. (1942): La Guana (*Sterculia cubensis* Urb.). *Rev. Agr.* **26**(20): 106–115.
- Fors, J. A. (1947): El Pino macho, *Pinus caribaea*, en las lomas de Trinidad, Cuba. *Caribb. Forester* **8**: 125–134.
- Forster, A. S. and Rodriguez, M. S. P. (1942): Field studies on *Microcycas calocoma*. *Mem. Soc. Cub. Hist. Nat.* **16**(2): 105–121.
- Fosberg, F. R. (1963): The Island Ecosystem. In: Fosberg, F. R. (ed.) *Man's place in the Island Ecosystem*. 1–6. Honolulu; Bishop Museum Press.
- Fraser, H. (1956): Forest preservation in the Windward Islands. *Caribb. Forester* **17**: 25–27.
- Fuentes, V. (1982): El género *Espadaea* A. Rich. *Rev. Jard. Bot. Nac. Cub.* **3**(3): 51–67.
- Furazzola-Bermudez, G., Judoley, C. M., Mikkailovskaya, M. S., Mirolubov, J. S. and Solsona, J. B. (1964): *Geología de Cuba*. Cons. Nac. de Cuba. La Habana: 239 pp.
- Gams, H. (1930): Über Reliktföhrenwälder und das Dolomitphänomen. *Bes. Geobot. Inst. Rübel* **6**: 32–80.
- Gantcheff, S. (1968): Kratka geobotanicheskaya karakteristika na rastitel'nata pokrivka v rezervata Kupejal v Kuba. *Izv. Bot. Inst. Bulg. Akad. Nauk.* **18**: 17–59.
- Gauch, H. G. Jr. and Whittaker, R. H. (1972): Comparison of ordination techniques. *Ecol.* **53**: 868–875.
- Gaussen, H. (1954a): Expression du milieu par des formules écologiques. Leur représentation cartographique. Les division écologiques du monde. *Coll. Intern. CNRS. (Paris) Ann. Biol.* **31**(5–6): 257–267.
- Gaussen, H. (1954b): Les possibilités forestières du monde en relation avec climat. *Act. du IV<sup>e</sup> Congr. Forest. Mond.* **3**. (Dehra Dun) 125–130.
- Gaussen, H. (1954c): Théories et classification des climats et microclimats. — *8<sup>me</sup> Congr. de Bot. Paris* Sect. 3–7.
- Gaussen, H. (1955a): Détermination des climats par la méthode des courbes ombrothermiques. *C. R. Acad. Sci.* **240**: 642–644.
- Gaussen, H. (1955b): Les climats analogues à l'échelle du monde. *R. C. Acad. Sci. Agr. Paris* **41**.
- Gaussen, H. (1963): Les cartes bioclimatiques et de la végétation. Principes directeurs et emploi de la couleur. *Science du Sol*. 1963. Mai. No. 1. 1–14.
- Gaussen, H. (1968a): Relations des sols et des climats avec la végétation et les paysages agraires. *Mém. Doc. Centre Réch. Cart. Géogr.* **6**: 35–39.
- Gaussen, H. (1968b): Carte de végétation comme base pour l'utilisation du sol et la production d'aliments. Manuscript 1–18.
- Gaussen, H. and Bagnouls, F. (1952): L'indice xérothermique. *Bull. de l'Assoc. Géogr.* (1952) 10–16.
- Gentilli, J. (1953): Une critique de la méthode de Thornthwaite pour la classification des climats. *Ann. de Géogr.* **351**: 180–185.
- Gentry, A. H. (1969): A comparison of some leaf characteristics of tropical dry forest and tropical wet forest in Costa Rica. *Turrialba* **19**: 419–428.
- Gigon, A. and Rorison, I. H. (1972): The response of some ecologically distinct plant species to nitrate and to ammonium nitrogen. *J. Ecol.* **60**: 93–102.
- Gill, T. (1931): *Tropical Forests of the Caribbean*. Washington, 318 pp.
- Gleason, H. A. and Cook, T. M. (1926): Plant ecology of Puerto Rico. **7**: 1–178. In: Scientific Survey of Puerto Rico and the Virgin Islands. Washington.
- Glover, J., Robinson, R. and Henderson, J. (1954): Provisional maps of the reliability of annual rainfall in East-Africa. *Q. Journ. Roy. Meteorol. Soc.* **80**: 602–609.
- Glover, P. E. (1968): The role of fire and other influences on a savanna habitat, with suggestions for further research. *East. Afr. Wildl. J.* **6**: 131–137.
- Göhlert, F. (1928): Die Flora über Eisenkarbonat. *Biol. Generalis (Wien)* **4**: 333–336.
- Goldstein, R. A. and Grigal, D. F. (1972a): *Computer Programs for the Ordination and Classification of Ecosystems*. ORNL, Oak Ridge.
- Goldstein, R. A. and Grigal, D. F. (1972/b): Definition of vegetation structure by canonical analysis. *J. Ecol.* **60**: 277–284.

- Golley, F. B., McGuinis, J. T., Clements, R. G., Child, G. I. and Duever, M. J. (1969): The structure of tropical forests in Panama and Colombia. *Bio Science* **19**: 693–696.
- Golley, F., Odum, H. T. and Wilson, R. F. (1962): The structure and metabolism of a Puerto Rican red mangrove forest in May. *Ecology* **43**: 10–19.
- Gómez de la Maza, M. and Roig, J. T. (1914): *La Flora de Cuba*. Rambla, Bonza y Margall, Habana. 182 pp.
- Gómez-Pompa, A. (1967): Some problems of tropical plant-ecology. *J. Arn. Arb.* **48**: 105–121.
- Good, R. (1954): *The Geography of the Flowering Plants*. 2nd ed Longmans Green and Co., London--New York. 452 pp.
- Goodland, R. (1970): The savanna controversy. *McGill Univ. Savanna Res. Ser.* **15**: 1–66.
- Goreczynski, W. (1943): Aridity factor and precipitation ratio and their relation to world climats. *Bull. Pol. Inst. Arts and Science* **1**, New York (3).
- Gorden, A. and Lipman, C. B. (1926): Why are serpentine and other magnesian soils unfertile? *Soil. Sci.* **22**: 291–302.
- Gradman, R. (1916): Wüste und Steppe. *Geogr. Zschr.* **22**: 503.
- Gradstein, R., Pócs, T. and Vana, J. (1983): Disjunct hepaticae in tropical America and Africa. *Acta Bot. Hung.* **29**: 127–171.
- Graham, A. (1974): *Vegetation and Vegetation History of Northern Latin America*. Elsevier, New York.
- Gregory, R. P. G. and Bradshaw, A. D. (1965): Heavy metal tolerance in populations of *Agrostis tenuis* Sibth. and other grasses. *New Phytol.* **64**: 131–143.
- Greig-Smith, P. (1952): Ecological observations on degraded and secondary forest in Trinidad, British West Indies. II. Structure of the communities. *J. Ecol.* **40**: 316–330.
- Greig-Smith, P. (1971): Application of numerical methods to tropical forests. In: Patil, G. P., Pielou, E. C. and Waters, W. E. (eds): *Statistical Ecology*. Vol. 3. *Many species populations, ecosystems and system analysis*. The Pennsylvania State Univ. Press, University Park. 195–204.
- Greig-Smith, P., Austin, M. P. and Whitmore, T. C. (1967): The application of quantitative methods to vegetation survey I. Association analysis and principal component ordination of rain forest. *J. Ecol.* **55**: 483–503.
- Griffiths, J. F. (1962): The climate of East Africa. In: *The Natural Resources of East Africa*. Russel, Nairobi. 144 pp.
- Grisebach, A. R. (1857): Über die Vegetation der Kariben. *Abh. Königl. Ges. Wiss. Göttingen* **7**: 151–286.
- Grisebach, A. R. (1860): Plantae Wrightianae e Cuba Orientali. Pars I. *Mem. Amer. Acad. Sci. N. S.* **8**: 20–192.
- Grisebach, A. R. (1862): Plantae Wrightianae e Cuba Orientali. Pars II. *Mem. Amer. Acad. Sci. N. S.* **8**: 503–536.
- Grisebach, A. R. (1864): *Flora of the British West Indian Islands*. Lovell Reeve and Co. London. 789 pp.
- Grisebach, A. R. (1866): *Catalogus Plantarum Cubensium*. Lipsiae, 292 pp.
- Grubb, P. J., Lloyd, J. R., Pennington, T. D. and Whitmore, T. C. (1963): A comparison of montane and lowland rainforest in Ecuador I. The forest structure, physiognomy and floristics. *J. Ecol.* **51**: 567–601.
- Grudzinskaya, I. A. (1975): Shrub formation (Fourrés) of Cuba. Abstr. XII. Intern. Bot. Congr., Leningrad. 113.
- Guerra, A. T. (1953): Formação de lateritos sobre floresta equatorial amazônica. *Rev. Brasil. Geogr.* **14**: 407–426.
- Guevara, S. S. and Gómez-Pompa, A. (1972): Seeds from surface soils in a tropical region of Veracruz, Mexico. *J. Arn. Arb.* **53**: 312–335.
- Hadač, E. (1970): Novitates Florae Cubanæ. *Folia Geobot. Phytotax.* **5**: 429–433.
- Hadač, E. (1975): Species diversity of mangrove and continental drift. Abstr. Xllth Intern. Bot. Congr., Leningrad. 113.
- Hadač, E. and Hadačová, V. (1969): Notes on the ecology and distribution of *Bidens pilosa* L. in Cuba. *Folia Geobot. Phytotax.* **4**: 165–173.
- Hadač, E. and Hadačová, V. (1971): The Association *Blechno serrulati-Aceoloraphetum wrightii* in the Remates de Guane, West-Cuba, and its ecology. *Folia Geobot. Phytotax.* **6**: 369–388.
- Hammen, Th. van der. (1961): The quaternary climatic changes of Northern South America. *Ann. New York Acad. Sci.* **95**: 676–683.

- Hammen, Th. van der. (1972): Historia de la vegetación y el medio ambiente del Norte Sudamericano. *Mem. I. Congr. Latinoam. Bot.* (Mexico) 119–134.
- Hämet-Ahti, L., Ahti, T. and Koponen, T. (1974): A scheme of vegetation zones for Japan and adjacent regions. *Ann. Bot. Fennici* **11**: 59–88.
- Hardy, F. (1940): A provisional classification of the soils of Trinidad. *Trop. Agr.* (Trinidad) **17**: 153–158.
- Hardy, F. (1945): The soils of South America. In: Verdoorn F.: *Plants and Plant Science in Latin America*. Vol. 2. 322–326 pp.
- Harman, H. H. (1967): *Modern Factor Analysis*. Wiley, New York.
- Harshberger, J. M. (1958): *Phytogeographic Survey of North America* (2nd ed.) Weinheim, Engelmann; 790 pp.
- Hatheway, W. H. (1971): Contingency-table analysis of rainforest vegetation. In: Patil, G. P., Pilou, E. C. and Waters, W. E. (eds): *Statistical Ecology*, Vol. 3. *Many Species Population, Ecosystems and Systems Analysis*, The Pennsylvania State Univ. Press, University Park, Pa. 207–307 pp.
- Hattori, S. (1955): Hepaticae occurring on serpentine on Mt Apoi (Hokkaido). *Bot. Magazine*, Tokyo, **68**: 320–323.
- Hawkes, J. G. and Smith, P. (1965): Continental drift and the age of *Angiosperm* genera. *Nature* **207**, No. 4992, 48–50.
- Hayek, F. (1926): *Allgemeine Pflanzengeographie* Bornträger. Berlin.
- Hernandez, A. (1966): Importancia del estudio genetico de los suelos: su aplicación en la prov. de Las Villas. *Tech. Agropec.* No. 2.
- Hess, H. H. and Maxwell, J. C. (1953): *Caribbean Research Project*. **5**: 707–773.
- Hewetson, C. E. (1956): A discussion on the climax concept in relation to the tropical rain and deciduous forest. *Emp. For. Rev.* **35**: 274–291.
- Hills, T. L. (1965): Savannas: A review of a major research problem in tropical geography. *Canad. Geographer* **9**: 216–228.
- Hills, T. L. (1969): The savanna landscapes of the Amazon basin. *McGill Univ. Savanna Res. Ser.* **14**: 1–41.
- Hills, T. L. and Randall, R. E. (eds) (1968): The ecology of the forest-savanna boundary. *McGill Univ. Savanna Res. Ser.* **13**: 1–128.
- Hirt, P. (1926): Die Isonotiden. *Peterm. Geogr. Mitteil.* **72**.
- Hitchcock, A. S. (1936): *Manual of the Grasses of the West Indies*. Washington, U. S. Govt. Print. Off. 439 pp.
- Holdridge, L. R. (1940): Some notes on the mangrove swamps of Puerto Rico. *Caribb. Forester* **1**(4), 19–29.
- Holdridge, L. R. (1962): The determination of atmospheric water movements. *Ecology* **43**: 1–9.
- Holdridge, L. R. (1967): *Life Zone Ecology*. San José. 206 pp.
- Holdridge, L. R. and Budowski, G. (1956): Report on an ecological survey of the Republic of Panama. *Caribb. Forester* **17**: 92–110.
- Holdridge, L. R. and Budowski, G. (1957): Informe sobre un levantamiento ecológico de Panamá. *Caribb. Forester* **18**: 12–32.
- Holdridge, L. R., Gronke, W. C., Hatheway, W. H., Liang, T. and Tosi, J. A., Jr. (1971): *Forest environments in tropical life zones. A pilot study*. Wiley and Son, New York 747 pp.
- Hollick, A. (1924): A paleobotanical survey of the Antillean region. *Bull. Torr. Bot. Club* **51**: 19–34.
- Hopkins, B. (1956a): Observation on savanna burning in the Olokemeji Forest Reserve, Nigeria. *J. Appl. Ecol.* **2**: 367–381.
- Hopkins, B. (1956b): *Forest and Savanna*. Heinemann, Ibadan and London, 100 pp.
- Horn, E. F. (1947): The grasslands, savanna forests and dry forests of Brazil. *Caribb. Forester* **8**: 135–144.
- Horn, E. F. (1948): The Amazon Hylea, some notes on its development. *Caribb. Forester* **8**: 361–372.
- Howard, A. R. (1947): Notes on some plants of Cuba. *J. Arn. Arb.* **28**: 117–126.
- Howard, A. R. (1949): Studies on the genus *Coccoloba*. The genus *Coccoloba* in Cuba. *J. Arn. Arb.* **30**: 388–424.
- Howard, A. R. (1968): The ecology of an elfin forest in Puerto Rico. I. Introduction and composition studies. *J. Arn. Arb.* **49**: 381–418.
- Howard, A. R. (1969): The ecology of an elfin forest in Puerto Rico. VIII. Studies of stem growth and form and of leaf structure. *J. Arn. Arb.* **50**: 225–262.

- Howard, A. R. (1974): The vegetation of the Antilles. In: Graham, A. (ed.): *Vegetation and Vegetation History of Northern Latin America*. Elsevier, New York. 1–38.
- Howard, A. R. (1975): Modern problems of the years 1492–1800 in the Lesser Antilles. *Ann. Mo. Bot. Gard.* **62**: 368–379.
- Howard, A. R. (1979): Flora of the West Indies. In: Larsen, K. (ed.): *Tropical Botany*. Academic Press, London. 239–250.
- Howard, A. R. (1979): Flora of the Lesser Antilles (Leeward and Windward Islands). Vol. 3. Monocotyledoneae. Brooke Thompson Mills (ed). Arnold Arboretum Harvard Univ. Cambridge 586 pp.
- Howard, A. R. and Briggs, W. R. (1953): The vegetation on coastal dogtooth limestone in Southern Cuba. *J. Arn. Arb.* **34**: 88–95.
- Hueck, K. (1966): *Die Wälder Südamerikas*. Fischer, Jena 422 pp.
- Hueck, K. and Seibert, P. (1972): *Vegetationskarte von Südamerika*. Fischer, Stuttgart 69 pp.
- Hunter, J. G. and Vergnano, O. (1952): Nickel toxicity in plants. *Ann. Appl. Biol.* **39**: 279–284.
- Hunziker, A. T. (1979): South American Solanaceae; a synoptic survey the biology and taxonomy of the Solanaceae. *Linnean Soc. Symp. Ser.* No. 7.
- Ijzerman, R. (1931): *Outline of the Geology and Petrology of Surinam*. Kemink. Utrecht.
- Innes, R. R. (1971): Fire in West African Vegetation. *Proc. Ann. Tall Timbers Fire Ecology Conference*. **11**: 147–173.
- Isachs, B., Oliver, J. and Sykes, L. R. (1968): Seismology and the new global tectonics. *J. Geophys. Res.* **73**: No. 18., 5855–5899.
- Iturralde, M. A. (1975): Problemas de aplicación de dos hipótesis tectónicas modernas a Cuba y la región Caribe. *Rev. Technol.* **13**: 46–63.
- Iturralde-Vinent, M. A. (1977): Los movimientos tectónicos de la etapa de desarrollo platafórmico en Cuba. *Inf. Cien.-Tec. Acad. Cien. Cuba*. No. 20. 24. pp.
- Jacobsen, W. B. G. (1967): The influence of the copper content of the soil on the vegetation at Silverside North, Mangula area. *Kirkia* **5**: 259–277.
- Jacobsen, W. B. G. (1968): The influence of the copper content of the soil on trees and shrubs at Molly South Hills, Mangula. *Kirkia* **6**: 63–84.
- Jaeger, F. (1945): Zur Gliederung und Benennung des tropischen Grasslandgürtels. *Verh. Naturf. Ges. (Basel)* **56**: 509–520.
- Janzen, D. H. (1970): Promising directions of studies in tropical animal-plant interactions. *Ann. Mo. Bot. Gard.* **64**: 706–736.
- Járai-Komlódi, Magda (1964): Trópusi vegetációtanulmányok Kubában (Studies on tropic vegetation in Cuba). *Bot. Közlem.* **51**: 143–168.
- Jennings, O. E. (1917): A contribution to the botany of the Isle of Pines. *Ann. Carnegie Mus.* **11**: 19–290.
- Johannessen, C. L. (1963): Savannas of interior Honduras. *Ibero-Americana* **46**: 1–160.
- Johnson, C. M., Pearson, G. A. and Stout, P. R. (1952): Molybdenum nutrition of crop plants. II. *Plant and Soil* **4**: 178–196.
- Jones, C. F. (1930): Agricultural regions of South America. *Instalment VIII. Econ. Geogr.* **6**: 1–36.
- Jovet-Ast, S. (1949): Les groupements de Muscinaées épiphytes aux Antilles Françaises. *Rev. Bryol. Lichen.* **18**: 125–146.
- Judd, S. W. (1981): A monograph of *Lyonia* (Ericaceae). *J. Arn. Arb.* **62**: 63–209, 315–436.
- Judoley, C. M. and Furrazola-Bermudez, G. (1967): La posición de Cuba en la estructura geológica de la región del Caribe. *Rev. Technol.* Vol. 5., No. 6., 10–19.
- Judoley, C. M. and Furrazola-Bermudez, G. (1968): *Estratigrafía y Fauna del Jurásico de Cuba*. Havana, Acad. Cienc. Cuba, Inst. Cubano Rec. Min., 126 pp.
- Judoley, C. M. and Furrazola-Bermudez, G. (1971): *Geología del Área del Caribe y de la Costa del Golfo de México*. La Habana, Ministerio de Min. Comb. y Met. Dpto. Geología, 286 pp.
- Kapos, V. and Tanner, E. V. J. (1985): Water relations of Jamaican upper montane rainforest trees. *Ecology* **66**: 241–250.
- Keay, R. W. J. (1959): *An Outline of Nigerian Vegetation*. Govern. Printer, Lagos.
- Kennan, T. C. D. (1971): The effects of fire on two vegetation types at Matopos, Rhodesia. *Proc. Ann. Tall Timbers Fire Ecology Conference*. No. **11**: 53–98.

- Kershaw, K. A. (1968): Classification and ordination of Nigerian savanna vegetation. *J. Ecol.* **56**: 467–482.
- Khudoley, K. M. and Meyerhoff, A. A. (1971): Paleogeography and geologic history of Greater Antilles. *Geol. Soc. Amer. Mem.* **129**: 200 pp.
- Kinloch, J. B. (1940): Mapping vegetational types in British Honduras from aerial photographs. *Caribb. Forester* **1**: 1–4.
- Kissin, I., Semjonov, E. and Izquierdo, A. (1968): *La Evaporación en Cuba*. Inst. Nac. Recurs. Hidraul. La Habana. 93 pp.
- Kitamura, S. (1950): Adaptation and isolation on the serpentine areas. *Acta Phytotax. Geobot.* **12**: 178–185.
- Klausing, O. (1958): Untersuchungen über Vegetation und Wasserhaushalt am Volcan de San Salvador. *Ber. Deutsch. Bot. Ges.* **71**: 439–452.
- Klimes-Szmik, A., Szébenyi-Nagy, E. and Trémol, A. (1975): *Suelos de Cuba*. Vol. I. Química del suelo. Dir. Nac. Suel. Fertil. INRA. Orbe, Havana. 352 pp.
- Klimes-Szmik, A., Suárez, O. D., Mesa A. N. and Pena J. (1980): *Suelos de Cuba*. Vol. II. Física del suelo. Orbe, Ciudad de la Habana. 328 pp.
- Klinge, H. (1965): Podzol soils in the Amazon basin. *J. Soil. Sci.* **16**: 95–103.
- Klinge, H., Rodrigues, W. A., Brunig, E. and Fittbau, J. (1975): Biomass and structure in a Central Amazonian rain forest. In: Golley, F. B. and Medina, E. (eds): *Tropical Ecological Systems. Ecological Studies* **11**: 115–122.
- Klotz, G. (1978): Arcaltypen cubanischer Pflanzensippen I. *Beitr. Phytotax.* **6**: 7–78.
- Knapp, R. (1964): Höhere Vegetations-Einheiten einiger Gebiete der holarktischen und neotropischen Floren-Reiche. *Geobot. Mitteil.* **28**: 1–11.
- Knapp, R. (1965): *Die Vegetation von Nord- und Mittelamerika*. Fischer, Stuttgart-Jena, 373 pp.
- Knapp, R. (1973): *Die Vegetation von Afrika*. Jena, 626 pp.
- Knight, H. D. (1975): A phytosociological analysis of species-rich tropical forest on Barro Colorado Island, Panama. *Ecol. Monogr.* **45**: 259–284.
- Komarek, E. V. Sr. (1971): Lighting and Fire Ecology in Africa. *Tall Timbers Fire Ecol. Conf.* **11**: 473–511.
- Kool, R. G. A. (1953): *L'agriculture tunisienne*. Wageningen 188 pp.
- Kozáry, M. (1968): Ultrabasic rocks in thrust zones of north-western Oriente province Cuba. *Amer. Assoc. Petroleum Geol. Bull.* Vol. 52: No. 12, 2298–2317.
- Köppen, W. (1900): Versuch einer Klassifikation der Klimate vorzugsweise nach ihrem Beziehungen zur Pflanzenwelt. *Geogr. Zschr.* **6**: 593–611, 657–679.
- Köppen, W. (1918): Klassifikation der Klimate nach Temperatur, Niederschlag, und Jahresverlauf. *Peterm. Geogr. Mitteil.* **64**: 193–203, 243–248.
- Köppen, W. and Geiger, R. (1930–1936): *Handbuch der Klimatologie*. I–V., Berlin.
- Krause, W. (1958a): Pflanzendecke und Standort auf Serpentin. *Handbuch der Pflanzenphysiol.* **4**: 755–798.
- Krause, W. (1958b): Boden und Pflanzengesellschaften. *Handbuch der Pflanzenphysiol.* **4**: 807–850.
- Kruckeberg, A. R. (1954): The ecology of serpentine soils. III. Plant species in relation to serpentine soils. *Ecol.* **35**: 251–258.
- Küchler, A. W. (1947): A geographical system of vegetation. *Geogr. Rev.* **37**: 233–240.
- Küchler, A. W. (1949): A physiognomic classification of vegetation. *Ann. Assoc. Amer. Geogr.* **39**: 20–210.
- Küchler, A. W. (1953): Some uses of vegetation maps. *Ecology* **34**: 629–636.
- Labroue, L., Legris, P. and Viart, M. (1965): *Bioclimats du souscontinent Indien*. Inst. Franc. de Pondichéry **3**: 1–33.
- Lam, H. J. (1927): En plantengeografisch Dorado. *Handg. IV. Nederl. Ind. Naturw. Congr.* 386–397.
- Lamotte, M. (1975): The structure and function of a tropical savanna ecosystem. In: Golley, F. B. and Medina, E. (eds): *Tropical Ecological Systems. Ecol. Studies* **11**: 179–223.
- Lamprecht, H. (1962): Ensayo sobre unos métodos para el análisis estructural de los bosques tropicales. *Acta Cient. Venez.* **13**: 57–65.
- Lang, G. E., Knight, D. H. and Anderson, D. A. (1971): Sampling the density of tree species with quadrats in a species-rich tropical forest. *For. Sci.* **17**: 395–400.
- Lang, R. (1915): Versuch einer exakten Klassifikation der Böden in klimatischer und geologischer Hinsicht. *Internat. Mitteil. Bodenkunde* **5**.

- Lanjouw, J. (1936): Studies on the vegetation of the Surinam savannas and swamps. *Nederl. Kruidk. Arch.* **46**: 823–851.
- Lawley, D. N. and Maxwell, A. E. (1963): *Factor Analysis as a Statistical Method*. Butterworths, London.
- Lemée, G. (1959): Effects des caractères du sol sur la localisation de la végétation en zones équatoriales et tropicales humides. In: *Tropic Soils and Vegetation. Proc. Abidjan Symp.* Paris: UNESCO 1961.
- León, H. Sauget (1923): Una excursion al Pico Turquino. *Rev. de la Salle* **11**:(101): 349–357.
- León, H. Sauget (1926): Cuba: Natural areas and regions. — In: *Naturalist's Guide to the Americas*. Shelford Baltimore 761 pp.
- León, H. Sauget (1931): Contribución al estudio de las palmas de Cuba. *Copernicia I. Rev. Soc. Geogr. Cuba* **3**: 33–59.
- León, H. Sauget (1934): La Guana y su aprovechamiento (*Sterculia cubensis* Urb.) *Mem. Soc. Cuba Hist. Nat.* **2**(8): 63–68.
- León, H. Sauget (1936): Contribución al estudio de las palmas de Cuba. *Copernicia. II. Mem. Soc. Cuba Hist. Nat.* **10**: 203–226.
- León, H. Sauget (1939): Contribución al estudio de las palmas de Cuba III. Género *Coccothrinax*. *Mem. Soc. Cuba Hist. Nat.* **13**: 107–156+lam. 10–21.
- León, H. Sauget (1940): Contribución al estudio de las palmas de Cuba IV. Un corojo cubano nuevo para la ciencia. *Mem. Soc. Cuba Hist. Nat.* **14**: 51–54.
- León, H. Sauget (1941): Contribución al estudio de las palmas de Cuba V. Novedades en el género *Hemitrinax*. *Mem. Soc. Cuba Hist. Nat.* **15**: 378–384.
- León, H. Sauget (1942): Vegetación de las terazas de Maisi. *Rev. Soc. Geogr. Cuba* **15**(2): 33–49.
- León, H. Sauget (1943): Contribution à l'étude de palmiers de Cuba VI. Nouveautés dans les palmiers royaux de Cuba Oriental. *Mem. Soc. Cuba Hist. Nat.* **17**: 1–14.
- León, H. Sauget (1944): Contributions to the study of the Cuban Palms VII. The genus *Calyptrogyne* in Cuba. *Contrib. Ocas. Mus. Hist. Nat. Col. La Salle* No. 3. 1–12.
- León, H. Sauget (1946): Flora de Cuba. Vol. 1. — *Contrib. Ocas. Mus. Hist. Nat. Col. La Salle*, No. 8. 441 pp. Havana.
- León, H. Sauget and Alain H. Liogier (1951–1957): Flora de Cuba Vols. II. III. IV. *Contrib. Ocas. Mus. Hist. Nat. Col. La Salle* Havana. No. 10. 456 pp; No. 13. 502 pp; No. 16. 556 pp.
- Léonard, J. (1952): Les diverses types de forêts du Congo-Belge. *Lejeunea* **16**: 81–93.
- Leroy, J. F. (1979): Composition, origin and affinities of the Madagascan vascular flora. *Ann. Mo. Bot. Gard.* **65**: 535–589.
- Levins, R. (1968): *Evolution in Changing Environments. Some Theoretical Explorations* Princeton Univ. Press, Princeton N. J. 120 pp.
- Lind, E. M. and Morrison M. E. S. (1974): *East African Vegetation*. Longman, London, 257 pp.
- Lindeman, J. C. (1953): *The Vegetation of the Coastal Regions of Suriname*. Utrecht. 416 pp.
- Lipman, C. B. (1926): The bacterial flora of the serpentine soils. cit. In: Bennett, H. H., Allison, R. V.: *The Soils of Cuba*. — Trop. Plant Res. Found. Washington.
- Lippold, H. (1970): Die *Hypericum*-Arten Cubas. *Wiss. Zschr. Schiller Univ. Jena Math. Nat. R.* **9**: 377–382.
- Lippold, H. (1975): Die Apocynaceen-Gattung *Angadenia* auf Cuba. *Wiss. Zschr. Schiller Univ. Jena* **24**: 387–395.
- Lippold, H. (1978a): Die Gattung *Dodonaea* Miller (Sapindaceae) in America. *Beitr. Phytotax. (Jena)* **6**: 79–126.
- Lippold, H. (1978b): Zur Verbreitung der Sapindaceen und einiger Vertreter aus der Familie der Apocynaceen auf Cuba. *Beitr. Phytotax. (Jena)* **6**: 127–169.
- Lippold, H. (1978c): Zur Verbreitung der Sapindaceen und einiger Vertreter der Familie der Apocynaceen auf Cuba. *Beitr. z. Phytotax. (Jena)* **6**: 171–221.
- Lippold, H. (1979): Die Gattung *Plumeria* L. (Apocynaceae) auf Cuba. *Feddes Repert.* **90**: 193–215.
- Longman, K. A. and J. Jenik (1974): *Tropical Forest and its Environment* Longman Group, London. 196 pp.
- Lötschert, W. (1955): La vegetation de El Salvador. *Com. Inst. Trop. Inv. Cien.* **4**: 65–79.
- Lötschert, W. (1958): Die Übereinstimmung von geologischer Unterlagen und Vegetation in der Sierra de los Organos (West-Cuba). *Ber. Deutsch. Bot. Ges.* **71**: 55–70.
- Lötschert, W. (1959): Vegetation und Standortsklima in El Salvador. *Bot. Studien*, **10**: 1–88.

- Loveless, A. R. (1960): The vegetation of Antigua. W. Indies. *J. Ecol.* **48**: 495—527.
- Loveless, A. R. and Asprey G. F. (1957): The dry evergreen formations of Jamaica. I. *J. Ecol.* **45**: 799—822.
- Loveless, C. M. (1959): A study of the vegetation in the Florida everglades. *Ecology* **40**: 1—9.
- Lugo, A. E., Evink, G., Brinson, E. E., Broce, A. and Snedaker, S. C. (1975): Diurnal rates of photosynthesis, respiration and transpiration in mangrove forests of South Florida. In: Golley, F. B. and Medina, E. (eds): *Tropical Ecological Systems. Ecol. Studies* **11**: 335—350.
- Lyon, G. L., Peterson, P. J., Brooks, R. R. and Butler, G. W. (1971): Calcium, magnesium and trace elements in a New Zealand serpentine flora. *J. Ecol.* **59**: 421—429.
- Mabberley, D. J. (1975a): The giant Lobelias: pachycauly, biogeography, ornithophily and continental drift. *New Phytol.* **74**: 365—374.
- Mabberley, D. J. (1975b): The giant lobelias: toxicity, inflorescence and tree building in the Campanulaceas. *New Phytol.* **75**: 289—295.
- MacGillavry, H. J. (1970): Geological history of the Caribbean. *Koninkl. Nederlandse Akad. Wetensch. Proc., Ser. B.*, **73**: 64—96.
- Major, J. (1963): A climatic index to vascular plant activity. *Ecology* **44**: 483—498.
- Malaisse, F. (1973): Contribution à l'étude de l'ecosystème forest claire (Miombo). Note: 8. Le projet Miombo. *Ann. Fac. Sci. Abidjan Ser. Ecol.* **6**: 277—250.
- Malaisse, F., Freson, R., Goffinet, G. and Malaisse-Mousset, M. (1975): Litter fall and litter breakdown in Miombo. In: Golley F. B. and Medina, E. (eds): *Tropical Ecological Systems. Ecol. Studies* **11**: 137—152.
- Malfait, B. and Dinkelman, M. (1972): Circum-Caribbean tectonic and igneous activity and the evolution of the Caribbean plate. *Geol. Soc. Amer. Bull.* **83**: 251—272.
- Mangenot, G. (1951): Sur formule simple permettant de caractériser les climats de l'Afrique intertropical dans leurs rapports avec la végétation. *Rev. Gen. Bot.* **58**: 353—372.
- Margalef, R. (1957): La teoria de la información en la ecología. *Mem. R. Acad. Cienc-Artes Barcelona* **32**: 373—449.
- Margalef, R. (1968): *Perspectives in the Ecological Theory*. University Chicago Press, Chicago.
- Marie-Victorin, F. (1942): *Dracaena cubensis*. *Contr. Inst. Bot. Univ. Montreal*. No. 43: 1—16.
- Marie-Victorin, F. (1943a): Las vallisoletas américaines. *Contr. Inst. Bot. Univ. Montreal* No. 46: 1—38.
- Marie-Victorin, F. (1943b): Les haut pinèdes d'Haiti. *Contr. Inst. Bot. Univ. Montreal* No. 48: 47—60.
- Marie-Victorin, F. (1944): Études taxonomiques sur la flore de Cuba III. L'Acacia serpentinicole de Moa. *Contr. Inst. Bot. Univ. Montreal*. No. 49: 57—67.
- Marie-Victorin, F. (1948a): Le genre *Hedyosmum* (Chloranthacées) avec description d'une espèce nouvelle. *Contr. Inst. Bot. Univ. Montréal*. No. 63: 5—10.
- Marie-Victorin, F. (1948b): Nouvelles études taxonomiques sur la Flore de Cuba III. Le genre *Spathelia* (Rutaceae). *Contr. Inst. Bot. Univ. Montreal*. No. 63: 14—48.
- Marie-Victorin, F. (1948c): Nouvelles études taxonomiques sur la Flore de Cuba IV. Le genre *Purdiaeia* (Cyrillaceae). *Contr. Inst. Bot. Univ. Montreal*. No. 63: 49—62.
- Marie-Victorin, F. and León, H. Saguet (1942, 1944, 1956): Itinéraires botaniques dans l'île de Cuba. *Contr. Inst. Bot. Univ. Montreal*. No. 41: 496 pp.; No. 50: 410 pp.; No. 68: 227 pp.
- Martonne, De E. (1928): Aérisme et l'indice d'aridité. *R. C. Acad. Sci.* **182**: 1395—1398.
- Mason, H. L. (1946): The edaphic factor in narrow endemism. I. The nature of environmental influence. *Madroño* **8**: 209—226.
- Mattson, P. (1973): Middle Cretaceous nappe structures in Puerto Rican ophiolites and their relation to the tectonic history of Greater Antilles. *Geol. Soc. Amer. Bull.* Vol. 84., No. 1. 21—38.
- Maul, F. (1973): Bevezető a trópusi-szubtrópusi talajok ismeretéhez. (Introduction to the knowledge of tropical-subtropical soils). In: Pál, I. (ed.): *Meleggövi növénytani és állattani ismeretek* University Notes. Agricult. Univ. Gödöllő. 16—33.
- Máthé, I. (1942): Magyarország flóraelem-esportjainak életformáösszetétele. (Prozentuelle Verteilung der Lebensformen in den Gruppen der Florenelemente). *Tisia* **5**: 39—43.
- McCulloch, J. S. G. (1963): Tables for the rapid computation of the Penman estimate of evaporation (cyclostyled). East. Afr. Agr. Forest. Res. Organ.
- Medina, E. (1970): Estudios eco-fisiológicos de la vegetación tropical. In: *Ciencia en Venezuela 1970*. Univ. Carabobo, Venezuela. 63—88.
- Meher-Homji, V. M. (1963): Les bioclimats du sub-continent Indien et leurs types analogues dans le monde. *Doc. pour les Cartes des Productions végétales*. Toulouse **1**: 387 pp.

- Melville, R. (1966): Continental drift, Mesozoic continents and the migrations of the Angiosperms. *Nature* **211**: No. 5045, 116–120.
- Mészáros, Z. (1976): The *Melocactus* species of Cuba. *Acta Bot. Acad. Sci. Hung.* **22**: 127–147.
- Mészáros, Z. (1978): Additional contributions to the knowledge of the *Melocactus* species in Cuba. *Acta Bot. Acad. Sci. Hung.* **24**: 301–305.
- Meyer, A. (1926): Über einige Zusammenhänge zwischen Klima und Boden in Europa. *Chemie der Erde* **2**.
- Meyerhoff, A. A. (1967): Future hydrocarbon provinces of Gulf of Mexico-Caribbean region. *Gulf Coast Assoc. Geol. Soc. Trans.*, **17**: 217–260.
- Meyerhoff, A. A. and Meyerhoff, H. A. (1972): Continental drift IV.: The Caribbean “plate”. *J. Geology*. Vol. 80., No. 1, 34–60.
- Meyerhoff, A. A., Meyerhoff, H. A. and Briggs, R. (1972): Continental drift V.: Proposed hypothesis for Earth tectonics. *J. Geology*. Vol. 80., No. 12., 663–692.
- Meyerhoff, A. A. (1954): Antillean tectonics. *New York Acad. Sci. Trans.*, Ser. 2., Vol. 16., No. 3., 149–155.
- Minguzzi, G. and Vergnano, O. (1953): Il contenuto di elementi inorganici delle piante della formazione ophiolitica dell'Impruneta. *Nuova Giorn. Bot. Ital. N. S.* **60**: 287–319.
- Molnár, P. and Sykes, L. R. (1969): Tectonics of the Caribbean and Middle America regions from focal mechanisms and seismicity. *Geol. Soc. Amer. Bull.* Vol. 80, No. 9, 1639–1684.
- Monasterio, M. (1968): Observations sur les rythmes annuels de la savanne des “Llanos” du Venezuela. Thesis, Montpellier, France.
- Monasterio, M. (1971): Ecología de las sabanas de América tropical. I. Análisis macroecológico de los Llanos de Calabozo, Venezuela. *Cuadernos Geogr.* **4**: 1–126.
- Monasterio, M. and Sarmiento, G. (1968): Análisis ecológico y fisiológico de la sabana de la Estación Biológica de Los Llanos. *Bol. Soc. Ven. Cien. Nat.* No. 113–114, 477–524.
- Monasterio, M. and Sarmiento, G. (1971): Ecología de las sabanas de América tropical. II. Caracterización ecológica del clima en los Llanos de Calabozo, Venezuela. *Rev. Geogr.* **21**: 5–38.
- Monod, Th. (1960): Notes botanique sur les îles de Sao Thomé et de Príncipe. *Bull. de l'I. F. A. N.* **22**: Sér. A. No. 1, 19–83 p.
- Montgomery, G. G. and Sungquist, M. E. (1975): Impact of sloths on neotropical forest energy flow and nutrient cycling. In: Golley F. B. and Medina E. (eds) *Tropical Ecological Systems. Ecol. Studies* **11**: 69–98.
- Morgan, W. J. (1968): Rises, trenches, great faults and crustal blocks. *J. Geophys. Res.* Vol. 73, No. 6, 1959–1982.
- Moscoso, R. M. (1943): *Catalogus Florae Domingensis*. Univ. de Santo Domingo, New York, 732 pp.
- Moss, P. P. (1969): The ecological background to land use studies in Tropical Africa, with special reference to the West. In: *Environment and Land Use in Africa*. Thomas and Whittington 193–238.
- Mueller-Dombois, D. (1975): Some aspects of island ecosystem analysis. In: Golley, F. B. and Medina, E. (eds): *Tropical Ecological Systems. Ecological Studies*. **11**: 353–366.
- Müller, C. H. (1955): The origin of *Quercus* of Cuba. *Rev. Soc. Cubana Bot.* **12**(3): 41–48.
- Müller, C. H. (1961): *Quercus oleoides* C. et S. var. *sagraeanas*. *Amer. Midl. Nat.* **65**: 37.
- Muñiz, O. (1970): *Endemismo en la Flora. Atlas Nac. de Cuba* 57–60.
- Muñiz, O. and Borhidi, A. (1981): Palmas nuevas del género *Coccothrinax* Sarg. en Cuba. *Acta Bot. Acad. Sci. Hung.* **27**: 3–4.
- Muñiz, O. and Borhidi, A. (1982): Catálogo de las palmas de Cuba. *Acta Bot. Acad. Sci. Hung.* **28**: 309–345.
- Murça Pires, J., Dobzhansky, Th. and Black, G. A. (1953): An estimate of the number of species of trees in an Amazonian forest community. *Bot. Gaz.* **114**: 467–477.
- Myers, J. G. (1936): Savannah and forest vegetation of the interior Guiana Plateau. *J. Ecol.* **24**: 163–184.
- Nagy, E. (1972): Perfil transversal esquemático de Oriente desde el punto de vista de la tectónica de placas (hipótesis). *Acad. Cienc. Cuba. Inst. Geol. Actas.* No. **2**, 63–65.
- Novák, F. A. (1928): Quelques remarques relatives au problème de la végétation sur les terrains serpentiniques. *Preslia* **6**: 42–71.
- Novák, V. and Pelišek, J. (1940): Rendzinaböden auf Serpentingestein. *Ann. Fac. Agron. Bucarest* **11**.

- Nuñez, J. A. (1959): *Geografía de Cuba*. Havana, (Hungarian edition) 1966. Budapest. Gondolat 221 pp.
- Observatorio Nacional de Cuba (1965): *Boletín de Meteorología y Climatología*, Ep. 5., No. 1. 1–56., No. 2., 1–64.
- Obst, E. (1923): Das abflusslose Rumpfschollenland in nordöstlichen Deutsch-Ostafrika. Teil. II. *Mitt. Geogr. Ges. Hamburg* **35**:
- Odum, E. P. (1959): *Fundamentals of Ecology*. (2 ed.) Saunders New York, 546 pp.
- Odum, E. P. (1962): Relationships between structure and function in the ecosystem. *Jap. J. Ecol.* **12**: 108–118.
- Old, Sylvia, M. (1969): Microclimate, fire and plant production in an Illinois prairie. *Ecol. Monogr.* **39**: 355–384.
- Orlóci, L. (1967): An agglomerative method for classification of plant communities. *J. Ecol.* **55**: 193–206.
- Orlóci, L. and Mukkattu, M. M. (1973): The effect of species number and type of data on the resemblance structure of a phytosociological collection. *J. Ecol.* **61**: 37–46.
- Palmer, R. H. (1945): Outline of the geology of Cuba. *J. Geol.* **53**: 1–34.
- Parkhurst, D. F. and Loucks, O. L. (1972): Optimal leafsize in relation to environment. *J. Ecol.* **60**: 505–538.
- Parsons, J. J. (1955): The Miskito pine-savanna of Nicaragua and Honduras. *Ann. Assoc. Amer. Geogr.* **45**: 36–63.
- Parsons, J. J. (1972): Spread of African pasture grasses to the American tropics. *J. Rge. Mgt.* **25**: 12–17.
- Pelton, W. L., King, K. M. and Tanner, C. B. (1960): An evaluation of the Thornthwaite and mean temperature methods for determining potential evapotranspiration. *Agron. J.* **52**: 387–395.
- Penman, H. L. (1948): Natural evaporation from open water, bare soil and grass. *Proc. Roy. Soc. (Lond.) Biol. A.* **193**: 210–245.
- Penman, H. L. (1956): Evaporation: an introductory survey. *Neth. J. Agric. Sci.* **4**: 8–29.
- Phillips, E. P. (1930): Fire: its influence on biotic communities and physical factors in South and East Africa. *South Afr. J. Sci.* **27**: 352–367.
- Phillips, E. P. (1938): Deterioration in the vegetation of the Union of South Africa and how this may be controlled. *South Afr. J. Sci.* **35**: (Symp. Repr.) 24–32.
- Phillips, E. P. (1965): Fire — as master and servant: Its influence in the bioclimatic regions of Trans-Saharan-Africa. *Proc. Ann. Tall Timbers Fire Ecol. Conf.* No. 4: 7–109.
- Phillips, J. (1930): Some important vegetation communities in the Central Province of Tanganyika Territory (Formerly German East Africa). *J. Ecol.* **18**: 193–234.
- Phillips, J. D. and Forsyth, D. (1972): Plate tectonics paleomagnetism and the opening of the Atlantic. *Bull. Geol. Soc. Amer.* **83**: 1579–1600.
- Pijmans, K. (1970): An analysis of four tropical rainforest sites in New Guinea. *J. Ecol.* **58**: 77–101.
- Pitt-Schenkel, C. J. W. (1938): Some important communities of warm temperate rainforest at Magamba, West Usambara, Tanganyika Territory. *J. Ecol.* **26**: 50–75.
- Pócs, T. (1974): Bioclimatic studies in the Uluguru Mountains (Tanzania, East Africa). *Acta Bot. Acad. Sci. Hung.* **20**: 115–135.
- Pócs, T. (1976a): Bioclimatic studies in the Uluguru Mountains (Tanzania, East Africa) II., Correlation between orography, climate and vegetation. *Acta Bot. Acad. Sci. Hung.* **22**: 163–183.
- Pócs, T. (1976b): Vegetation mapping in the Uluguru Mountains (Tanzania, East Africa) *Boissiera* **24**: 477–498.
- Ponce-León, P. de (1952, 1953, 1954): Evolución de las formaciones vegetales transitorias en Cuba. *Rev. Soc. Cub. Bot.* **9**: 91–93; **10**: 20–23; **11**: 69–72.
- Ponce-León, P. de (1955): La vegetación de la Laguna de Ariguanabo. *Rev. Soc. Cub. Bot.* **12**: 48–53.
- Précésényi, I. (1972): *Természetes terresztris ökoszisztemák primér produkcíójá*. Akadémiai Kiadó, Budapest 127 pp.
- Proctor, J. (1970): Magnesium as a toxic element. *Nature*, London **227**: 742–743.
- Proctor, J. (1971a): The plant ecology of serpentine II. Plant response to serpentine soils. *J. Ecol.* **59**: 397–410.
- Proctor, J. (1971b): The plant ecology of serpentine III. The influence of a high magnesium/calcium ratio and high nickel and chromium levels in some British and Swedish serpentine soils. *J. Ecol.* **59**: 827–842.

- Proctor, J. and Woodell, S. R. J. (1971): The plant ecology of serpentine I. Serpentine vegetation of England and Scotland. *J. Ecol.* **59**: 375–395.
- Puig, H. (1972): La sabana de Hunnanguillo, Tabasco, Mexico. *Mem. I. Congr. Latinoamer. Bot. (Mexico)*, 389–411.
- Pulle, A. A. (1906): *Enumeration of the Vascular Plants Known from Surinam*. Leyden.
- Pulle, A. A. (1938): Exploraciones botánicas de Surinam. — *Ann. Reuniao Sul. Amer. Bot.* **1**: 239–248.
- Puri, G. S. (1960): *Indian Forest Ecology. A Comprehensive Survey of Vegetation and its Environment in the Indian Subcontinent*. 1–2, 710 pp.
- Rand, A. S. (1973): Long-term stability in a tropical environment: Year to year fluctuations on B. C. I. *II. Intern. Symp. on Trop. Ecol. Caracas*. Venezuela (abstracts).
- Randall, E. R. (1970): Vegetation and environment on the Barbados coast. *J. Ecol.* **58**: 155–172.
- Raunkjaer, C. (1980): *Livsformernes Statistik Som Grundlag for biologisk Plantegeografi*. *Bot. Tidskr.* 1–34.
- Raunkjaer, C. (1934): *The Life-form of Plants and Statistical Plant Geography*. Clarendon Press, Oxford, 632 pp.
- Raven, P. H. (1973): The evolution of the Mediterranean floras. In: DiCastri, F. and Mooney, H. A. (eds): *Mediterranean Type Ecosystems, Origin and Structure*. Springer, Berlin, 213–224 pp.
- Rawitscher, F. (1948): The water economy of the vegetation of the “Campos Cerrados” in Southern Brazil. *J. Ecol.* **36**: 238–268.
- Read, W. (1975): The genus *Thrinax* (Palmace: Coryphoideae) *Smiths. Contrib. Bot.* No. **19**: 1–38.
- Rensburg, H. J. van (1971): Fire: Its effect on grasslands, including swamps — Southern, Central and Eastern Africa. *Proc. Ann. Tall Timbers Fire Ecol. Conf.* **11**: 175–199.
- Reyes, D. (1982): Estudio taxonómico y fitogeográfico de la subfamilia más importante de las Hepáticas epífilas (Lejeuneaceae: Cololejeuneoideae) en Cuba. Thesis, C. Sc. Budapest, 1–12 pp.
- Richards, P. W. (1936): Ecological observations on the rainforest of Mount Dulit, Sarawak. *J. Ecol.* **24**: 1–37.
- Richards, P. W. (1952): *The tropical rainforest*, Cambridge, Univ. Press 450 pp.
- Richards, P. W. (1955): The secondary succession in the tropical rainforest. *Sci. Prog.* **43**: 45–57.
- Richards, P. W. (1959, 1961): The types of vegetation of the humid tropics in relation to soil. In: *Tropical Soils and Vegetation. Proc. Abidjan Symps.* 1959. Paris, UNESCO 1961.
- Richards, P. W. (1969): Speciation in the tropical rainforest and the concept of niche. *Biol. J. Linn. Soc.* **1**: 149–153.
- Richards, P. W. (1973): Africa, “The Odd Man Out.” In: Maggers, J. B. J., Ayensu, C. S. and Duckworth, W. D.: *Tropical Forest Ecosystems in Africa and South America: A Comparison Review*. Smiths. Inst. Washington pp. 21–26.
- Richards, P. W., Tansley, A. G. and Watt, A. S. (1940): The recording of structure, life form and flora of tropical forest communities as a basis for their classification. *J. Ecol.* **28**: 224–229.
- Richardson, W. D. (1963): Observations on the vegetation and ecology of the Aripo Savannas. Trinidad. *J. Ecol.* **51**: 295–313.
- Robinson, W. C., Edgington, and Byers, H. (1935): Chemical studies on infertile soils derived from rocks high in magnesium and generally high in chromium and nickel. *U. S. Pept. Agricult. Techn. Bull.* 471.
- Roig, T. J. (1919): Las maderas de Isla de Pinos. *Rev. Agr. Comerc. Trab.* **2**: 498–500.
- Roig, J. T. (1922): Reseña sobre una excursión botánica a Isla de Pinos. *Mem. Soc. Cub. Hist. Nat.* **5**: 72–76.
- Roig, J. T. (1928): *Diccionario Botánico de Nombres Vulgares. Cubanias. Cons. Nac. Univ.* La Habana I–II, 1142 pp.
- Rollet, B. (1969): La régénération naturelle en forêt dense humide semperfivente de plaine de la Guyane Vénézuélienne. — *Bois For. Trop.* **124**: 19–38.
- Rosenkrantz, F. (1936): Klimacharakter und Pflanzendecke. *Öst. Bot. Zschr.* **85**: 183–212.
- Rune, O. (1953): Plant life on serpentine and related rocks in the North of Sweden. *Acta Phytogeogr. Suecica* **36**: 1–139.
- Russel, M. B. and Hurlbut, L. W. (1959): The agricultural water supply. *Adv. in Agricult.* **11**: 6–19.
- Rutten, M. G. (1956): Geology of the northern part of the Province of Santa Clara (Las Villas), Cuba. *Geogr. en Geol. Med., Physiogr., Geol. Reeks.*, Ser. 2., Utrecht, No. 11., 1–59.
- Saakov, S. G. (1970): Endemic palms of Cuba. *Bot. Zhurn.* **55**: 196–221.

- Sagra, R. de La (1845, 1850): Historia, fisica, politica y natural de la Isla de Cuba II. *Hist. Nat.* Paris X–XI.
- Samek, V. (1967): Regiones naturales de Isla de Pinos. *Acad. Cien. Cuba, Serie Isle de Pinos* 3: 1–36.
- Samek, V. (1968a): La protección de la naturaleza en Ciba. *Acad. Cien. Cuba, Ser. Transform. Natur.* 7: 1–23.
- Samek, V. (1968b): Regiones naturales de la provincia Pinar del Rio bajo el aspecto de la planificación física. *Acad. Cien. Cuba, Ser. Pinar del Rio* 15: 1–23.
- Samek, V. (1969): La vegetación de la Isla de Pinos. *Acad. Cien. Cuba, Ser. Isle de Pinos No.* 28: 1–28.
- Samek, V. (1973a): Pinares de Cajalbana. *Acad. Cien. Cuba, Ser. Forestal* 13: 1–56.
- Samek, V. (1973b): Pinares de la Sierra de Nipe. *Acad. Cien. Cuba, Ser. Forestal* 14: 1–58.
- Samek, V. (1973c): Regiones fitogeográficas de Cuba. *Acad. Cien. Cuba, Ser. Forestal* 15: 1–60.
- Samek, V. (1973d): Vegetación litoral de la costa norte de la provincia de la Habana. *Acad. Cien. Cuba, Ser. Forestal* 18: 1–85.
- Samek, V. and Duck, J. J. (1967): Nombres topográficos derivados de árboles y formaciones vegetales en la Provincia de Pinar del Rio. *Acad. Cien. Cuba, Ser. Pinar del Rio* 2: 11 pp.
- Samek, V. and Moncada, M. (1971): Comunidades vegetales de las lagunas blancas de Pinar del Rio. *Cuba. Acad. Cien. Cuba, Ser. Pinar del Rio*, 27: 1–48.
- Samek, V. and Travieso, A. (1968): Climaregiones de Cuba. *Rev. Agricult. Havana* 2: (1). 5–23.
- Samkova, H. and Samek, V. (1967): Bibliografía Botánica Cubana. *Acad. Cien. Cuba, Ser. Biológica* 1–36.
- San José, J. J. and Medina, E. (1975): Effect of fire on organic matter production and water balance in a tropical savanna. In: Golley, F. B. and Medina, E. (eds): *Tropical Ecological Systems. Ecol. Studies* 11: 251–264.
- Santamaría, E. and Bónazzi, A. (1963): Factores edáficos que contribuyen a la creación de un ambiente xerofítico en el Alto Llano de Venezuela: El arrecife. *Bol. Soc. Venez. Cienc. Nat.* 106: 9–17.
- Sarasin, F. (1917): *Neu-Caledonien und die Loyalty-Inseln*. Georg. Co., Basel.
- Sarmiento, G. (1968): Correlación entre los tipos de vegetación de América y dos variables climáticas simples. *Bol. Soc. Venez. Cienc. Nat.* 113–114: 454–476.
- Sarmiento, G. (1972): Ecological and floristic convergences between seasonal plant formations of tropical and subtropical South America. *J. Ecol.* 60: 367–400.
- Sarmiento, G. and Monasterio, M. (1969): Studies on the savanna vegetation of the Venezuelan llanos. I. The use of association-analysis. *J. Ecol.* 57: 879–898.
- Sarmiento, G. and Monasterio, M. (1975): A critical consideration of the environmental conditions associated with the occurrence of savanna ecosystems in tropical America. In: Golley, F. B. and Medina, E. (eds) *Tropical Ecological Systems. Ecol. Studies* 11: 223–250.
- Sarukhan, K. J. (1964): Estudio sucesional de un área talada en Tuxtepec. *Oac. Publ. Especial Inst. Nac. Invest. For. Mexico* 3: 107–172.
- Sauvalle, F. A. (1868–1872): Flora Cubana. *Acta Acad. Havana* VI–VII–VIII.
- Sauvalle, F. A. (1873): *Flora Cubana*, Havana, 414 pp.
- Sawyer, J. O., Jr. and Lindsay, A. A. (1964): The Holdridge bioclimatic formations of Eastern and Central United States. *Proc. Indiana Acad. Sci.* 72: 105–112.
- Schimper, A. F. W. (1898): *Pflanzengeographie auf physiologischer Grundlage*. I–II. Fischer, Jena, 839 pp.
- Schnell, R. (1970–1971): *Introduction à la Phytogéographie des pays tropicaux*. I–II. Ed. Gauthier-Villars, Paris, 951 pp.
- Schubert, R. and Danert, S. (1969): Über die Vegetationsverhältnisse in zwei Naturschutzgebieten Kubas. I. Das Naturschutzgebiet von El Beral. *Arch. Naturschutz Landschaftsforsch.* 9: 271–299.
- Schubert, R. and Danert, S. (1971): Über die Vegetationsverhältnisse in zwei Naturschutzgebieten Kubas. II. Das Naturschutzgebiet Cupeyal. *Arch. Naturschutz Landschaftsforsch.* 11: 17–41.
- Schuchert, C. (1935): *Historical Geology of the Antillean-Caribbean Region*. John Wiley and Sons, New York, 811. pp.
- Schulz, J. P. (1960): Ecological studies on rainforest in northern Suriname. *Verh. Kon. Ned. Akad. Wet.* 53: 1–267.
- Schulz, J. P. (1967): La regeneración natural de la selva mesofítica tropical de Surinam después de su aprovechamiento. *Bol. Inst. For. Lat. Amer. Invest. Cap.* 23: 3–27.

- Scott, J. D. (1934): Ecology of certain plant communities of the Central Province, Tanganyika Territory. *J. Ecol.* **22**: 177–229.
- Scott, V. J. (1971): Veld Burning in Natal. *Tall Timbers Fire Ecol. Conf.* **11**: 33–51.
- Seifriz, W. (1940): Die Pflanzengeographie von Cuba. *Bot. Jahrb.* **70**: 441–462.
- Seifriz, W. (1943): The plant life of Cuba. *Ecol. Monogr.* **13**: 375–426.
- Shanbag, G. Y. (1956): The climates of India and its vicinity according to a new classification. method *Ind. Geogr. J.* **31**:
- Silva, G. (1979): *Murcielagos de Cuba* Acad. Cien. Habana, 389 pp.
- Simeonovich, B. (1964): Grafico de corelación entre la temperatura y la altura. La Habana, Inst. Nac. Rec. Hidraulicos. 1–18.
- Siolli, H. and Klinge, H. (1962): Soils, tipos de vegetação e aquas na Amazonia. *Bol. Mus. Paraense Goeldi, Avulsa* **1**: 27–41.
- Smith, E. E. (1954): The forests of Cuba. *Maria Moors Cabot. Found. Publ.* **2**: 1–98.
- Smythc, N. (1970): Relationships between fruiting seasons and seed dispersal methods in a neotropical forest. *Amer. Naturalist* **104**: 25–35.
- Snow, D. W. (1968): Fruiting seasons and bird breeding seasons in the New World Tropics (Abstracts). *J. Ecol.* **56**(2): 5–6.
- Soane, B. D. and Sauder, D. H. (1959): Nickel and chromium toxicity of serpentine soils in Southern Rhodesia. *Soil Sci.* **88**: 322–329.
- Specht, R. L. (1969): A comparison of the sclerophyllous vegetation characteristics of Mediterranean type climates in France, California and Southern Australia. I. Structure, morphology and succession. *Austral. J. Bot.* **5**: 52–85.
- Stark, N. (1970): The nutrient content of plants and soils from Brasil and Surinam. *Biotropica* **2**: 51–60.
- Starker, A. L. (1950): Vegetation zones of Mexico. *Ecology*, **31**: 507–518.
- Stearn, W. T. (1968): Jamaican and other species of *Bumelia* (Sapotaceae). *Arn. Arb.* **49**: 280–289.
- Stearn, W. T. (1969): A synopsis of Jamaican Myrsinaceae. *Bull. Brit. Mus. (Nat.-Hist.) Bot.* **4**: No. 4. 145–178.
- Stearn, W. T. (1971): A survey of the tropical genera *Oplonia* and *Psilanthele* (Acanthaceae) *Bull. Brit. Mus. (Nat.-Hist.) Bot.* **4**, No. 7: 261–323.
- Stebbins, G. L. (1952): Aridity as a stimulus to plant evolution. *Amer. Naturalist* **86**: 33–44.
- Stehlé, H. (1941): Conditions éco-sociologiques et evolution des forêts des Antilles Françaises. *Caribb. Forester* **2**: 154–159.
- Stehlé, H. (1943): La végétation muscinale des Antilles Françaises et son intérêt dans la valorisation sylvicole. *Caribb. Forester* **5**: 20–43.
- Stehlé, H. (1944): Les petites associations épiphiles en forêt hygro-sciophile aux Antilles Françaises. *Caribb. Forester* **5**: 136–137.
- Stehlé, H. (1945): Forest Types of the Caribbean Islands. Part I. *Caribb. Forester* **6**: Suppl. 273–422.
- Steila, D. (1966): An evaluation of the Thornthwaite and Holdridge classification as applied to the Mediterranean borderland. *Profess. Geographer* **18**: 358–364.
- Stepanov, I. (1964): Clasificación preliminar de los suelos de Cuba. Univ. Habana, No. 28: 166–167.
- Stoffers, A. L. (1956): *The Vegetation of The Netherlands Antilles*. Balkema Utrecht, 142 pp.
- Sugden, A. M. (1983): Determinants of species composition in some isolated neotropical cloud forests. *Trop. Rain For.* **2**: 43–56.
- Suess, E. (1908): *Das Antlitz der Erde*. F. Tempsky, Wien, **V**(1) 778 pp.
- Swan, F. R. (1970): Post-fire response of four plant communities in south-central New York state. *Ecology* **51**: 1074–1082.
- Szántó, I. (1949): Erdőgazdaságunk éghajlati adottságai (Climatic conditions of Hungarian forestry). *Erd. Kisér. 49*: 1–58.
- Száva-Kováts, J. (1927): Einige Bemerkungen über die Isonotiden. *Peterm. Geogr. Mitteil.* **73**:
- Száva-Kováts, J. (1931): A klíma nedvességgének problémája (Das Problem der klimatischen Feuchtigkeit.). *Időjárás* **35**: 41–49., 76–77.
- Takhtajan, A. L. (1970): *Proiskoshdenie i rasselenie cvetkovith rasteni*. Izd. Nauka Leningrad. 145 pp.
- Tamayo, F. (1961): Exploraciones botánicas en el estado Bolívar. *Bol. Soc. Venezol. Cien. Nat.* **22**: Nos 98–99: 25–180.
- Tamayo, F. (1962): Adaptaciones de la vegetación pirófila. *Bol. Soc. Venezol. Cien. Nat.* No. 101: 49–58. pp.

- Tanner, C. B. (1960): Energy balance approach to evapotranspiration from crops. *Soil Sci. Soc. Amer. Proc.* **24**: 1–9.
- Tanner, E. V. J. (1977): Four montane rainforests of Jamaica. *J. Ecol.* **65**: 883–918.
- Tanner, E. V. J. (1982): Species diversity and reproductive mechanisms in Jamaican trees. *Biol. J. Linn. Soc.* **18**: 263–278.
- Taylor, B. W. (1963): An outline of the vegetation of Nicaragua. *J. Ecol.* **51**: 27–54.
- Taylor, R. A. (1967): A tropical hammock on the Miami (Florida) limestone — a twenty-five-year study. *Ecology* **48**: 863–867.
- Thai-van-Trung (1962): Ecology and classification of the forest vegetation of Vietnam. (Thesis of D. Sc., in Russian). Leningrad, 41 pp.
- Thai-van-Trung (1967): Classification "Ecosystmatique" de la végétation forestière du Viet-Nam. *Beitr. trop. subtrop. Landwirtsch Tropenveterinärmed* **5**: 277–292.
- Thompson, P. T. (1966): A test of the Holdridge model in midlatitude mountains. *Profess. Geographer* **15**: 286–292.
- Thorne, R. F. (1973): Floristic relationships between tropical Africa and tropical America. In: Meggers, B. J. and Ayensu, C. S.: *The Flora of North and South America: a Comparative Review*. Smiths. Inst., Washington, 27–47.
- Thornthwaite, C. W. (1931): The Climates of North America according to a new classification. *Geogr. Rev.* **21**: 633–655.
- Thornthwaite, C. W. (1933): The climates of the Earth. *Geogr. Rev.* **23**: 433–440.
- Thornthwaite, C. W. (1948): An approach towards a rational classification of climate. *Geogr. Rev.* **38**: 55–94.
- Thornthwaite, C. W. and Mather, J. R. (1955): The water balance. *Publ. in Climate* Drexel Inst. Techn. Center. **8**: No. (1.)
- Thornthwaite, C. W. and Mather, J. R. (1957): Instructions and the water balance. *Publ. in Climate* **10**: (3) 183–311.
- Tikhomirov, I. N. (1967): Formaciones magmáticas de Cuba y algunas particularidades de su metalogenia. *Rev. Technol.* **5**: (4); 13–22.
- Tixier, P. (1966): *Flore et végétation orophiles de l'Asie tropicale*. Sedes, Paris, 240 pp.
- Tosi, J. A., Jr. (1964): Climatic control of terrestrial ecosystems: A report of the Holdridge model. *Economic Geogr.* **4**: 173–181.
- Tosi, J. A., Jr. (1971): Forest inventory and demonstrations, Panama. Life zones: An ecological basis for silvicultural research and forest inventories in the Republic of Panama. FAO Report No. Fo: SF (PAN 6. Inf. Tecnico **2**, 123 pp.
- Trapnell, C. G. (1959): Ecological results of woodland burning experiments in Northern Rhodesia. *J. Ecol.* **7**: 129–168.
- Trapnell, C. G. and Langdale-Brown, I. (1971): The natural vegetation of East Africa. In: *The Natural Resources of East Africa*, 92–102.
- Trewartha, G. T. (1954): *An Introduction to the Climate*. McGraw-Hill, New York.
- Troll, C. (1935): Gedanken und Bemerkungen zur ökologischen Pflanzengeographie. *Geogr. Zschr.* **4**: 380 pp.
- Trusov, I. I. (1965): Mapa de isolinias de las precipitaciones en período húmedo. La Habana, Inst. Nat. Rec. Hidráulicos.
- Trusov, I. I. (1967): *Las precipitaciones en la Isla de Cuba*. La Habana Inst. Nac. Rec. Hidráulic. 1–63 pp.
- Tüxen, R. and Diemont, H. (1937): Klimaxgruppe und Klimaxschwarm. Ein Beitrag zur Klimaxtheorie. *Jahresber. Naturhist. Ges. (Hannover)* **88–89**: 73–87.
- Urban, I. (1898): Bibliographia Indiae occidentalis botanica. *Symb. Ant.* **1**: 3–195.
- Urban, I. (1923): Zur Pflanzengeographie von Hispaniola. *Symb. Ant.* **9**: (1): 1–54.
- Urban, I. (1924–1928): Plantae cubenses novae vel rariores a clo. E. L. Ekman lectac. *Symb. Ant.* **9**: (1): 55–176. (2.): 177–272 (1924). — **9**(3): 398–432. (1925). — **9**(4): 435–543 (1928).
- Vales, M. A., Borhidi, A. and Del-Risco, E. (1982): Anatomía de la madera de Myricaceae en Cuba: Consideraciones ecológicas. *Acta. Bot. Acad. Sci. Hung.* **28**: 241–253.
- Vales, M. A. and Süss, H. (1985): Die Holzanatomie der in Kuba endemischen Rubiaceae *Acunaeanthus tinifolius* (Griseb.) Borhidi, *Ariadne shaferi* (Standl.) Urb. und *Neomazaea phialanthoides* (Griseb.) Krug et Urb. *Feddes Repert.* **96**: 215–225.
- Van Steenis, C. G. G. J. (1958a): Basic principles of rain forest sociology, In: *Study of Tropical Vegetation. Kandy Symp.* UNESCO, Paris, 159–163 pp.

- Van Steenis, C. G. G. J. (1958b): Rejuvenation as a factor for judging the status of vegetation types: The biological nomad theory. In: *Study of Tropical Vegetation*. Proc. Kandy Symp. 212–215. UNESCO, Paris.
- Van Steenis, C. G. G. J. (1962): The Land-Bridge Theory in Botany. *Blumea* **11**: 235–372.
- Vareschi, V. (1960): La estación biológica de los Llanos de la Sociedad Venezolana de Ciencias naturales y su tarea. *Bol. Soc. Venez. Cien. Nat.* **21**: No. 96: 107–117.
- Vareschi, V. (1962): La quema como factor ecológico en los Llanos. *Bol. Soc. Venez. Cien. Nat.* **23**, No. 101: 9–26.
- Vesey-FitzGerald, D. F. (1963): Central African grasslands. *J. Ecol.* **51**: 243–274.
- Vesey-FitzGerald, D. F. (1965): The utilization of natural pastures by wild animals in the Rukwa valley, Tanganyika. *East Afr. Wildl. J.* **3**: 38–48.
- Vesey-FitzGerald, D. F. (1970): The origin and distribution of valley grasslands in East-Africa. *J. Ecol.* **58**: 51–76.
- Vesey-FitzGerald, D. F. (1971): Fire and animal impact on vegetation in Tanzania National Parks. *Proc. Ann. Tall Timbers Fire Ecol. Conf.* **11**: 297–317.
- Vézda, A. and Samek, V. (1972): Cladonien der Kieferwälder im NO der Provinz Oriente, Cuba. *Acta Mus. Siles. Ser. A* **21**: 1–18.
- Vidal, J. (1960): *La végétation du Laos*. Trav. Lab. Forest. Toulouse, 513 pp.
- Vidal, J. (1961): Application au Laos de deux méthodes bioclimatiques. *R. C. Soc. Biogeogr.* **334**: 53–64.
- Vidal, J. (1966): Aspects biogéographiques du Sud-Est asiatiques. *R. C. Soc. Biogeogr.* **378–379**: 130–140.
- Voronov, A. G. (1970): Mapa de vegetación. *Atlas Nacional de Cuba*. Havana, pp. 56–59.
- Vuilleumier, B. S. (1971): Pleistocene changes in the fauna and flora of South America. *Science* **173**: 772–780.
- Wadsworth, F. H. (1947): Growth in the lower mountain rain forest of Puerto Rico. *Caribb. Forester* **8**: 27–44.
- Wadsworth, F. H. (1954): Tropical rain forest. The *Dacryodes-Sloanea* association of West-Indies. *Act. du IV<sup>e</sup> Congr. Forest. Mond.* **3**, Dehra Dun.
- Wadsworth, F. H. (1959): Growth and regeneration of white mangrove in Puerto Rico. *Caribb. Forester* **20**: 59–72.
- Waibel, L. (1943): Place names as an aid in the reconstruction of the original vegetation of Cuba. *Geogr. Rev.* **33**: 376–396.
- Waibel, L. (1948): Vegetation and land-use in the Planalto Central of Brazil. *Geogr. Rev.* **38**: 529–544.
- Walker, R. B. (1954): Factors affecting plant growth on serpentine soils. *Ecology* **35**: 259–266.
- Walter, H. (1951): *Einführung in die Phytologie III/1*. Standortslehre. Fischer Stuttgart.
- Walter, H. (1955): Die Klimadiagramme als Beurteilung der Klimaverhältnisse für ökologische, vegetationskundliche und landwirtschaftliche Zwecke. *Ber. Deutsch. Bot. Ges.* **68**: 331–344.
- Walter, H. (1962): Die Vegetation der Erde in ökophysiologischer Betrachtung. I. *Die tropischen und subtropischen Zonen*. Fischer, Stuttgart–Jena, 538 pp.
- Walter, H. (1970a): *Vegetationszonen und Klima*. Fischer Jena, 244 pp.
- Walter, H. (1970b): El problema de la sabana. *Bol. Soc. Venez. Cien. Nat.* **115–116**: 121–144.
- Walter, H. and Lieth, H. (1960–1966): *Klimadiagramm-Weltatlas*. Fischer Jena.
- Walter, H. and Medina, E. (1971): Características climáticas de Venezuela sobre la base de clímadigramas de estaciones particulares. *Bol. Soc. Ven. Cien. Nat.* No. **119–120**: 211–240.
- Warming, E. (1982): Lagoa Santa. *Kgl. Danske Vidensk. Selsk. Skr. Afd.* **6**: 150–487.
- Warming, E. and Graebner, P. (1918): *Lehrbuch der ökologischen Pflanzengeographie*. Berlin.
- Watt, A. S. (1947): Pattern and process in the plant community. *J. Ecol.* **35**: 1–22.
- Webb, L. J. (1959): A physiognomic classification of Australian rainforests. *J. Ecol.* **47**: 551–570.
- Webb, L. J., Tracey, J. G., Williams, W. T. and Lance, G. N. (1967): Studies in the numerical analysis of complex rain-forest communities. I. A comparison of methods applicable to site/species data. *J. Ecol.* **55**: 171–191.
- Webb, L. J., Tracey, J. G., Williams, W. T. and Lance, G. N. (1970): Studies in the numerical analysis of complex rain-forest communities. V. A comparison of the properties of floristic and physiognomic-structural data. *J. Ecol.* **58**: 203–232.
- Webb, L. J., Tracey, J. G. and Williams, W. T. (1972): Regeneration and pattern in the subtropical rainforest. *J. Ecol.* **60**: 675–695.

- Webster, G. L. (1956–1958): A monographic study of the West Indian species of *Phyllanthus*. *J. Arn. Arb.* **37**: 217–268, 340–359, **38**: 51–79, 170–198, 295–373, **39**: 49–100, 111–212.
- Wegener, A. (1929): *The Origin of Continents and Ocean Basis*, Deser. N. Y. 217 pp.
- Welch, J. R. (1960): Observations on deciduous woodland in the eastern province of Tanganyika. *J. Ecol.* **48**: 557–573.
- West, O. (1965): Fire in Vegetation and its Use in Pasture Management with Special Reference to Tropical and Subtropical Africa. Commonwealth Bureau of Pastures and Field Crops, Hurley, Berkshire, England. No. 1/1965.
- West, O. (1971): Fire, man and wildlife as interacting factors limiting the development of climax vegetation in Rhodesia. *Proc. Ann. Tall Timber Fire Conference*, No. **11**: 121–145.
- Weyl, R. (1966): *Geologie der Antillen*. Gebrüder Borntraeger, Berlin. 410 pp.
- Whittaker, R. H. (1953): A consideration of Climax Theory: The Climax as a population and pattern. *Ecol. Monogr.* **23**: 41–78.
- Whittaker, R. H. (1954): The vegetation response to serpentine soils. *Ecology* **35**: 275–288.
- Whittaker, R. H. (1965): Dominance and diversity in land plant communities. *Science* **147**: 250–260.
- Whittaker, R. H. (1970): Communities and ecosystems. In: *Curr. Conc. in Biol. Series*. Macmillan, London.
- Wilcock, A. A. (1951): Potential evapotranspiration: a simplification of Thornthwaite's method. *Proc. Roy. Soc. Victoria (Australia)* **63**: 25–30.
- Wild, H. (1968a): Geobotanical anomalies in Rhodesia I. The vegetation of copper bearing soils. *Kirkia* **7**: 1–71.
- Wild, H. (1968b): Phytogeography in South Central Africa. *Kirkia* **6**: 197–222.
- Wild, H. (1970): Geobotanical anomalies in Rhodesia III. The vegetation of nickel-bearing soils. *Kirkia* **7**: (Suppl.) 1–62.
- Wiltshire, G. H. (1972): Effect of nitrogen source on translocation of nickel in some crop plants and weeds. *Kirkia* **8**: 103–123.
- Wiltshire, G. H. (1974): Growth of plants on soils from two metalliferous sites in Rhodesia. *J. Ecol.* **62**: 501–525.
- Woodring, W. P. (1954): Caribbean land and sea through the ages. *Geol. Soc. America Bull.* **65** (8): 719–732.
- Wyk, P. van (1971): Veld Burnings in the Kruger National Park, an interim report on some aspects of research. *Proc. Ann. Tall Timbers Fire Ecol. Conf.* **11**: 9–31.
- Zólyomi, B. (1942): A középdunai flóraválasztó és a dolomitjelenség. (Die Mitteldonau-Florenscheide und das Dolomitphänonomen) *Bor. Közlem.* **39**: 209–224.
- Zólyomi, B. (1958): Budapest és környékének természetes növénytakarója. (Natural vegetation of Budapest and its surroundings. In Hungarian) In: Pécsi, M. (ed.): *Budapest természeti képe*. Budapest, Akadémiai Kiadó 508–642 pp
- Zonn, S. V. (1968): Osobennosti pochvoobrazovania i glavnje tipi poehv Kubi. Genet.i Geogr. Poehv. Dokl. IX. Mezdunar. Kongr. Pochvoved. Moskva, 51–152.

## SUBJECT INDEX

(f=figure, t=table)

- Acacia-Krugiodendron* association 423  
*Acacio belairioidi-Spirotecometum holguinensis* 425  
*Acacio-Caesalpinion coriariae* 617  
*Acacio-Capparidetalia* 617  
*Achlaenetalia piptostachyae* 613  
*Achlaenion piptostachyae* 613  
*Acoelorrapihe wrightii* savannas 445  
*Acoelorrapho-Colpothrinacetalia* 614  
*Acoelorrapho-Colpothrinacion* 614  
*Acoelorrapho-Pinion tropicalis* 431, 445, 577  
*Acrosticho-Conocarpetum erectae* 518 f, 519 t,  
*Acrosticho-Schoenoplectetum americanac* 490,  
491 t,  
administrative distribution 29–31  
adventive elements 233–234  
  extratropical 234  
  neotropical 233  
  paleotropical 233–234  
*Agavo cajalbanensi-Pinetum* 198, 205, 318, 319,  
433, 583  
*Agavo-Melochietum tomentosae* 621 f,  
*Agavo (pendentatae)-Mitracarpetum acunae* 404,  
642  
*Agavo shaferi-Pinetum* 198, 206, 436, 437, 589.  
591 t,  
*Alchorneo-Pseudolmedion spuriae* 530  
*Aldrovando-Utricularietalia* 473  
*Aldrovando-Utricularion* 448, 474  
allochthonous character of rivers 36  
alluvial forests 78, 79, 80, 82, 112, 446, 449, 480  
*Amarantho spinosi-Parthenietum hysterophori*  
644  
*Ambrosio hispidae-Ipomoeetum pedis-caprae* 507  
*Ambrosio hispidae-Ipomoeetum stoloniferae* 507  
American species 233, 237  
amphiatlantic species 233  
*Andropogo-Copernicietalia* 442  
*Andropogoni-Sabalion* 613  
*Andropogo-Pinion cubensis* 439, 593  
*Anemio-Pinetum cubensis* 198, 206, 435, 588, 589 t,  
Antillean-Bahamean species 225  
Antillcan species 224–225, 235  
*Antirheo-Acrosynanthesum minoris* 360, 597  
arborescent forbs and grasses 188  
archipelago phase 268–270  
*Ariadno-Phyllanthesum* 425, 596  
*Ariadno-Phyllanthion* 198, 206, 597  
arid thorny scrubs 423–431, 460, 559–571, 595–611  
  limestone 423–425, 460, 559–571  
  semi-desert 460, 427–431, 617–621  
  serpentine 425, 460, 595–611  
atmospheric humidity 39–40  
  annual distribution 40  
  horizontal distribution 39  
  vertical distribution 39  
*Avicennietalia* 517  
*Avicennionoccidentalis* 517  
axeric tropical climate 72  
*Axonopetum compressi* 644  
*Azollaeo-Salvinion* 470  
Bahamean elements 225–226  
*Baracoense* 349, 361–362  
*Batidetum maritimae* 514  
*Batidi-Avicennietum germinantis* 517 t,  
*Batidion maritimae* 514  
*Batidi-Salicornietalia ambiguae* 513  
*Batidi-Salicornietea* 513  
Beard's savanna concept 153–154  
bioclimate and vegetation transects 77–85  
  of Central Cuba 79–80  
  of Havanna, Matanzas and Camagüey 78–79  
  of Pinar del Rio 77  
  of the mountains of Oriente 81–88  
bioclimate types of Cuba 62, 75–91  
bioclimatic indexes of Meher Homji 103–105  
bioclimatic systems 56  
  of Gaussem 57–60  
  of McHer-Homji 103–105  
biotype exhaustion 126  
bipolarity 246–247, 273–274

- bisectorial distribution 248  
 bixeric climates 93–96  
*Blechno-Acoelorrhaphetum* 448, 582  
*Blechno-Acoelorrhaphion wrightii* 579  
*Bletio purpureae-Andropogonetum gracilis* 613  
*Bombacopsi-Burserion simarubae* 550  
*Bombacopsi-Catalpetum punctatae* 199, 206, 412,  
 413 f, 552  
*Bombacopsi-Gaussietum principis* 553  
*Bombacopsi-Thrinacetalia* 417, 552  
*Bombacopsi-Thrinacetum morrisii* 418, 555, 556 t.  
*Borrichio-Mallotonietalia* 508  
*Borrichio-Mallotonietum* 454, 508 t.  
*Borrichio-Mallotonion* 453, 508  
*Borrichio-Rachicallietalia* 455, 511  
*Borrichio-Rachicallietum* 454, 512  
*Brachiario-Dactyloctenietum aegyptii* 644  
*Brasenietum schreberi* 480, 481 t.  
*Bucido-Fraxinetum cubensis* 528  
*Bucido-Pseudocarpidietum ilicifolii* 604  
*Burmanno-bicolori-Hypericetum fasciculati* 498,  
 500, 501 t.  
*Bursereto-Thrinacetum radiatae* 454, 572, 575, 576 t.,  
*Buxo foliosae-Acrosynanthetum revoluti* 637, 639 t.,  
*Byrsinimo-Andropogonetalia teneris* 615  
*Byrsinimo-Andropogonetum teneris* 616  
*Byrsinimo-Andropogonion teneris* 616  
*Byrsinimo minutifoliae-Myrtetum acunae* 597  
*Byrsinimo-Pinetea caribaeae* 576  
*Byrsinimo-Pinetum tropicalis-caribaeae* 579, 580,  
 581 t.
- Cabombetum piauhensis* 472, 473 t.,  
*Cabombo-Eichhornietea* 469, 476  
*Cabombo-Najadetalia* 477  
*Cabombo-Nymphaeeta* 476  
 Ca-deficiency 124, 133  
 Cajalbana Mountains 316–318, 319, 320  
*Cajalbanense* 291, 316–318, 319, 320  
*Calophyllo-Carapetum guianensis* 198, 206, 389,  
 390 f, 529  
*Calophyllo-Dipholian* 528  
*Calophyllo-Coccobolbetum retusae* 530  
*Calophyllo-Guettardion valenzuelanae* 530  
*Calophyllo-Swietenietum mahagoni* 539  
*Calyptronomo clementis-Cyrillietum racemiflorae*  
 631  
*Calyptronomo clementis-Cyrillion racemiflorae*  
 631  
*Calyptronomo-Cyrilletalia* 626  
*Calyptronomo-Cyrillietum racemiflorae* 319, 630  
*Calyptronomo-Cyrillion racemiflorae* 630  
*Camagüeyense* 336, 339–341  
*Camagüeyicum* 323, 336–337  
 Ca/Mg ratio 124, 125, 133, 134
- Canavalo-Ipomoeetalia* 506  
*Cappari-Ritterocereetum hystricis* 620  
 caribbean elements 224–228, 235–237  
 north-caribbean 226–227, 235  
 pan-caribbean 227–228, 237  
 south-caribbean 227, 235  
*Cascajalense* 323, 330–331  
*Caseario-Pinetea cubensis* 587  
*Casildense* 323, 331, 332  
*Cassio lineatae-Aristidetum neglectae* 614  
*Cassio lineatae-Aristidion neglectae* 614  
*Catalpa-Swietenia* association 410  
 Cauto basin 346  
*Cautoënse* 336, 346  
*Ceibetea occidentalis* 534  
*Ceibo-Roystonion* 442, 612  
*Celtidi-Hebestigmetum cubensis* 555  
*Centrosmo-Panicetum maximae* 644  
*Ceratopyxidi-Gaussietum principis* 199, 206, 553  
*Cercidi-Cereetea* 617  
*Cercidi-Prosopidetea* 616  
*Chaetolepidi-Rhynchosporetum filifoliae* 499  
 chamaephytes 189, 194, 204, 205  
*Chamaesyci-Cynodontetum dactylonis* 644  
 chorographical (chorological) spectrum 212–215,  
 457, 463–465  
*Chrysobalano-Annonetalia glabrae* 519  
*Chrysobalano-Annonetea glabrae* 519  
*Chrysobalano-Annonetum glabrae* 448, 520 f, 521 t.  
*Chrysobalano-Annonion glabrae* 199, 206, 520  
*Chrysobalano-Colpothrinacetum wrightii* 582  
 Ciego de Ávila-Alto Cedro-La Maya Plains 345–  
 346
- circumpolar species 233  
*Cladietea jamaicensis* 485  
*Cladietum jamaicensis* 450  
*Cladion jamaicensis* 496  
*Claraënse* 323, 336, 338–339  
 classification of vegetation 23, 456–465  
*Clethro-Pinetum maestrensis* 441 f, 623, 624–625 t.,  
 climate system 55–56  
 of Köppen 55  
 of Thornthwaite 56  
 of Trewartha 55  
 climate-vegetation correlations 88–93  
 climatic climax 136, 137, 147  
 climatic equivalence 137  
 climatic grassland concept 148–149  
 climatic water balance 109–110  
 climax formations 106  
 cloudforests 398–400, 639–641  
*Clusio-Ilicetalia* 400, 643  
*Clusio-Ilicetea* 400, 643  
*Clusio-Ilicetum shaferi* 644  
 coastal area of Belic-Pilón 378  
 coastal area of Cienfuegos-Casilda 331, 332

- coastal area of Marea Portillo-Nima nima 378–379  
 coastal terrace belt of Santiago-Guantanamo-Maisi-Baracoa 379–385  
 coastal vegetation 451–455, 503–518  
*Coccolobetalia uviferae* 572  
*Coccolobetea uviferae* 571  
*Coccolobetum uviferae* 454, 572  
*Coccolobion uviferae* 572  
*Coccolobo diversifoliae-Burseretum* 199, 206, 415  
*Coccolobo-Thrinacetum radiatae* 454, 572, 574 t.  
*Coccohrinaci-Bucidetumophiticola* 604, 606 t.  
*Coccohrinaci-Copernicietea* 442  
*Coccohrinaci-Plumerietea* 545  
*Coccohrinaci-Tabebiuetum vibracteolatae* 199, 206, 418, 557, 558 t.  
*Coccohrinaci-Tabebiun lepidotae* 164, 166, 199, 206, 425, 602, 614  
*Combretalia* 518  
 competition on serpentine 126, 135–136  
 coniferous forests 431–440, 460, 576–584, 587–594  
*Conocarpo-Bucidetum palustris* 527 t, 528  
*Conocarpo-Laguncularietum racemosae* 518  
*Conocarpo-Laguncularion* 518  
*Conocarpo-Rachicallietum americanae* 454, 511, 512 t,  
*Consoleo-Colubrinetum ellipticae* 620  
*Consoleo-Ritterocereion hystricis* 617  
 continuity of vegetation 457  
*Copernicia* agricultural savannas 442–443, 613  
*Copernicio-Bucidetum spinosae* 520, 522 t.  
*Copernicion giganti-rigidiae* 613  
*Copernicio-Tabebiuetum trachycarpace* 164, 165, 166, 199, 206, 425, 606, 609 t,  
*Cordio-Crotonetum stenophylli* 568  
 correlation  
 between soil and vegetation types 122–123  
 cosmopolitan species 233, 237, 238  
*Crino-Cladietum jamaicensis* 448, 496, 497 t,  
*Cristalense* 349, 356, 357  
*Crotomo-Lantanion* 566  
*Curatello-Byrsinimeta* 613  
 cyclones 49–50  
*Cyperetum articulati* 488, 490 t.  
*Cyperetum gigantei* 494, 495 t.  
*Cyperion heterophylli* 486  
*Cypero heterophylli-Pennisetetalia* 486  
*Cyrillo-Myrsinion* 640  
*Cyrillo nipensi-Pinetum cubensis* 198, 206, 626  
*Cyrillo-Pinion cubensis* 398, 626  
  
*Dacryodes-Sloanea* community 391  
*Dalbergio-Rhizophoretum manglis* 516, 516 t.  
 deciduous tropical forests 94, 97, 106, 107, 541–542, 544–545, 546, 552–553  
 descendance of montane vegetation 138–139, 141  
 desert 90, 97  
*Dichanthietalia annulata* 644  
*Dichanthion annulata* 644  
*Dipholi-Calophylletalia* 528  
 discontinuity of vegetation 457  
 disjunction 245–249  
*Distichlio-Spartinetalia* 514  
*Distichlion spicatae* 514  
 distribution of savannas 173–174  
*Dracaeno-Pinetum cubensis* 198, 206, 360, 435, 590, 592 f, 593 t.  
 drainage savanna concept 149, 153  
 dry forests 77–79, 82–87, 90–91, 98, 419–423, 460, 545–547  
 deciduous tropical 90  
 evergreen 77, 79, 94, 98, 419, 420–423, 460, 545–547  
 limestone 82, 83, 84, 86, 87, 449  
 littoral 77, 78, 79, 449  
 subtropical 90–91  
 dry season 47–48  
 dry shrubwoods 78–80, 82–83, 98, 123, 423–425, 460, 548–550, 559–566, 571–576, 595–611  
 littoral limestone 78, 80, 122, 449, 571–576  
 lowland serpentine 78, 79, 83, 123, 425, 460, 595–611  
 thorny limestone 80, 82, 83, 98, 123, 423–425, 449, 460, 548–550, 559–566  
 dry thickets 123, 371, 567–571  
 dwarf palm savannas 79, 82, 107, 123, 162–163, 174, 443–444, 614  
 dwarf succulents 190  
 dysgeogenic effect 126  
  
 eastern central Cuba 336–337  
 eastern Sierra Maestra 372–373  
 ecological effects of bixeric climates 93–96  
 ecological coordinate systems 97–98, 106–108  
 ecological equivalences 108  
     of Meher Homji 106–108  
     of Walker 97–98  
 edaphic climax 137, 147, 153  
 edaphic extrazonality 138  
*Eichhornietum azureae* 448, 471, 472 t.  
*Eichhornietum crassipedis* 448, 471 t,  
*Eichhornio heterospermae-Nymphoidetum aureae* 482, 483 t.  
*Eichhornion azureae* 471  
*Ekmanio-Erythroxyletum baracoënsis* 557, 559 t.,  
 elements of temperate North America 269–270  
*Eleocharitetum intersinctae* 492, 493 t.  
*Eleocharasetum cellulosae* 487, 489 t.  
 elfin forest 82, 91, 106, 122, 139, 360, 371  
 elfin thickets 82, 90, 91, 106, 123, 139, 360, 371,  
     372

- semi-dry (semi-arid) serpentine 123, 360, 400–402, 643–644  
 wet 123, 360, 371, 402–404, 641–642  
*Encopello-Hydrolaeetum nigricaulis* 501, 504 t.,  
 endemic genera 128–129, 288–290  
   of Cuba 288–289  
   of Hispaniola 289  
   of Jamaica 290  
   of Puerto Rico 290  
   of serpentines in Cuba 128–129  
 endemic speciation 242–246  
 endemics 216–224, 234–235, 238, 239–241, 242, 243, 244, 245  
 horizontal distribution 243–244  
   of Camagüey 219–220  
   of eastern Cuba 220–221  
   of Habana 219  
   of Isla de Pinos 217–218  
   of Las Villas 219  
   of Matanzas 219  
   of Oriente 222  
   of Pinar del Rio 216–217, 218  
   of western Cuba 218–219  
 pan-Cuban 222–223  
 vertical distribution 243, 245  
 epiphytes 187, 190, 195, 196, 201–204  
 ferns 195–196  
 herbs 190  
 woody 187  
*Eragrosti cubensi-Pinetum* 301, 431, 578  
*Eriocaulo-Paepalanthion* 499  
*Eriothali-Strumpfietum maritimae* 512, 513 t.,  
*Ernodeo-Surianetum maritimae* 509  
*Erythroxyl-Coccolobetum armatae* 600, 601–602 t.  
*Erythroxyl-Coccothrinacetum alexandri* 566, 568, 569 f., 570–571 t.,  
*Erythroxyl-Spirotecometum holguinensis* 610, 611 t.  
*Eu-Camagüeyicum* 336, 337  
*Eugenio-Capparidion* 548, 551 f.  
*Eugenio-Coccothrinacetum roseocarpae* 604  
*Eugenio-Crotonetum rosmariniodis* 568  
*Eugenio-Cupheetum lobelioidis* 633  
*Eugenio-Erthalietum fruticosae* 454, 561, 563–564 t.,  
*Eugenio-Moacrotonetum trigonocarpi* 600  
*Eugenio-Metopietalia toxiferi* 420, 547  
*Euphorbio helenae-Lyonietum myrsinifoliae* 597, 600 t.,  
*Euphorbio helenae-Pinetum cubensis* 198, 206, 437, 588, 590 t., 598  
*Eu-Pinaricum* 304  
 evergreen oakwoods 77, 431–432, 460, 585–587  
 evergreen shrub-woodland 106–107, 123,  
 evergreen tropical forest 90, 122, 296, 297, 363, evolutionary centres 272–279  
   of limestone vegetation 275–277  
   of mega-xerotherm elements 277–278  
   of montane rainforest elements 278–279  
   of serpentine vegetation 272–275  
*Exostemo-Ginorietum americanae* 633, 635 f., 636 t.,  
*Exostemo-Leucocrotonetum stenophylli* 637 f., 638 t.,  
 extratropical elements 233, 234, 238  
 extrazonality on serpentine 138–139  
*Fimbristyli-Salicornietum perennis* 513, 514 f.,  
*Fimbristyli-Salicornion perennis* 513  
 fire ecology 150  
 fire savanna concept 149–150  
 flatlands and hills  
   of Pinar del Rio and Isle of Youth 295  
 Florida-Antillean-Bahamean elements 226, 238  
 freshwater vegetation 297, 448–450, 469–485  
   free floating 469–472, 473–475  
   rooted emerged 478–485  
   rooted submerged 478–485  
 gallery forest 446–447  
 genetic drift 136  
 geophytes 189, 194  
*Gerонense* 291, 308–310  
*Gibarens* 336, 347–349  
*Gran Piedra* 372–373  
 grassland 80–83, 90–91, 151–152, 171–175, 440–445, 611–615  
   agricultural 442–443  
   anthropic 151, 152, 171–172, 175, 440–445, 611–615  
   artificial 80, 82, 83, 90, 91, 172  
   climatic 169, 171  
   edaphic 171, 613  
   grassland patterns 172–175  
   swamp 171, 444, 613  
 Greater Antillean genera 286–287  
 Greater Antillean species 224, 235, 238  
*Grimmeodendro-Gochnatietum sagraeanae* 454, 550  
*Guaimarense* 336, 345–346  
*Guaimaricum* 336, 343  
*Guamuhaya Massif* 331–333  
*Guanaocabibes peninsula (Guanaocabibense)* 291–292  
*Guantanamense* 349, 379–385  
*Guayaco-Pithecellobietum guadelupensis* 548, 549–550 t.,  
*Guazumo-Ceibetum* 450, 535–538 t.,  
*Guazumo-Cupanion* 534  
*Guettardo-Coccothrinacetum munizii* 564, 566 f., 567 t., 568  
*Guettardo-Jacarandion cowellii* 425, 604

- Guettardo-Pinion cubensis* 435, 587  
*Guettardo valenzuelanae-Pinetum caribaeae* 318, 319, 582  
*Guillemineo-Aristidion refractae* 616  
*Güninense* 323, 329–330  
*Gundlachio-Bryetum subinermis* 638  
*Gundlachio-Bryon subinermis* 638  
*Gynerio-Bambusetalia* 485  
*Gynencion sagittati* 485  
*Gynerietum sagittati* 485
- Halodulo-Ruppietum maritimae* 505 t.  
*Havanense* 323, 328–329  
*Havanicum* 323, 326  
 heavy metal toxicity 127, 132, 133, 135  
*Heliotropio (ternatae)-Pithecellobietum hystricis* 568  
 helo-hydatophytes 189, 194  
 hemicryptophytes 189, 194, 204, 205  
 hemi-therophytes 190  
*Hibisco-Calophylletum antillanae* 538  
*Hibisco-Swietenion mahagoni* 538  
*Holguinense* 336, 341–343  
 holocenotic savanna concept 150–151  
 Honduras flora 265–266  
*Hydrocotyletum umbellati* 479 t.  
*Hydrolaeetalia nigricaulis* 499  
*Hydrolaeetaea nigricaulis* 501  
*Hydrolaeion nigricaulis* 501  
*Hydromistrietum stoloniferae* 505  
*Hyeronimo-Sloanetum curatellifoliae* 397 f., 625, 627–629 t.  
*Hyparrhenietum rufae* 644  
*Hyptido-Paspaletalum conjugati* 644  
*Hyptido-Paspaletum conjugati* 644  
*Hyptido-Paspalion conjugati* 644
- Ilici-Laplaceetum benitoënsis* 644  
*Ilici-Laplaceion moaensis* 198, 206, 400, 644  
*Ilici (turquinensi)-Myricetum cacuminis* 402, 642  
*Ilici-Myricion cacuminis* 198, 206, 402, 641  
 incomplete soil profiles 120  
*Indiosense* 291, 301–303  
 inversion 253–254  
*Ipomoeo-Canavalieturn maritimae* 507  
*Ipomoeo-Canavalion maritimae* 506  
*Ipomoeo-Mallotonietea* 506  
*Ipomoeo-Mallotonion* 453  
*Ipomoeo-Paspaletum distachyi* 507  
*Ipomoeo-Philoxeretum* 453  
*Ipomoeo-Tournefortietea* 506
- Jacquinio-Tabebuietum linearis* 597  
*Jarucoënsis* 323, 326–328
- karstic forests 334, 416–419, 460, 553–559  
 eastern Cuban 418–419, 555–557, 558–559  
 montane 334, 419, 557, 559  
 western Cuban species-poor 418, 554–555  
 western Cuban species-rich 417, 553–554  
 karstic peninsulas (*Peninsularicum*) 290–291  
*Krugiodendro-Drypetetum* 420, 550
- lakes of Cuba 36  
 land-bridge phase 265–368  
*Lantano-Cordielatalia* 559  
*Lantano-Cordion* 560  
 latosolization 144–145  
 leaf-size classes 132, 137, 196–197, 205–206  
 leaf-size index 205–206, 207, 208  
*Lemairocereo-Lasiocrotonetum bahamensis* 568  
*Lemairocereo-Phyllostyletum brasiliensis* 568  
*Lemno-Azolletum caroliniana* 448, 470 t.  
*Leucaenetum leucocephala* 576  
*Leucaenion leucocephala* 574  
 lianes 187, 190, 194–195, 201–203  
 herb 190, 194, 195  
 woody 187  
 life-forms 179–210  
 affinity to vegetation types 198–202  
 analysis 181–183, 192–193  
 categories 180–181, 191–196, 461–462  
 in classification of vegetation 461–463  
 spectrum 182–183, 191–193, 196, 457  
 vicariancy 201–205  
 life-zones 108–114  
 life-zone map of Cuba 111–114  
 limestone karsts of Nipe-Baracoa (*Yatericum*) 349, 362  
 limestones and serpentines of Bahia Honda-Limonar 326–328  
 limestone thorn scrub-woodland 83, 84, 86, 87  
*Limnobietum laevigati* 505, 506 t.  
*Limnocharietum flavae* 485 t.  
*Linociero-Savietum bahamensis* 560–561 t.  
*Lithophilo-Trianthemetum portulacastri* 454, 510 t.  
 littoral dry forests 449, 548–551  
 littoral rock pavement 77, 78, 79, 82, 87, 297, 449, 454, 455  
 littoral sand vegetation 82, 123, 297, 449, 453, 454  
 littoral scrubs and woodlands 82, 559–566, 571–576  
 littoral thickets 82, 123, 297, 566–571  
 littoral zone and islands of Cayo Francés-Turigüano-Cebolla 347–349  
*Lonchocarpo-Ceibetalia* 534  
*Lonchocarpo-Ceibetum* 450  
 lowlands, limestone hills and shores of eastern central Cuba 343  
 lowland serpentine scrub woodlands 79, 80, 82, 123, 595–611  
 low palm grassland 79, 80, 82, 107, 123

- Lysilomo-Burseretum simarubae* 199, 206, 420, 421 f, 542  
*Lysilomo-Burserion simarubae* 541  
*Lysilomo-Metopietum brownii* 539
- Macrocopternio-Sabaletalia* 613  
 Madrean-Tethyan flora 267  
 Madrean Tethyan vegetation 266–268  
*Maëstricum* 349, 368–369  
*Magnolio-acunae-Cyrilletum racemiflorae* 198, 206, 395, 622  
*Magnolia cubensis* 622  
*Magnolio-Laplaceetum angustifoliae* 198, 206, 394 f, 622  
 mangroves 77, 78, 79, 80, 82, 83, 84, 87, 122, 296, 297, 303, 368, 449, 451–453, 515–518  
*Mastichodendro-Cupanietum glabrae* 539, 540–541 t.  
*Mastichodendro-Diphloietum salicifoliae* 538  
*Mastichodendro-Trichiliatum hirtiae* 541  
*Mataybaeo-Pseudolmedictum spuriae* 407 f, 408 f, 530, 531–534 t,  
*Mayacetalia fluviatilis* 475, 492  
*Mayacetum fluviatilis* 475, 476 t,  
*Mayacetum wrightii* 475, 476 t,  
*Mayacion fluviatilis* 475  
 megaphanerophytes 184  
 mesophanerophytes 184, 194  
 Mg toxicity 124, 134  
*Miconia ibaguensi-Quercetum sagraeanae* 585, 586–587 t,  
 micranthia 255  
 microphanerophytes 184, 194  
 microphyllia 254–255  
 migration 270, 271–279  
 mixed oak-pine forests 431–432, 583–585  
 mixed treefern-pine woods 440, 622–623, 624–625  
*Moaënse* 349, 357–361  
*Moanicum* 349, 351–353  
 mocarrero 119, 123, 152, 162, 442, 616  
 mogote-complex 77, 78, 79, 82, 83, 84, 321, 416  
 mogote formation 80, 363, 367, 374  
 mogotes of Baire and northern Sierra Maestra 373, 374, 375, 376  
 mogotes of the Nipe-Yateras zone 364–368  
 monoclimax 147  
 monsoon forest 90, 94, 97  
 montane evergreen serpentine scrub 80, 82, 83, 84, 86, 87, 90, 123  
 montane evergreen thicket 334  
 montane tropical climate 72–75  
 mossy forest 82, 106, 398–400, 639–641  
 mountains of Cuba 33  
   of Camagüey Province 32  
   of Isla de Pinos (Isle of Youth) 34–35  
   of Las Villas Province 32  
   of Oriente Province 34  
   of Pinar del Rio Province 32, 310–311  
   multisectorial areas 249  
*Myrico-Baccharideea* 400, 641  
*Myrico-Baccharidetalia vacciniodis* 400  
*Myrico-Bourrieretum pauciflorae* 401  
*Myrico-Lyonietalia jamaicensis* 400, 641  
*Myriophylletum sparsiflori* 480 t,  
*Myrsino-Nectandretum reticularis* 198, 206, 400, 641  
 nanophanerophytes 184–185  
 natural savannas 151–152, 153, 169–171, 175  
*Nelumbio-Nymphaeion amplae* 480  
*Nelumbonetum luteae* 481, 482 t,  
*Neomazaeo-Phyllanthetum orbicularis* 602, 603–604 t,  
*Neomazaeo-Pinetum caribaeae* 198, 206, 318, 319, 433, 582  
*Neomazaeo-Pinion caribaeae* 433, 582  
*Neomazaeo-Reynosion reusiae* 318, 319, 599  
 neotropical elements 228–231, 233, 237, 238  
*Nipe-Baracoa Massif* 349, 351–353  
*Nipense* 349, 353–355  
 Ni toxicity 125, 132, 133  
 non-serial factors 147, 148  
 North and Central American species 233  
 Northern Las Villas 344  
*Nupharitetum macrophylli* 481, 482 t.  
 nutrient deficiency 124, 127, 133, 149  
*Nymphaeetalia amplae* 478  
*Nymphaeetum amplae* 480, 481 t,  
*Nymphaeo-Cabombetum piauhensis* 472, 473 t,  
*Nymphaeo-Potametum nodosi* 478, 479 t,  
*Nymphaoidetum aureae* 483 t,  
*Nymphaoideum grayanae* 483, 484 t,  
*Ocoteo ekmanii-Cyrilletum racemiflorae* 622  
*Ocoteo-Magnolieta* 392, 621  
*Ocoreo-Magnolieta* 621  
 oligotrophic swamp vegetation 497–503, 504  
*Opuntio-Coccobolbeum uviferae* 572, 573 t,  
*Osmundo-Annoneum glabrae* 448  
*Osmundo-Chrysobalanetum icaconis* 520  
*Oxandro-Burseretalia* 410, 539  
*Oxandro-Burserion* 539  
*Oxandro-Diphloietum jubillae* 199, 206, 405, 406 f, 530  
*Pachyantho poiretii-Pinion caribaeae* 431, 583  
 pachycaul elements 181, 188  
*Paepalantho-Eriocauletalia* 499  
*Paepalantho-Pinetum tropicalis* 164, 166, 198, 206, 303, 431, 498, 577, 578 t,  
 pan-tropical elements 231–233, 237  
 paraclimax 107, 146

- Parthenio-Bidentetalia pilosae* 644  
*Parthenio-Bidentia pilosae* 644  
*Parthenio-Dichamptetea annulata* 644  
*Purvicopernicio-Coccothrinacetalia* 614  
*Purvicopernicio-Coccothrinacion* 443, 614  
*Parvirhynchosporo-Eriocauletea* 497, 499  
*Paspalidieum paludivagi* 450, 487, 488 t.  
*Paspalo-Roystonietalia* 612  
*Paspalo-Sesuvietum portulacastri* 507  
*Peninsularicum* 290–291  
 peri-Afro-American elements 262–265  
 phanerophytic pteridophytes 191–194  
*Philoxero-Unioletum virgatae* 453, 507  
*Phyllantho-Neobracetalia valenzuelanae* 425, 598  
*Phyllantho-Neobracetea valenzuelanae* 595  
*Phyllantho selbyi-Aristidetum neglectae* 615  
*Phyllostylon-Acacia* association 423  
 phytogeographic analysis  
     of savanna communities 163–167  
     of the flora 212–215  
 phytogeographic districts 291–385  
     of central Cuba 326–349  
     of eastern Cuba 353–385  
     of western Cuba 291–321  
 phytogeographic elements 211–239, 463–465  
 phytogeographic sectors and subsectors  
     of central Cuba 326–343  
     of eastern Cuba 351–377  
     of western Cuba 290–311  
 phytogeographic sub-provinces of Cuba  
     central Cuba (Centro-Cubanicum) 321–325  
     eastern Cuba (Oriente-Cubanicum) 349–351  
     western Cuba (Occidente-Cubanicum) 290  
*Picrodendro-Burseretum simarubae* 549  
*Piedraíne* 349, 372–373  
*Pilonense* 349, 378  
*Pinarense* 291, 304–308  
*Pinaricum* 291, 295  
 pine forests 77, 82, 83, 107, 318, 320, 322, 360,  
     367, 431–441, 460, 576–584, 587–594, 576–584,  
     587–594, 622–623, 624–625  
     montane 82, 83, 371, 440, 441, 622–623  
 pine savannas 163, 164, 166, 167, 174, 444  
*Pinetalia cubensis* 587  
*Pinetalia occidentalis-maestrensis* 440, 622  
*Pinetalia tropicalis-caribaeae* 576  
*Pinetum caribaeae* 583  
*Pinetum tropicalis-caribaeae* 164, 166, 198, 206,  
     578  
 pine woodland 77, 83, 303, 318  
*Pinion maestrensis* 623  
*Pino-Aristidion neglectae* 444  
*Pinus caribaea* forests 107, 431–433, 578, 582–583,  
     616  
*Pinus cubensis* forests 107, 435–439, 587–594  
     mesophilous 439–440, 593–594  
     montane 439–440, 593–594  
     xerothermic 435–439, 587–593  
*Pinus maestrensis* montane forest 107, 440, 622,  
     623, 624–625  
*Pinus tropicalis* forest 107, 577–582, 616  
*Pistietum stratiotidis* 448, 471, 472 t.  
*Pithecellobion lentiscifolii* 542  
 plain of Artemisa-Colon 329–330  
 plate phase 261–265  
 plate tectonics 258–260  
*Podocarpo-Bonnetietum cubensis* 198, 206, 626  
*Podocarpo-Byrsimonion orientensis* 398, 623  
*Podocarpo-Sloanetalia* 396, 623  
 polyclimax 147  
*Polygal omissae-Rhynchosporetum diodonis*  
     616  
*Polygoneto-Thalietum geniculatae* 494, 495 t.,  
*Polygonetum densiflori* 484 t.,  
*Pontederio-Cladietum jamaicensis* 448  
*Potametum illinoensis-nodosi* 450, 478, 479 t.,  
*Potamion illinoensis* 478  
 potential evapotranspiration 56, 108–114  
 precipitation 44, 46–48  
     annual distribution 46–47  
     extreme values 48  
     fluctuation 48  
     map of regional distribution 44  
     predictability of zonal vegetation 96–99  
*Pseudocarpidio-Cordion* 199, 206, 562  
*Pseudocarpidio-Guettardetum cueroensis* 564, 565 t.,  
 pseudoxeromorphism 136, 143  
 pseudovirginicity 181  
*Psidium montanum-Calophyllum jacquinii* com-  
     munity 391  
*Purdiaeo-Rondeletion pachyphyllae* 634  
*Puriatense* 349, 362–364  
  
*Quercetalia oleoidis* 583  
*Quercion sagraeanae* 199, 206, 585  
*Querco-Pinetum tropicalis* 583  
*Querco sagraeanae-Pinetum caribaeae* 198, 206,  
     432, 583, 584–585 t.,  
  
 Rachicalli-Borrichion 511  
 rainforest flora of Guyana 269  
 rainforests 90, 91, 94, 97, 98, 122, 139, 389–396,  
     460, 528–539  
     sclerophyllous (serpentine) 83, 86, 87, 91, 122,  
     139  
     seasonal lowland 91, 122, 404–405, 530, 534–  
     539  
     seasonal submontane 91, 321, 322, 405, 530–534  
     semi-arid (semi-dry) montane (serpentine) 122,  
     396–398, 623–625, 626, 627–629  
     submontane 84, 85, 87, 90, 91, 106, 122, 139,  
     363, 367, 389–391, 528–530

- wet montane 80, 82, 83, 86, 87, 90, 91, 106, 107, 122, 139, 321, 334, 374, 392–395, 621–622
- rainy season 47
- relict character 256
- relict cryptogams 261
- relict phanerogams 261
- Rhizophoretalia* 515
- Rhizophorion occidentalis* 515
- Rhizophoro-Avicennietea germinantis* 515
- Rhizophoro-Avicennietum germinantis* 517
- Rhizophoro-Chrysobalanetum icaonis* 525, 526 t
- Rhynchosporo-Eleocharitetum interstinctae* 492, 493 t.
- Rhynchosporo-Xyridetalia* 492, 498
- Rhynchosporo-Eleochariton interstinctae* 491
- Rhynchosporo-Pinetum cubensis* 198, 206, 439, 594
- Rhynchosporo-Xyridion* 498
- Ritterocereetalia hystricis* 617
- Ritterocereo-Guetardetum cueroensis* 620
- Ritterocereo-Lasiocrotonetum bahamensis* 620 f., riverside evergreen scrub 631, 633–639
- riverside forests 334, 389, 446, 626, 630–631, 632
- rivers of Cuba 37
- Roigello-Andropogonetum multinervosi* 616
- Roigello-Andropogonion multinervosi* 616
- Rondeletio-Gesnerietea* 631
- Rondeletio-Ginorietalia* 631
- Rondeletio-Guetardetum clarensis* 164, 165, 166, 199, 206, 425, 605, 608 t,
- Rondeletio microphyllae-Gesnerietum humilis* 633
- Rondeletio microphyllae-Ginorion* 631
- Rondeletio-Purdiaeetalia* 634
- Rosaricum* 291, 310–311
- Rosariense* 291, 319, 321, 322
- rosulate mesophanerophytes 185–187
- rosulate microphanerophytes 187
- royal palm savannas 107, 157, 158, 159, 174, 442, 612
- Roystonea* grassland 77, 78, 80, 82, 83, 367
- Roystonea* marsh forest 348
- Ruppieritalia maritimae* 503
- Ruppition maritimae* 505
- Sabaloënse* 291, 300–301
- Sabalo-Indiosense* 299–300
- Sabalo-Roystonietea* 440, 611
- Sabal* savannas 107, 161–162, 174, 444–445, 613
- Sabal* wet woodland 348
- Sagittario-Eleocharitetum interstinctae* 487, 488 t.
- Saguense* 323, 336, 344
- Salicetum carolinianae* 521, 522
- Salicion carolinianae* 521
- Salicornio-Distichlion spicatae* 515
- salt marsh 123, 513–514
- salt meadow 123
- salt swamp 123, 296, 297
- salt water vegetation 503–518
- coastal meadows 123, 514–515
  - mangroves 515–518
  - rock pavement 454–455, 509–513
  - salines 513–514
  - sandy beaches 79, 80, 84, 453–454, 506–509
  - submarine rooted swards 503, 506
- Salvinietum auriculatae* 448
- Salvinio-Eichhornietalia* 469
- Salvinio-Eichhornietea* 469
- Samaneo-Roystonion* 442, 612
- sampling of vegetation 22–23
- Saneti Spiritus Heights* 335
- sandy seashore 79, 80, 84, 453–454, 506–509
- Santiagicum* 349, 374–375, 377
- Sarcomphalo-Reynosietum septentrionalis* 550
- savannas 90, 91, 94, 97, 107, 123, 148–175, 440–445, 611–616
- agricultural 442–444
  - deciduous 445
  - natural edaphic 172–175, 444, 613
  - seasonally flooded 123, 444, 613, 615
  - secondary 161–175
  - semi-anthropic 171, 174, 175, 322
  - short grass 91, 123, 157, 443–445, 613–616
  - tall grass 91, 123, 157, 442–443, 612
  - treeless 157, 445, 616
- Schoenoplectetum validi* 490, 491 t.
- Schuebert's theory 258
- Scirpenum americanum* 450
- Scirpo-Eleocharitetalia interstinctae* 487
- Sclerio interruptae-Rhynchosporenum semiberbis* 616
- sclerophyllous evergreen scrub (thicket) 94, 123, 360, 425–426, 460, 595–611
- dry thorny 123, 425, 598–611
  - semi-dry 123, 426, 595–598
- sclerophyllous montane rainforest 82, 83, 84, 94, 122, 360, 367
- sclerophyllous vegetation 80, 82, 83, 84, 86, 87, 90, 91, 94, 95
- sclerophylls 142
- scrub 90, 91, 94, 123, 360, 425–426, 460, 595–611
- seashore belt of Bahia Honda-Hicacos 328–329
- seasonal evergreen forest (seasonal rainforest) 77, 78, 80, 82, 83, 84, 86, 87, 97, 98, 106, 107, 122, 139, 296, 297, 321, 322, 363, 371
- lowland 122, 139, 404–405
- submontane 122, 321, 322, 334, 405–410
- semi-anthropic grasslands 171, 174, 175, 322
- semi-arid (semi-dry) montane serpentine shrub-woods 322, 400–402, 460, 643–644
- semi-deciduous forest 77, 78, 79, 80, 82, 86, 87, 90, 91, 94, 97, 98, 106, 107, 122, 296, 297, 303,

- 321, 334, 367, 371, 376, 410–416, 449, 460,  
 539–541, 543  
 mesophytic 122, 321, 322, 334, 410–415  
 xerophytic 122, 415–416  
 semi-desert  
 cactus scrub 84, 86, 87, 98, 106, 377, 427–431, 460  
 climate 62  
 vegetation 90, 91, 97, 98, 106, 617–621  
 semi-dry serpentine vegetation  
     elfin thicket 400–402, 643–644  
     lowland shrublands (thickets) 360, 426–427,  
         460, 595–598  
 serpentine areas of Baracoa and Jauco 361–362  
 serpentine areas of Motembo-Holguín 337  
 serpentine effects on cryptogams 140–142  
 serpentine flora 129–132  
 serpentine mountains of the Nipe-Baracoa Massif  
     (Eu-Moanicum) 349, 353  
 serpentine pine woodland 82, 83, 84, 86, 87, 318  
 serpentine preference 140  
 serpentine savannas 162, 164, 165, 166, 167, 170  
 serpentines of Holguín 341–343  
 serpentines of northern Camagüey 339–341  
 serpentines of Santa Clara 338–339  
 serpentine thickets 123, 318, 425–426, 460, 595–  
     611, 643–644  
*Sesuvio-Ipomoeetum pes-caprae* 453, 507 t.  
*Sesuvio-Rachicallictea* 455, 510  
*Sesuvio-Rachicallaltetum americanae* 511  
*Shafero-Pinetum cubensis* 198, 206, 439, 594  
 siallitization 144–145  
 Sierra del Cristal (Cristalense) 349, 356–357  
 Sierra de los Organos 291, 311–316  
 Sierra del Purial 349, 362–364  
 Sierra del Rosario 291, 319, 321, 322  
 Sierra de Micara 349, 356–357  
 Sierra de Nipe 349, 353–355  
 Sierra Maestra 349, 368–369  
 Sierras de Moa and Toa 349, 357–361  
 silicate heights of Western Cuba 291, 304  
 slatey heights and alluvial plains of Pinar del Rio  
     291, 304–308  
*Sloanea berteriana-Ormosia krugii* association 391  
 soil classifications 117–120  
 soil succession 144–146  
 soil-vegetation relationships 119–120  
 South America–Cuba elements 229–230  
 South American–Antillean elements 228–229  
 southern coasts of Oriente (Santiagicum) 349,  
     374–375, 377  
 southern Isle of Youth (Isla de Pinos) 291, 292–293  
*Spathelio-Bourrierietum pauciflorae* 198, 206, 597  
*Spathelio-Gaussian* 553  
*Spigelio sphagnicolae-Paepalanthetum seslerioidis*  
     498, 502–503 t.  
*Spirituense* 323, 335  
*Spirodeto-Salvinietum auriculatae* 470 t.,  
*Spondias-Oreodoxa* association 389  
*Spondiato-Pithecellobietum lentiscifolii* 542, 544–  
     545 t.,  
*Spondiato-Roystonietum* 538  
*Sporobolo-Spartinetum junceae* 508  
*Sporoboletum indici* 644  
 structure profile of vegetation units  
     deciduous lowland forest 421  
     montane pine forest 441  
     montane semi-dry serpentine rainforest 397  
     semideciduous littoral shrubwood 551  
     semideciduous lowland forest 413  
     semi-desert communities 620, 621  
     submontane rainforest 390  
     submontane seasonal evergreen forest 406, 407,  
         408, 409, 414  
         wet montane rainforest 394  
 succession of vegetation types 144–146  
 succulent phanerophytes 187  
*Sudpineroense* 291, 292–293  
*Surianetum maritimae* 454, 509  
*Suriano-Baccharidetum halimifoliae* 509 t.  
*Suriano-Baccharidion halimifoliae* 509  
 swamp forests 78, 80, 122, 296, 297, 449, 518–528  
 swamp vegetation 122, 296, 297, 303, 448, 449,  
     485–497, 518–528  
*Swietenio-Brosimetea* 528  
*Swietenio-Metopietum brownei* 199, 206, 420  
*Syngonantho-Paepalanthetum alsinoidis* 499  
*Syringodio-Thalassietum* 454, 505, 506 t.,  
*Syringodio-Thalassion* 505  
  
*Tabebuio angustatae-Bucidetum palustris* 448,  
     450, 523–525 t.,  
*Tabebuio-Bucidetalia* 522  
*Tabebuio-Bucidetum palustris* 523  
*Tabebuio-Burseretalia* 541  
*Tabebuio-Burseretea* 541  
*Tabebuio-Coccothrinacetalia* 417, 555  
*Tabebuio-Coccothrinacion* 418, 557  
*Tabebuio sauvalei-Garryetum* 199, 206, 419, 557  
*Tabebuio-Viticetum guanahacabibensis* 561, 562 t.  
*Tablaense* 349, 373–374, 375, 376  
 temperature 39–45  
     annual distribution 39, 45  
     extreme values 39–40  
     fluctuation 40–41  
     physiological effect 41  
     vertical distribution 42  
     vertical gradient 42–43  
*Thalassio-Syringodietalia filiformis* 505  
 thermobixeric climate 67–72  
 therophytes 189–190  
 thorn index 208–210  
 thorn woodlands 83, 90, 97, 338, 376, 616–617

- Thrinacion morrisii* 554  
*Thrinacion punctulatae* 418  
*Tournefortietum gnaphaloidis* 508  
*Trianthemo-Sesuvietalia* 455, 511  
*Trianthemo-Sesuvietum portulacastri* 454, 511.  
 511 t,  
*Trianthemo-Sesuvion* 510  
*Trinidadense* 323, 333–335  
 Trinidad Heights or Escambray 333–335  
*Trinidadicum* 323, 331–333  
 trisectorial distribution 248  
*Turnereto-Unioletum virgatae* 508  
*Turquinense* 349, 369–372  
*Typhetum domingensis* 450, 494 t,  
  
*Utricularietum foliosae* 474 t.  
*Utricularietum juncea* 474 t.  
*Utricularietum resupinatae* 474, 475 t.  
*Uveroënse* 349, 378–379  
  
*Vallisnerietum americanae* 477 t.,  
*Vallisnerietum neotropicalis* 448, 450 f, 477, 478 t.  
*Vallisnerion americanae* 477  
 vegetation map  
     Cajalbana range 318  
     Isle of Youth (Isla de la Juventud) 303  
     Zapata Peninsula 449  
 vegetation of coastal rock pavement 303, 454–455,  
     509–513  
 vegetation of sandy beaches 303, 453–454, 506–509  
 vegetation transects  
     Cajalbana range 319  
     Casilda Peninsula 332  
     Cerro de Yamanigüey 360  
     Cerro Pelo Malo 339  
     eastern Sierra del Rosario 322  
     eastern Zapate swamp 297  
     limestone terrace at Aguadores 383  
     limestone terrace at Siboney 383  
     Meseta El Toldo (Moa) 360  
     mogote of Pan de Guajaibon (Sierra del Rosario)  
     321  
     mogote of Pico Potrerillo (Sierra de Escambray)  
     334  
     mogotes of Rio Mogote (Sierra Maestra) 374  
     Monte Libano 367  
     Pico Turquino (Sierra Maestra) 371  
     Sierra de Iberia (Moa) 398  
     swamp of Laguna de Leche (Moron) 348  
     western Zapata swamp 296  
     Yunque de Baracoa 363  
 vertical vegetation belts 139  
 Vesey-FitzGerald's grassland classification 154–155  
 vicariance 201–203, 239, 249–253  
     ecological 250–252  
     epiphytes vs lianes 201–203  
     geographic 249–250, 251, 252, 253  
     hemicryptophytes vs chamaephytes 204–205  
     xerophilous vs hygrophilous epiphytes 203–204  
*Vinalense* 291, 311–316  
*Vrieseo-Bombacopsisidetum cubensis* 417  
*Vrieseo-Ceratopyxidetum verbenaceae* 553, 554 t,  
 vulnerability 256  
 water surplus 99–103  
     efficiency 99  
     gradient 100–103  
     index 99  
  
 Wegener's theory 257  
*Weinmannio-Cyrilletalia* 398, 640  
*Weinmannio-Cyrilletea* 639  
 western and central Sierra Maestra 349, 369–372  
 western Central Cuba 326  
 West Indian genera 285  
 white sandlands 299–303  
     of Isle of Youth 301–303  
     of western Pinar del Rio 300–301  
 white sand savannas 163  
  
 xerochimenic climate 64–67  
 xcromorphism (xcromorphy) 136, 137, 138, 143  
 xerotheric climates 60–61  
 xerotherotropical climate 63–64  
*Xyridi-Hypericetum fasciculati* 498  
*Xyridi-Paepalanthetum seslerioidis* 499  
  
*Yaguaramas-Cascajal plain* 330–331  
*Yaterense* 349, 364–368  
 Yucatan flora 269  
 Yunque de Baracoa 362–364  
  
*Zanthonyo-Burseretum simarubae* 199, 206, 415  
*Zanthonyo-Jacquinietum shaferi* 164, 165, 166,  
     199, 206  
*Zanthonyo-Reynosietum retusae* 602  
 Zapata peninsula (Zapatense) 294–298  
 zonality of savannas 107–108  
 zonal vegetation  
     bioclimatic classification 103  
     ecological parameters 106–108  
     predictability 93–99  
     profiles 77–88  
     types 62–75  
     vertical belts 106, 139  
 zonation  
     coastal rock pavement 454  
     coastal vegetation of SE-Cuba 568  
     fresh water communities 450  
     litorral sand vegetation 332  
     mangrove 451  
     sandy beaches 454  
*Zosteretea* 503

## REGISTER OF LATIN PLANT NAMES

occurring in the text, without the phytosociological tables  
(f=figure, m=map)

- Abrus precatorius* 232, 412  
*Acacia* 184, 267, 423, 443, 613  
– *belairioides* 343, 610  
– *bucherii* 358  
– *cowellii* 382, 423, 617  
– *cupeynsis* 348, 423, 617  
– *curbelotii* 348, 423, 617  
– *daemon* 218, 327, 604  
– *farnesiana* 232, 420, 617  
– *lutea* 423, 617  
– *macracantha* 383 f.  
– *polypyrogenes* 331  
– *roigii* 348, 423, 617  
– *seifriziana* 385, 423  
– *zapaensis* 294  
*Acalypha* 254  
– *huichinonii* 331  
– *mogotensis* 314  
*Acanthodesmos* 290  
*Achlaena pictostachya* 613  
*Acidocroton* 254, 286  
– *trichophyllus* 343  
*Acisantherina quadrata* 498  
*Acoelorraphe* 157, 163, 167, 303  
– *wrightii* 167, 301, 302, 431, 446, 448, 450 f., 519,  
577, 614  
*Acroporium pungens* 142  
*Acrostichum aureum* 194, 237, 451, 490, 517  
– *danaifolium* 194, 237, 446, 451, 517  
*Acrosanthus* 287, 350, 643  
– *minor* 360 f., 426, 597  
– *ovatus* 356  
– *revolutus* 636  
– *trachyphyllus* 401, 644  
*Actinostemon brachypodus* 552  
*Acunacanthus* 128, 130, 288, 310, 326  
– *tinifolius* 306, 317, 433, 530, 582  
*Adelia ricinella* 412, 420, 551 f.  
*Adenoua* 128 f., 268, 288  
– *cubensis* 596  
*Adiantopsis paupercula* 235  
*Adiantum* 194, 405  
– *capillus-veneris* 237  
– *cristatum* 235  
– *deltaeum* 235  
*Aechmea nudicaulis* 391  
*Agave* 187, 314, 376 f., 617  
– *albescens* 384 f., 385, 429, 620 f.  
– *anomala* 343  
– *brittoniana* 338 f., 605  
– *cajalbanensis* 316, 433, 582, 583, 599  
– *furcroydes* 233  
*Agave legrelliana* 325, 455, 617  
– *pendentata* 372 f., 403 f., 404, 461, 642  
– *shaserae* 368, 437, 588, 590  
– *tubulata* 313, 416 f., 417, 418, 552  
– *underwoodii* 351, 374 f., 383 f., 424, 617  
*Ageratum maritimum* 190  
*Albizia* 265  
*Alchornea latifolia* 392, 407 f., 408 f., 414 f., 530  
*Aldrovanda vesiculosa* 469, 473  
*Alibertia edulis* 231, 585  
*Allophylus cominia* 225, 412, 414 f.  
– *roigii* 218 m., 291, 293  
*Alsophila* 622, 626  
– *aquilina* 194, 237, 396, 401, 440  
– *aspera* 194, 399, 623  
– *myosuroides* 237, 630  
– *strigillosa* 234  
*Alvaradoa amorphoides* ssp. *psilophylla* 226, 541  
– *arborescens* 222, 350, 418  
*Alysicarpus vaginalis* 612  
*Alyssum bertoloni* 126  
*Anaioua corymbosa* 231, 585  
*Ambrosia elatior* 233  
*Ampelocera* 287  
*Amphiolanthus* 288, 326  
*Amphiolanthus arenarioides* 300  
*Anrysia* 267, 350  
– *balsamifera* 227, 412, 420, 539, 545  
– *diatrypa* 348, 545, 559  
– *elemifera* 230, 412, 539

- *lineata* 223, 248
- *polymorpha* 378
- *stromatophylla* 253
- Anacaona* 289
- Anacardium edule* 233, 442, 612
- Ananas comosus* 233
- Ancistranthus* 288, 310, 313
  - *harpochilooides* 313, 553
- Andira inermis* 231, 405, 412, 414 f, 539, 612
- Andrachne brittonii* 385
- Andropogon* 189, 435, 443, 444, 576, 613
  - *bicornis* 443
  - *breviflorus* 443
  - *caricosus* 612
  - *glomeratus* 496
  - *gracilis* 433, 576, 613, 615
  - *hirtiflorus* 444, 615
  - *multinervosus* 325, 444, 615
  - *nashianus* 439
  - *pertusus* 442, 612
  - *reedii* 443
  - *reinoldii* 436
  - *saccharoides* 612, 613
  - *tener* 615
  - *virgatus* 614
  - *virginicus* 233, 442, 443, 612, 613
- Anemia* 140, 194, 261, 273
  - *adiantifolia* 237
  - *cajalbanica* 234, 273 m, 327, 425, 433, 582
  - *coriacea* 234, 249, 253, 273 m, 317, 337, 353, 435, 587
    - - *ssp. moaensis* 273 m, 437
  - *cuneata* 234
  - *nipensis* 435
  - *wrightii* 237
- Angelonia pilosella* 444
- Anguria* 190
  - *pedata* 419
- Anisantherina hispidula* 443
- Annona* *bullata* 223, 323, 425, 595
  - *cascarilloides* 313, 417
  - *crystalensis* 356, 397 f
  - *glabra* 448, 519, 520 f
  - *muricata* 233
  - *sclerophylla* 401, 402 f, 596
- Anthurium* *venosum* 313, 416 f, 417
- Antillia* 266, 288
- Antirhea* 268
  - *abbreviata* 401, 426, 596
  - *aristata* 337
  - *lucida* 228
  - *minutifolia* 343
  - *ophiticola* 343
  - *orbicularis* 426
  - *radiata* 530
- Apassalus parvulus* 385
- Arcoa* 289
- Ardisia* *baracoensis* 362
- Argythamnia microphylla* 348
- Ariadne* 128, 129 f, 268, 288, 351
  - *shaferi* 437, 588, 596
  - - *ssp. shaferi* 401
- Aristida* 189, 444, 577, 614, 615
  - *brittonorum* 300
  - *fragilis* 301
  - *laevigata* 351, 437
  - *neglecta* 444, 614
  - *pradana* 351
  - *refracta* 433, 444, 576, 616
  - *vilifolia* 444, 576, 616
- Aristolochia* 190
  - *clematis* 425
  - *lindeniana* 351
  - *passiflorifolia* 614
- Arthrostylidium* 398
- *angustifolium* 364
- *capillifolium* 225, 446
- *cubense* 446
- *fimbriatum* 401
- *multispicatum* 404
- *pinifolium* 401
- Artocarpus* *incisa* 234
- Arundinella* *deppiana* 576
- Arundo* *donax* 446, 485
- Asciadium* 288
- Ascyrum* 642
- Asplenium* 194, 391, 405
  - *cristatum* 237
  - *cuneatum* 137
  - *dentatum* 237
  - *phyllitidis* 406
- Aster* 446, 488, 631
  - *burgessii* 631
  - *grisebachii* 498, 579, 615
- Atelcia* 269
  - *apetala* 415
  - *cubensis* 412
  - *gummifera* 420
  - *parvifoliola* 345
- Ateranmnus* *lucidus* 184, 228, 383 f, 405, 412, 420, 424, 545
- Ateranmnus* *recurvus* 342, 596, 597
- Auerodendron* 267, 286, 350
  - *acuminatum* 319, 553
  - *cubense* 424
  - *glaucescens* 368
  - *northropianum* 347
  - *reticulatum* 557
- Avicennia* 161, 451
  - *germinans* (*nitida*) 230, 296, 332 f, 451 f, 517
- Axonopus compressus* 576
- Ayenia cajalbanensis* 598

- *euphrasifolia* 444, 614
- *violacea* 568
- Azolla caroliniana* 194, 237, 469, 470
- Baccharis* 254, 509, 641
  - *halimifolia* 453
  - - *ssp. angustior* 491
  - *orientalis* 348
  - *scoparioides* 439, 587
  - *shaferi* 396, 401, 587
- Bacopa* 446
  - *longipes* 300
  - *monnierii* 490
- Bactris cubensis* 352, 353 m, 389, 396, 440, 529, 594, 626, 631
- Bambusa vulgaris* 188, 234, 446, 485
- Banara acunae* 319
  - *glaberrima* 334
  - *wilsonii* 348
- Banisteria pauciflora* 223, 414 f
- Barbieria pinnata* 585
- Barleriola* 287, 349
  - *saturejoides* 382 m
  - - *ssp. acunae* 382 m, 385
  - - *ssp. hirsuta* 382 m, 559
  - - *ssp. saturejoides* 382 m
- Batis maritima* 230, 451 f, 513, 516, 518 f
- Bauhinia* 187
  - *divaricata* 412 f
  - *jenningsii* 291
- Befaria* 267, 290
  - *cubensis* 254, 270, 300, 305 f, 432, 577, 579
- Begonia* 419
  - *cowellii* 378
  - *libanensis* 368
  - *lomensis* 404
  - *Behaimia* 160, 288, 324
  - *cubensis* 325, 605
  - *Beilschmiedia pendula* 621
  - *Belairia* 160, 288
    - *mucronata* 323, 415 f, 443, 613
    - *parvifolia* 378
    - *savannarum* 613
    - *ternata* 420
  - *Belloniu* 287, 349
  - *spinosa* 385, 415, 424, 559
  - *Belotia* 269
  - *Bembidium* 288, 362
  - *Berberis* 267
  - *tenuifolia* 419
  - *Bergia sessiliflora* 345-346
  - Bernardia dichotoma* 224, 419
  - Bisgoeppertia* 287
    - *scandens* 587
  - Blechnum* 194
    - *occidentale* 237
  - *serrulatum* 237
  - - *ssp. roigii* 576
  - *ruerckheimii* 400
  - Bletia florida* 401, 435
    - *purpurea* 228, 401, 435
    - *violacea* 593, 594 f
  - Bocconia frutescens* 230, 418
  - Bolbitis* 194
    - *nicotianifolia* 237
  - Bombacopsis* 416, 553
    - *cubensis* 187, 218, 275, 313, 314 f, 326, 412, 413 f, 416 f, 417, 418, 541, 550, 552 f
  - Bonania* 286, 350
    - *elliptica* 375
    - *microphylla* 348
  - Bonnetia* 230 m, 265
    - *cubensis* 222, 231 f, 358, 396, 626
  - Borreria eritrichoides* 604
    - *matanzasiae* 219
    - *ocimoides* 442, 443
    - *strumpfioides* 499
    - *thymocephala* 443
    - *verticillata* 612
  - Borrichia* 285, 454 f
    - *arborescens* 453, 455, 508, 511
    - *cubana* 219, 329, 508, 511
  - Botrychium* 194
    - *underwoodianum* 235
    - *virginianum* 237
  - Bourreria* 160, 184, 614
    - *badia* 599
    - *cassinifolia*
    - *divaricata* 595, 599
    - *microphylla* 595, 599
    - *mucronata* 314, 553
    - *pauciflora* 401
    - *polyneura* 313, 417, 553
    - *setoso-hispida* 599
    - *succulenta* 420, 545, 548, 551 f
    - *taylorii* 221
  - Brachiaria extensa* 223
  - Brasenia peltata* 476
    - *schreberi* 448, 476, 478
  - Broughtonia* 286
  - Brunellia comocladiifolia* 231, 395, 441 f, 621
  - Brunfelsia* 354
    - *clarensis* 330
    - *nitida* 326
  - Brya* 160, 286, 338
    - *chrysogonii* 337, 638
    - *depressa* 579
    - *ebenus* 224, 338 f, 415, 423, 433, 602
    - *hirsuta* 342
    - *microphylla* 621
    - *subinermis* 358, 398, 638
  - Buchenavia capitata* 308, 389, 390, 539

- Buchnera elongata* 443, 616  
*Bucida* 296, 297, 446  
 – *buceras* 324, 405, 414 f, 446, 519  
 – *ophiticola* 143, 327, 602, 605 f  
 – *palustris* 294, 298, 448, 450 f, 519, 523, 528  
 – *spinosa* 225, 420, 519  
 – *subinermis* 330, 446, 519, 523, 613  
*Bulbostylis* 475, 490  
 – *junciformis* 616  
 – *paradoxa* 230, 303, 613  
 – *setacea* 443, 490, 616  
 – *tenuifolia* 616  
*Bumelia* 161, 545  
 – *celastrina* 547  
 – *cubensis* 360, 437  
 – *glomerata* ssp. *horrida* 547  
 – (*Diphollis*) *jubilla* 350, 390, 406  
 – *retusa* 291  
 – (*Diphollis*) *salicifolia* 227  
*Bunchosia linearifolia* 424  
*Burmannia bicolor* 230  
*Bursera* 160, 265, 297  
 – *angustata* 599  
 – *glaucia* 350, 559  
 – *shaferi* 313, 553  
 – *simaruba* 184, 228 m, 268, 332 f, 383 f, 408 f, 412, 415, 416 f, 420, 421 f, 442, 539, 541, 546 f, 551 f, 571, 612  
*Buxus* 136, 140, 185, 222, 267, 350, 358, 595, 631, 643  
 – *aneura* 354  
 – *bahamensis* 225  
 – *baracoënsis* 361  
 – *crassifolia* 644  
 – *flaviramea* 125, 327, 604  
 – *foliosa* 636  
 – *gonoclada* 219, 327  
 – *heterophylla* 343  
 – *historica* 644  
 – *imbricata* 356  
 – *olivacea* 356  
 – *retusa* 644  
 – *shaferi* 644  
 – *wrightii* 433, 599  
*Byrsonima* 218, 254  
 – *biflora* 396, 401, 596, 623  
 – *bucherae* 358, 427  
 – *coriacea* 396  
 – *crassifolia* 167, 169, 185, 228, 431, 432, 444, 576, 613, 616  
 – × *hybrida* 579  
 – *minutifolia* 439, 596, 597  
 – *motembensis* 325  
 – *orientensis* 396, 397 f, 623  
 – *pinetorum* 301, 319, 432, 577, 579  
 – *verbascifolia* 230, 303, 577, 613  
 – *wrightiana* 431, 577, 578  
*Byttneria microphylla* 248, 419  
*Cabomba* 477  
 – *aquatica* 476  
 – *caroliniana* 226  
 – *piauhyensis* 448, 476  
*Caesalpinia bahamensis* 225, 420, 617  
 – *bonduc* 187, 232, 423, 548 f  
 – *coriaria* 617  
 – *crista* 187, 232, 420, 423  
 – *decapetala* 621  
 – *glandulosa* 385  
 – *glaucophylla* 344  
 – *guanensis* 314  
 – *hermeliæ* 343  
 – *hornet* 345  
 – *pauciflora* 420, 429, 560, 620  
 – *pinnata* 380, 429, 620  
 – *savannarum* 344, 613  
 – *subglaуca* 382, 617  
 – *vesicaria* 227, 420  
*Cakile maritima* 506  
*Calliandra* 267  
 – *colletioides* 350, 424  
 – *orientalis* 350  
*Callicarpa* 222, 354, 359  
 – *areolata* 362  
 – *buchereri* 375, 564, 568  
 – *cuneifolia* 590  
 – *ferruginea* 441  
 – *floccosa* 373  
 – *grisebachii* 375  
 – *hitchcockii* 225  
 – *lancifolia* 401, 437  
 – *nipensis* 437  
 – *ob lanceolata* 358, 437, 588  
 – *roigii* 291, 550  
 – *wrightii* 588  
*Callitricha occidentalis* 326  
*Calophyllum* 157, 161, 265, 296, 297, 626  
 – *antillanum* 184, 225, 405, 407 f, 408 f, 409 f, 414 f, 534, 630  
 – – ssp. *rivularis* 446, 630, 631  
 – *calaba* ssp. *pinetorum* 530, 583  
 – *jacquinii* 391  
 – *utile* 184, 389, 390 f, 396, 528, 623  
*Calycogonium* 140, 285, 350, 358, 396, 643  
 – *grisebachii* 593  
 – *microphyllum* 318  
 – *moanum* 596  
 – *plicatum* 364, 557  
 – *rosmarinifolium* 253, 254 m, 353, 401, 596, 597  
 – – ssp. *brachyphyllum* 254 m  
 – – ssp. *moanum* 254 m

- - *ssp. rosmarinifolium* 254 *m*
- *saxicola* 314
- Calycophyllum* 269, 296
- *candidissimum* 406, 534
- Calypogeia venezuelana* 140, 142
- Calyptranthes* 184, 217, 219, 222, 254, 265, 332, 350, 354, 358, 368, 626, 634, 643
- *calyprata* 626
- *cardiophylla* 626
- *caroli* 530
- *clementis* 630
- *dodecandra* 547
- *ferruginea* 630
- *gracilipes* 318
- *lomensis* 333
- *moaënsis* 636
- *monocarpa* 626, 636
- *pallens* 547
- *paradoxa* 366
- *peninsularis* 294
- *pinetorum* 577
- *pseudomoaënsis* 636
- *punctata* 626
- *toaensis* 636
- Calyptronoma* 287, 446, 621, 626
- *clementis* 350, 631
- - *ssp. orientensis* 389, 398, 632 *f*
- *dulcis* 305, 446, 630
- *intermedia* 630
- *microcarpa* 333, 630
- *orientensis* 626
- Calyptropsisidium sartorianum* 630
- Cameraria* 286
- *microphylla* 220, 341
- *retusa* 613
- Campylopus porphyrodetion* 142
- *saxatilis* 142
- Canavalia ekmanii* 412
- *maritima* 231, 453, 506
- Canella winterana* 226, 412, 415, 420, 421 *f*, 545
- Capparis* 420, 545, 617
- *cynophallophora* 230, 314 *f*, 420, 429, 548, 550, 551 *f*
- *ferruginea* 620
- *flexuosa* 230, 420, 429, 548, 619 *f* 620
- *grisebachii* 620
- Carapa* 269
- *guianensis* 184, 229, 361, 389, 390 *f*, 528
- Cardamine porphyrophylla* 334
- Caribaea* 288, 375, 380
- *litoralis* 385, 512
- Carica papaya* 187
- Carpodiptera* 262
- *cubensis* 415, 420
- Casabitoa* 289
- Casasia* 297
- *calophylla* 223, 323, 412, 414 *f*, 551 *f*
- *clusiifolia* 348, 507, 571
- - *var. hirsuta* 348
- Catasia jacquiniooides* 427, 597
- *nigrescens* 253 *m*, 353
- - *ssp. moaënsis* 253 *m*, 427
- - *ssp. nigrescens* 253 *m*
- Casearia* 160, 351, 587
- *bissei* 359, 437, 591
- *comocladiifolia* 385
- *crassinervis* 587
- *formosa* 331
- *guianensis* 414 *f*, 418
- *hirsuta* 412, 418, 539
- *moaënsis* 437, 587
- *ophiticola* 358, 587
- *spinescens* 539
- Cassia* 160, 265
- *aeschynomene* 612
- *bucherae* 358, 427
- *clarensis* 331
- *ekmaniana* 227
- *emarginata* 424
- *holguinensis* 343
- *insularis* 443
- *niqueroensis* 378
- *roiigii* 306
- *rotundifolia* 443
- *scleroxyla* 382
- *stenophylla* 617
- *turquinae* 404
- Cassipourea elliptica* 391
- Castela* 511, 547
- *calcicola* 249, 560
- *victorini* 380
- Castilloa elastica* 233
- Casuarina equisetifolia* 234
- Catalpa brevipes* 374, 378, 418
- *punctata* 412, 413 *f*, 541, 548, 550
- Catesbaea* 254, 268, 285, 511
- *gamboana* 345
- Catharanthus roseus* 234
- Catopsis* 392
- *nitida* 419
- Cattleyopsis lindenii* 190, 225
- Cecropia* 160, 161
- *peltata* 161, 230, 324, 339 *f*, 389, 405, 408 *f*, 421 *f*, 530
- Cedrela* 157, 161, 265
- *cubana* 534
- *odorata (mexicana)* 184, 231, 390 *f*, 405, 409 *f*, 412, 413 *f*, 421 *f*, 530, 534
- Ceiba* 157, 159, 160, 296, 410 *f*
- *pentandra* 184, 232, 268, 324, 405, 412, 413 *f*, 416 *f*, 442, 534, 612
- Celtis* 267

- *iguanea* 412, 553
- *trinervia* 184, 228, 418, 541, 554, 555
- Cenchrus distichophyllum* 499
- *tribuloides* 506
- Centella erecta* 490, 496, 501
- Ceratophyllum* 479
  - *denserum* 477
- Ceratopyxis* 266, 288, 310, 313, 553
  - *verbenacea* 313, 416 f, 417, 553
  - *verticillata* 416
- Cestrum bahamense* 225
  - *buxoides* 593
- Ceuthocarpus* 128, 268, 288, 358
  - *involucratus* 358 f, 427
- Chaetium* 288, 324
- Chaetocarpus* 593
  - *cordifolius* 356
  - *hunilis* 314
    - (*Mentenia*) *oblongatus* 591
- Chaetolepis* 267, 290
  - *cubensis* 295, 431, 498, 499, 577
- Chaunaesyce* 190
  - *biranensis* 346
  - *buxifolia* 511
  - *filicaulis* 343
  - *microclada* 382
  - *niqueroana* 378
  - *paredonensis* 347
  - *paucipila* 325
  - *pinarionia* 432
- Chaptalia* 354, 631, 634
  - *comptonioides* 378
  - *ekmanii* 316
  - *nipensis* 636
  - *shaferi* 636
  - *turquinensis* 372, 404, 462
- Chara* 448
- Chascotheca* 287
- Cheilanthes harrisii* 235, 400
- Cheiophyllum* 219, 287, 330, 444, 613
  - *macranthum* 341
  - *sphaerocarpum* 338
- Chimarrhis* 269
  - *cubensis* 631
- Chiococca alba* 231, 232 m, 412
- Chionanthus (Linociera) bumeloides* 541, 560
  - *domingensis* 396
- Chiome myrtifolia* 223
- Chloris cubensis* 613
  - *sagraeana* 451 f, 515
- Chlorophora* 265
  - *tinctoria* 230, 405
- Chrysobalanus* 297
  - *icaco* 228, 420, 448, 520
  - *pellocarpus* 432
- Chrysophyllum argenteum* 224, 405, 406
  - *clarens* 331
  - *oliviforme* 184, 414 f, 420, 442, 612
  - Chusquea abietifolia* 187, 372 f, 398, 404, 642
  - Ciceronia* 128, 288, 354
  - Cinnamodendron cubense* 369
  - Cinnamomum* 184
    - *elongatum* 225
    - (*Phoebe*) *montanum* 224, 389
    - *triplinerve* 407 f
  - Cionosycios* 287
  - Cissampelos pareira* 419
  - Cissus* 187
    - *dichroa* 314
    - *formosa* 227, 291, 412
    - *sicyoides* 412
    - *torreana* 325
  - Citharexylum caudatum* 227, 417
    - *matheanum* 419, 557
    - *ternatum* 370
  - Citrus* 234
  - Cladium* 296, 297, 348 f, 490
    - *jamaicense* 431, 446, 448, 450 f, 492, 496 f
    - *restioides* 398, 626
  - Clematis dioica* 412
  - Cleome gamboensis* 345
    - *tenuifolia* 345
  - Clerodendron calcicola* 314 f, 550
    - *nipense* 352, 435 f, 587
  - Clethra* 621, 622, 623
    - *cubensis* 222, 350, 440, 441 f, 623
  - Cleyera ekmanii* 369
    - *nimanimae* 350, 395, 622
  - Clidemia* 585
    - *capituliflora* 220, 333, 440
    - *neglecta* 432, 583
    - *strigillosa* 583
    - *wrightii* 333
  - Clitoria laurifolia* 432, 578
  - Clusia* 181, 350, 359, 643
    - *alainii* 217 m, 643
    - *brittonii* 314
    - *callosa* 400, 643
    - *minor* 585
    - *moaensis* 400, 643
    - *monocarpa* 400, 643
    - *nipensis* 354, 355 f, 401, 437, 590, 643
    - *rosea* 230, 389, 390, 391 f
    - *tetragyna* 392, 394 f, 623
  - Cneorum* 256, 264 m
    - *pulverulentum* 264 m
    - *tricoccum* 264 m
    - *trimerum* 264 m, 351, 369, 370
  - Cnidoscolus matosii* 385
  - Coccoloba* 188, 222, 297, 350, 351, 358, 508, 509, 595, 643
    - *acuna* 360, 427, 596

- *armata* 420, 599
- *coriacea* 599
- *costata* 397 f, 623
- *cowellii* 337, 610
- *diversifolia* 420, 545
- *geniculata* 220, 273, 275 m, 337, 605
- *microphylla* 599
- *nervosa* 644
- *nipensis* 343, 427, 596
- *oligantha* 437, 644
- *pallida* 326, 599
- *praestans* 596
- *reflexa* 401, 597
- *retusa* 323, 530
- *shaferi* 427, 587
- *uvifera* 230, 332 f, 453, 571, 572
- *wrightii* 333
- Coccothrinax* 107, 157, 162, 163, 170, 222, 267, 285, 297, 337, 346, 350, 365 m, 375, 418, 429, 545, 557, 559, 577
- *acunana* 365 m, 372
- *alexandri* 187, 365 m, 380, 429, 564, 566, 569 f
- - *ssp. nitida* 365 m
- *bermudezii* 365 m
- *borchidiana* 217 m, 219, 317 m, 329 f, 547, 562
- *camagüeyana* 220, 317 m, 340 f, 341, 610, 614
- *clarensis* 317 m, 337, 605, 614
- *crinita* 248, 317 m
- - *ssp. brevicrinitis* 317 m
- *cupularis* 560, 561
- *ekmanii* 565
- *elegans* 365 m, 374 f, 418, 557
- *fagildei* 365 m, 382
- *fragrans* 385, 424, 547
- *garciana* 187, 343, 365 m, 444, 605, 610 f, 614
- *guantanamensis* 365 m, 385
- *gundlachii* 351, 365 m, 379
- *hiorami* 187, 365 m, 380, 429, 564
- *leonis* 365 m
- *litoralis* 317, 326, 420, 547
- *microphylla* 365 m, 385, 424
- *miraguama* 187, 325, 425, 431, 443, 561, 614
- - *ssp. arenicola* 168, 576
- - *ssp. roseocarpa* 125, 327, 425, 604, 605 f
- *moaensis* 365 m, 443 f, 444, 598
- *munizii* 187, 365 m, 380, 424, 429, 564, 565, 566 f
- *muricata* 317 m, 348
- *nipensis* 365 m, 444
- *orientalis* 365 m, 434 f, 435, 437, 587
- *pauciflora* 346
- *pseudorigida* 187, 220, 317 m, 340 f, 341, 605, 610, 614
- *salvatoris* 337, 347
- *savannarum* 346
- *saxicola* 365 m, 378
- *trinitensis* 419
- *victorini* 365 m, 378
- *yunquensis* 364, 365 m, 557
- *yuraguana* 187, 316, 317 m, 433, 582
- - *ssp. moaensis* 437
- Cocos nucifera* 234
- Coeloneurum* 289
- Cololejeunea* 641
- Colpothrinax* 157, 163, 167, 168, 261, 295, 302
- *wrightii* 167, 218, 295, 300 m, 301, 431, 445 f, 557, 614
- Colubrina* 267
- *arborescens* 421 f, 547
- *cubensis* 547, 562
- *elliptica (reclinata)* 383 f, 417, 418, 424, 547, 548, 551 f, 621
- *obtusata* 385
- Columnea tincta* 359, 391 f, 528
- Commelinia elegans* 572
- Comocladia* 187
- *dentata* 324, 423, 583
- *mollifolia* 326
- *platyphylla* 547 f
- Comparettia falcata* 398
- Conocarpus* 297, 451, 455
- *erecta* 231, 332 f, 383 f, 446, 451 f, 455 f, 490, 511, 517, 571
- Conostegia xalapensis* 432, 576, 585
- Consolea* 285, 617
- *macracantha* 187, 222, 375, 380, 429, 430 f, 564, 617
- *millspaughii* 617, 619 f
- Copernicia* 107, 121, 157, 161, 162, 170, 219, 265, 324 m, 328 m, 331, 337, 344, 346, 416, 443
- *baileyana* 185, 221, 324 m, 325, 343, 415 f, 443, 613
- *brittonorum* 328 m
- × *burretiana* 324 m, 337, 344, 443
- *clarkii* 443
- *cowellii* 187, 220 f, 341, 444, 605, 610, 614
- *curbeloi* 337
- *curtissii* 187, 327, 328 m, 583
- *fallaensis (fallaense)* 187, 324 m
- *gigas* 185, 324 m, 337, 343, 443, 613
- Copernicia glabrescens* 302, 320 f, 328 m, 443, 519
- *hospita* 170, 324 m, 343, 443, 599, 605, 613
- - *ssp. clarenensis* 444
- *humicola* 378, 443
- *longiglossa* 345
- *macroglossa* 187, 324 m, 325, 443, 599, 614, 615 f
- *molineti* 337, 344, 443
- *oxycalyx* 348
- *pauciflora* 614
- - *var. ramosissima* 561
- *ramosissima* 604
- *rigida* 221 f, m, 337, 443, 613
- *roigii* 443

- *x shaferi* 337
- *x sueroana* 221, 324 m, 337, 343, 415, 443, 613
- *x textilis* 324 m, 337, 343, 344, 415, 443, 613
- *x vespertilionum* 187, 221, 324 m, 337, 343, 443, 613
- *yarey* 187, 337, 444, 614
- Cordia* 136, 160, 184, 222, 359, 429, 559
  - *alba* (*dentata*) 442
  - *alliodora* 418, 541
  - *angiocarpa* 326
  - *bahamensis* 347
  - *baracoënsis* 361
  - *brittonii* 424, 568
  - *collococca* 405, 409 f, 415, 539
  - *corallicola* 378, 568
  - *curbeloi* 348, 563
  - *duartei* 359, 511
  - *dumosa* 378
  - *galeottiana* 548, 560
  - *gerascanthus* 228, 405, 412, 413 f, 414 f, 415, 421 f, 442, 539, 541, 554, 555, 612
  - *globosa* ssp. *humilis* 418, 420, 424, 545, 560, 568
  - *grisebachii* 337
  - *holguinensis* 610
  - *intricata* 331
  - *leptoclada* 375, 424, 568
  - *leucosebestena* 375, 383 f, 424, 568
  - *linnaei* 585
  - *longipedunculata* 369, 400
  - *nitida* 415
  - *pedunculosa* 593
  - *pulverulenta* 424, 563
  - *sebestena* 548, 560
  - *sulcata* 390 f, 391, 529
  - *taylorii* 375
  - *toaensis* 593, 594
  - *valenzuelana* 319
- Coronopus didymus* 190
- Corymborkhis* 189
- Coussarea* 269
- Crescentia cujete* 405, 446, 541, 542 f
- *mirabilis* 220, 347–348
- Crinum* 189
  - *americanum* 484, 494, 496
  - *oliganthum* 326, 484, 496
- Crocodilus rhombifer* 298 f
- Crossopetalum rhacoma* 229, 545, 571
- ekmanii*
- *spathulifolium* 369
- *ternifolium* 596
- Crotalaria urbaniana* 345
- Croton* 185, 254, 268, 559, 566, 617
  - *acunae* 343, 605
  - *betulinus* 568
  - *bispinosus* 248
  - *borhidii* 440
- *brittonianus* 248
- *camagüeyanus* 337, 605, 610
- *cerinus* 577, 615
- *corallicola* 378, 568
- *craspedotrichus* 432, 577, 615
- *excisus* 385, 568
- *heteropleurus* 337, 605
- *holguinensis* 343, 610
- *litoralis* 223, 375, 568
- *ssp. *rugelianus** 329, 562, 568
- *lucidus* 225, 324, 420, 424, 547
- *micradenus* 375, 568
- *monogynus* 437
- *munizii* 375, 568
- *myricifolius* 337, 375, 424, 568
- *nephrophyllus* 337, 605
- *nummulariifolius* 189, 444, 614
- *pachyrhachis* 356
- *rosmarinoides* 225, 375, 424, 568
- *sagraeanus* 613
- *stenophyllus* 568
- *tenuiramis* 568
- *tropidophyllus* 362
- *yunquensis* 364, 557
- Cryptorhiza* 289
- Cryptostegia grandiflora* 234
- Cubacroton* 288, 369
- *mæstrensis* 369
- Cubanola* 266, 287
- Cubanthus* 287
- Cuervea integrifolia* 417, 553
- Cupania americana* 184, 389, 390 f, 406
- *glabra* 228, 390 f, 414 f, 421 f, 534
- *macrophylla* 268, 409 f
- Cuphea lobelioides* 631
- Cuphea melanium* 442
- *parsonisia* 442
- *pseudosilene* 499, 577
- Curatella americana* 231, 432, 576, 613
- Cyathea* 621, 622
  - *araneosa* 194, 234, 390, 391, 441 f
  - *arborea* 194, 235, 236 m, f, 237, 392, 394 f, 399, 623, 640
  - *balanocarpa* 392, 399, 640
  - *cubensis* 234, 392
  - *furfuracea* 235 m
  - *insignis* 640
  - *minor* 399, 626
  - (*Nephelea*) *pubescens* 235, 640
  - *tenera* 194, 235
- Cybianthus* 290
- Cylindropuntia hystrix* 429, 617
- Cynanchum* 190
  - *ophiticola* 425, 545
  - *orientensis* 595
  - *salinarum* 451

- *Savannarum* 613
- Cynodon* 488
  - *dactylon* 233
- Cynometra cubensis* 530
- Cyperus* 189, 234, 442
  - *acicularis* 446
  - Cyperus alternifolius* 374, 446
    - *articulatus* 446, 487
    - *diffusus* 487
    - *flavus* 442
    - *giganteus* 448, 492, 494
    - *haspan* 442, 612
    - *heterophyllus* 486
    - *iria* 442
    - *ligularis* 442
    - *surinamensis* 442, 486, 612
  - Cyrilla* 621, 626
    - *cubensis* 135, 222, 352, 401 f, 437
    - – *ssp. nipensis* 396
    - *nipensis* 437, 626, 640
    - *racemiflora* 135, 228, 392, 394 f, 399, 441 f, 582, 622, 631
- Cytisus* 140
- Dalbergia cubensis* 350, 529
- Dalbergia ecastaphyllum* 446, 516, 521
  - *monetaria* 521
- Dalechampia scandens* 412
- Danaea* 400
- Daphnopsis alainii* 334, 557
  - *guacacao* 319
  - *oblongifolia* 220, 337, 605
- Dasytropis* 128, 288, 354, 643
- Datura stramonium* 233
- Daucus carota* 234
- Davilla rugosa* 432
- Deherainia* 269
- Delonix regia* 234
- Dendrocereus* 288, 297
  - *nudiflorus* 187, 249, 331, 375, 377 f, 424, 429, 545, 546 f, 561, 564
- Dendrocoussinia* 290
- Dendropanax arboreus* 405, 407 f, 418, 419, 530
  - *cuneifolius* 187, 530, 630
  - *nervosus* 356
- Dendropemon* 187
- Dendrophthora* 187
- Dennstaedtia* 194, 399, 405, 640
  - *cicutaria* 237
- Desmanthus* 262
- Desmodium barbatum* 432
- Dichaea* 190
  - *hystricina* 225, 398
- Dichanthium (Andropogon) caricosum* 172
- Dichromena ciliata* 442
  - *colorata* 446
- Dichrostachys* 121, 338
  - *cinerea* 134, 162, 234, 324
- Didymopanax morototoni* 187, 231, 335 f, 389, 390 f, 397 f, 405, 406, 626
- Digitaria decumbens* 172
- Dilomilis* 190
  - *elata* 398
  - *montana* 224, 392
  - *oligophylla* 398, 401
- Dinema* 287
  - *cubincola* 398
- Diodia maritima* 506
  - *rigida* 444
  - *teres* 444
- Dioscorea* 190
- Diospyros* 268
  - *anisandra* 227, 291
  - *crassinervis* 412, 413 f, 550
  - *grisebachii* 223, 424, 545
  - *halesioides* 415
  - *tetrasperma* 291
- Dipholis* 265
  - *cubensis* 224, 397 f, 427
  - *ekmanii* 528
  - *jubilla* 184, 374 f, 389, 396, 405, 528
  - *neglecta* 530
    - *salicifolia* 405, 534
- Diplazium* 194, 405
  - *arboreum* 237
  - *unilobum* 235
  - *urbanii* 400
- Distichlis* 296
  - *spicata* 507, 515
- Distictis gnaphalantha* 425
  - *lactiflora* 425
  - *rhynchocarpa* 425
- Ditta* 286
  - *maestrensis* 395, 640
  - *myricoides* 224, 248
- Dodonaea* 267
- Doerpfeldia* 268, 288, 336
  - *cubensis* 337, 348
- Dorstenia* 219, 265, 368
  - *lanei* 338
  - *roccana* 335
  - *roigii* 314
- Doryopteris pedata* 237
- Dracaena* 256
  - *cubensis* 135, 187, 188, 222, 262, 651 f, 352, 359, 360 f, 437, 591, 592 f, 597
  - *reflexa* 262
  - *draco* 262
- Drepanolejeunea* 641
  - *suchaeta* 142
- Drosera brevifolia* 226
  - *intermedia* 233, 498

- *rotundifolia* 233
- *tenella* 230
- Drymaria glandulosa* 385
- Dryopteris* 405
  - *asterothrix* 237
  - *crypta* 234
  - *dentata* 237
  - *hastata* 235
  - *incisa* 235
  - *patens* 519
  - *sancta* 237
  - *serra* 519
- Dryptes alba* 407f, 408f
  - *lateriflora* 409f, 413f
  - *mucronata* 541
- Duranta arida* 385
  - *fletcheriana* 223, 400
- Echinochlaena* 262
- Echinodorus* 487, 488
- Ehretia tinifolia* 184
- Eichhornia* 469, 482
  - *azurea* 189, 469
  - *crassipes* 469
  - *diversifolia* 469
  - *heterosperma* 469
- Ekmania* 288, 364
  - *lepidota* 557
- Ekmanianthe* 287
  - *actinophylla* 313, 417, 553
- Ekmaniocharis* 289
- Ekmanochloa* 128, 288, 351
  - *aristata* 401
- Elaeagia* 269
- Elaeodendron attenuatum* 420
- Elaphoglossum* 195, 395, 400, 639
  - (*Hymenodium*) *crinitum* 391
  - *eggersii* 235
  - *firmum* 237
  - *hirtum* 237
  - *martinicense* 235
  - *revolutum* 237
- Eleocharis* 297, 348f, 494
  - *alveolata* 492
  - *articulata* 487
  - *capillacea* 498
  - *caribaea* 501
  - *cellulosa* 448, 487, 489f, 491
  - *interstincta* 448, 487, 488, 492
  - *minutissima* 492
  - *oligantha* 492
  - *parvula* 233
- Elephantopus* 498
  - *pratinus* 501
- Elleanthus linifolius* 229, 398
- Enallagma latifolia* 414f
- Encopella* 288
  - *tenuifolia* 501
- Encyclia* 405
  - *atropurpurea* 401
- Enhydra sessilis* 446
- Eosanthe* 128, 266, 288
  - *cubensis* 217m
- Epaltes mattfeldii* 190
- Epidendrum* 190, 228, 350, 405
  - *globosum* 398
  - *ionosmum* 230
- Equisetum giganteum* 237
- Eragrostis* 190
  - *cubensis* 614
  - *salzmannii* 230, 515
- Erianthus giganteus* 226, 492, 496
- Erigeron bellidoides* 631
  - *capillipes* 334, 631
  - *cuneifolius* 613
  - *paucilobus* 631
  - *taylori* 379
  - *thrincoides* 631
- Eriocaulon* 217, 218, 267, 290, 295, 301, 303
  - *arenicola* 499
  - *dioicum* 300
  - *echinospermum* 330
  - *fuliginosum* 498, 499, 501
  - *ovoideum* 499
  - *sphaerocephalum* 300
- Eriochloa ekmanii* 337
  - *setosa* 326, 613, 614
- Eriosema* 269
  - *crinitum* 432
- Erithalis fruticosa* 228, 424, 455, 507, 571
  - *vacciniifolia* 511, 512
- Ernodea* 285
  - *maritima* 453, 507
- Erythrina* 267
  - *cubensis* 417, 552
  - *leptopoda* 356
- Erythroxylon* 184, 420, 545
  - *areolatum* 225, 418, 420
  - *baracoense* 364, 557
  - *clarensse* 419, 557
  - *confusum* 225, 446, 522
  - *coriaceum* 596
  - *flavicans* 356
  - *havanense* 223
  - *horridum* 610
  - *longipes* 223
  - *minutifolium* 338f, 424, 583, 595, 599, 610
  - *pedicellare* 596, 597
  - *roigii* 218, 291
  - *rotundifolium* 420, 551f
- Espadaea* 266, 288
  - *amoena* 323

- Euchorium* 288, 310, 313  
*Eugenia* 160, 184, 185, 217, 219, 222, 254, 265,  
 268, 295, 354, 358, 369, 396, 559, 631  
 – *aceitillo*, 378  
 – *amblyophylla* 382  
 – *anafensis* 327, 554, 555  
 – *axillaris* 226, 412, 413 f, 420  
 – *banderensis* 597  
 – *bayatensis* 366  
 – *cajalbanica* 599  
 – *camarioca* 325 m, 425, 599, 602  
 – *clarensis* 219, 338, 605  
 – *cowellii* 424, 568  
 – *cristata* 553  
 – *cycloidea* 588  
 – *duplicata* 554, 555  
 – *excisa* 366  
 – *farameoides* 326, 585  
 – *galeata* 313, 417, 553  
 – *guanensis* 314  
 – *ignota* 308  
 – *itephylla* 382  
 – *laeteviridis* 370, 631  
 – *lomensis* 404  
 – *lucayana* 347  
 – *maestrensis* 370, 404, 631  
 – *malevolens* (*buxifolia*) 226, 324, 412, 415, 419,  
 420, 421 f, 545, 548, 551 f  
 – *melanadenia* 605  
 – – *ssp. santayana* 341  
 – *mensuraensis* 401  
 – *mollifolia* 329  
 – *naguana* 370  
 – *oligantha* 631  
 – *papayoensis* 623  
 – *peninsularis* 378  
 – *piedraensis* 354  
 – *pinetorum* 437, 587  
 – *pterooclada* 348  
 – *punicifolia* 585  
 – *ramonae* 587  
 – *ramonianana* 582  
 – *rhombea* 412  
 – *rigidifolia* 319, 433, 582, 583, 599, 602  
 – *rosariensis* 433  
 – *sauvalllei* 425, 602  
 – *serrei* 554  
 – *shaferi* 341  
 – *subdisticha* 219, 338  
 – *victorini* 302  
*Euleria* 288, 324, 332  
 – *tetramera* 557  
*Eupatorina* 289  
*Eupatorium* 160, 185, 222, 350, 362, 369, 396  
 – *ayapanoides* 404, 587  
 – (*Grisebachianthus*) *carsticola* 351, 557  
 – *dalea* 441 f, 623  
 – *grandiceps* 440, 593  
 – *grisebachianum* 316  
 – *helianthemooides* 375  
 – *lantanifolium* 439, 587  
 – *littoralis* 348  
 – *maestrense* 370  
 – *minutifolium* 360  
 – *nipense* 437  
 – *paucibracteatum* 370, 399, 402, 441 f  
 – *polystictum* 587  
 – *rhexioides* 398  
 – *turquinense* 641  
*Euphorbia* 267  
 – *sect. Euphorbiadendron* 395  
*Euphorbia cubensis* 317, 644  
 – *helena* 222, 317, 350, 437 f, 440, 587, 596, 598  
 – *lactea* 234  
 – *marchii* 566  
 – *munizii* 358, 359 f  
 – *podocarpifolia* 342 f, 343, 354, 401, 596  
 – *pulcherrima* 233  
*Evolvulus* 614  
 – *arbuscula* 189, 225  
 – *sericeus* 432, 444  
*Exostema* 268  
 – *caribaeum* 420, 551 f  
 – *longiflorum* 631, 633 f  
 – *purpureum* 427  
 – *shaferi* 361  
 – *spinulosum* 383, 385, 424  
 – *stenophyllum* 354, 636  
  
*Fadyenia hookeri* 235  
*Fagara* 265  
*Furanea occidentalis* 231, 405, 409 f, 416 f  
*Feddeaa* 128, 135, 187, 288, 358, 643  
 – *cubensis* 359, 401, 644  
*Feea* 639  
*Ficus* 181, 187, 405  
 – *aurea* 226, 418, 420, 442 f  
 – *bertero* 374, 390 f, 405, 406, 528  
 – *combsii* 323  
 – *crassinervis* 412, 413 f  
 – *havanensis* 339 f  
 – *jacquinifolia* 225, 418, 545  
 – *laevigata* 415  
 – *subscabrida* 223, 323, 409 f, 414 f, 446  
 – *wrightii* 389, 440, 528  
*Fimbristylis* 488, 490, 613  
 – *annua* 232, 443, 446  
 – *autumnalis* 613  
 – *complanata* 501  
 – *spadicea* 233  
 – *spathacea* 232, 446, 513  
 – *squarrosa* 231–232

- Fissidens weizii* 400  
*Flaveria linearis* 455, 511  
*Forchhamneria* 269  
 – *emarginata* 360  
 – *trifoliata* 227, 291  
*Forestiera* 267  
 – *rhamnifolia* 225, 551 f  
 – *segregata* 621  
*Forsteronia corymbosa* 407 f, 419  
*Fraxinus* 269  
 – *caroliniana* ssp. *cubensis* 226, 270, 295, 448, 522, 528  
*Freziera* 265, 621  
 – *conocarpa* 356  
 – *grisebachii* 621, 622  
*Fuertesia* 289  
*Fuertesiella* 287, 349  
*Fuirena umbellata* 484, 492  
  
*Galactia acunana* 291  
 – *cuneata* 375  
 – *galactioides* 599  
 – *herrdurensis* 308  
 – *isopoda* 295  
 – *jenningsii* 295  
 – *maisiana* 385  
 – *revoluta* 596  
 – *roundata* 348  
 – *rudolphiooides* 435, 587  
*Garcinia ophiitica* 388, 596  
 – *polyneura* 596  
 – *revoluta* 596  
 – *ruscifolia* 596  
 – *serpentinii* 583  
*Garrya* 267  
 – *fadyenii* 224, 333, 392, 394 f, 399, 419, 440, 441 f, 557, 640  
*Gastrococos* 288, 324  
 – *crispia* 185, 223, 323 m, 444  
*Gauussia* 287, 416, 553  
 – *princeps* 185, 313, 319, 416, 417, 553  
*Gayoides crispum* 425  
*Genipa americana* 442  
*Genista* 140  
*Geophila minutiflora* 385  
*Gesneria* 188, 189, 222, 285, 350, 362, 364, 368, 369, 557  
 – *brevifolia* 321  
 – *celsioides* 313, 416 f, 417  
 – *clarensis* 220, 333  
 – *cubensis* 359, 557  
 – *duchartreoides* 587  
 – *ferruginea* 317, 583  
 – *gibberosa* 351, 362, 557  
 – *heterochroa* 557  
 – *humilis* 631  
 – *libanensis* 367  
 – – ssp. *libanensis* 367 m  
 – – ssp. *lopezii* 367 m  
 – *nipensis* 354, 355 f, 437  
 – *norlindii* 587  
 – *pachyclada* 356, 587  
 – *purpurascens* 362, 367 m, 557  
 – – var. *purpurascens* 367 m  
 – – var. *yumuriensis* 364 f, 367 m  
 – *salicifolia* 557  
 – *viridiflora* 220, 557, 621  
 – – ssp. *colorata* 622  
 – *yumuriensis* 362  
*Ginoria americana* 446, 631  
 – *arborea* 638  
 – *ginoriooides* 631  
 – *microphylla* 337  
 – *montana* 638  
*Gleichenia* 194, 395, 399, 640  
 – *bifida* 194, 622  
 – *flexuosa* 398, 622  
 – *leonis* 398  
 – *palmata* 622  
 – *pectinata* 622  
*Glyricidia sepium* 324  
*Gnaphalium attenuatum* 190  
*Gochnatia* 136, 185, 350, 351, 429  
 – *calcicola* 375  
 – *cowellii* 337, 610  
 – *cubensis* 354  
 – *ekmanii* 300, 308, 321  
 – *intertexta* 317, 599  
 – *maisiana* 351  
 – *mantuensis* 306  
 – *microcephala* 617  
 – *parvifolia* 343  
 – *recurva* 439  
 – *sagracana* 329, 550  
*Goerziella* 288, 291  
*Goetzea* 266, 287  
*Gomidesia lindeniana* 392, 440  
*Gomphrena decumbens* 442  
*Gossypianthus brittonii* 616  
*Gossypiospermum praecox* 230, 415, 541, 554  
*Gouania lupuloides* 407 f  
 – *ekmani* 314  
*Graffenreidea rufescens* 395  
*Grammitis* 639  
*Grimmeodendron* 285  
 – *eglandulosum* 225, 383 f, 424, 545, 547, 548, 550  
*Grisebachianthus* 288  
*Guajacum officinale* 184, 230, 429 f, 545, 550, 559, 564, 617, 620  
 – *sanctum* 184, 227, 545, 550  
*Guapira (Torrubia)* 545  
 – *bracei* 347

- *cajubancensis* 317
- *leonis* 327
- *longifolia* 547
- *peninsularis* 294
- Guarea* 269
  - *guaranda* (*guara*) 184, 231, 389, 390 f, 405, 406, 408 f
  - *Guatteria* 265
    - *blainvillii* 333, 621, 622
    - *muralessii* 350, 397 f, 593, 623
  - *Guazuma ulmifolia* 405, 414 f, 534, 612
  - *Guenthera* 184, 265, 268, 351, 429, 559
    - *amblyophylla* 314, 541
  - *calcicola* 417, 553
  - *calyptata* 223, 323, 435, 440, 582, 599
  - *camagüeyensis* 220, 341, 605
  - *clarensis* 219, 338, 605
  - *combsii* 446, 522
  - *coxiiana* 375, 564
  - *crassipes* 437, 588
  - *cucuensis* 375, 563
  - *echinodendron* 599
  - *elliptica* 541
  - *ferruginea* 352, 437, 439, 440, 588
  - *languinosa* 220, 348
  - *monocarpa* 435, 588
  - *mutisii* 220
  - *nervosa* 334
  - *rígida* 564, 599
  - *roigiana* 219, 338, 605
  - *sciaphila* 596
  - *shaferi* 343 f, 588, 596
  - *undulata* 217 m, 219, 329
  - *urbaniana* 333
  - *valenzuelana* 530, 582
- *Guilleminia (Gossypianthus) heterophylla* 604
- Gundlachia* 285, 634
  - *apiculata* 638
  - *cubana* 440, 638
  - *lindeniana* 638
- *Guizmania erythrolepis* 392
- *lingulata* 391, 392, 419
- *monostachya* 391, 392, 394 f, 419
- Gymninda latifolia* 227, 420, 545
  - *orbicularis* 385
- Gymnogramma cubensis* 234
- Gynerium sagittatum* 188, 230, 446, 485, 486 f
- Gyrotaenia* 286
- Habenaria* 189
- Haematoxylum campechianum* 228, 617
- Haemianthus* 286
- Haitia* 289
- Halodule beaudetii* 503
- *wrightii* 503, 505
- Halophila aschersonii* 505
- *baillonii* 505
- Hamelia patens* 418
- Harnackia* 128, 187, 249, 268, 273, 288, 317, 354
  - *bisecta* 249 m, 401
- Harpalyce* 140, 256, 268, 350, 595
  - *cubensis* 317, 326, 327, 604
  - *macrocarpa* 338
  - *maisiana* 385
- Harrisia* 187, 297
  - *fernowii* 424, 429, 620
  - *taetra* 291, 292 f, 561
  - *taylori* 221, 564
- Hebestigma* 288
  - *cubense* 184, 541, 554, 555
- Hedychium coronarium* 255
- Hedyosmum* 265, 278
  - *crassifolium* 278 m, 398, 626
  - *cubense* 278 m, 370, 400, 640
  - *grisebachii* 220, 248, 278 m, 333
  - *nutans* 621, 640
  - *subintegrum* 278 m
- Heliconia caribaea* 188, 390, 391, 529
- Helicteres nípensis* 317
  - *semitriloba* 425, 620
  - *trapezifolia* 317, 599
- Helictia* 267
  - *glaucescens* 418
- Heliotropium antillanum* 291
- *curassavicum* 491, 515
- *humifusum* 455, 511
  - *myriophyllum* 347
  - *ternatum* 568
- Hemianthus* 446
- Hemianthus palmata* 237
- Hemitelia horrida* 194, 390, 391, 394 f, 621
- Hemithrinax* 250, 268, 336, 418
  - *rivularis* 360 f, 426
  - *savannarum* 426
- Henleophytum* 288, 324
- Hennozia* 266, 288, 336
  - *brittonii* 337, 620
  - *myrtifolia* 605, 610
- Henriettea cuabae* 361
  - *ekmanii* 370, 400, 640
  - *granularis* 319
  - *patrisiana* 585, 630
- Heptanthus* 189, 246, 247 m, 288, 631, 634
  - *brevispes* 247 m, 318
  - *cochlearifolius* 247 m 579
  - *cordifolius* 247 m, 435, 636
  - *lobatus* 247 m, f
  - *ranunculoides* 247 m, 295, 582
  - *shaferi* 247 m, 636
  - *yumuriensis* 247 m, 364
- Hernandia cubensis* 368
- Herodotia* 289

- Herpyza* 288, 295  
 – *grandiflora* 218, 432, 498  
*Heterotrichum umbellatum* 441 f  
*Hibiscus* 296, 297  
 – *elatus* 157, 224, 414 f, 446, 448, 534  
 – *urbani* 301  
*Hildegardia cubensis* 184, 221, 343, 541  
*Hillia* 529  
 – *parasitica* 187, 392 f  
 – *tetrandra* 391  
*Hippomane mancinella* 420, 545, 548, 551 f, 571  
*Hirtella* 269  
 – *americana* 626  
*Hohenbergia penduliflora* 190, 339 f, 413 f, 417  
*Homalopetalum* 287  
*Hottea* 289  
*Hydrocotyle* 488  
 – *oligantha* 366–367  
 – *umbellata* 476, 478, 479  
*Hydrolaeta nigricaulis* 501  
*Hyeronima* 184  
 – *crassistipula* 308  
 – *havanensis* 326  
 – *nipensis* 396, 397 f, 440, 593, 623  
*Hylocereus undatus* 187, 405  
*Hymenaea courbaril* 231  
 – *torrei* 221  
*Hymenocallis* 189  
*Hymenodium crinitum* 237  
*Hymenophyllum* 195, 395, 400, 639  
 – *abruptum* 401  
 – *axillare* 235  
 – *contortum* 237  
 – *lanatum* 235  
 – *polyanthos* 237  
 – *sericeum* 237  
*Hyparrhenia rufa* 442  
*Hypelate* 285  
 – *trifoliata* 226, 415, 420, 539, 541, 551 f  
*Hyperbaena acutifolia* 335  
 – *columbica* 306, 582  
 – *cubensis* 350, 418  
 – *domingensis* 229  
 – *racemosa* 551 f  
*Hypericum* 189, 642  
 – *incurvum* 308  
 – *ophiticola* 338  
 – *perforatum* 234  
 – *styphelioides* 223, 248, 432, 576  
 – – *ssp. moaënsis* 439  
 – – *ssp. styphelioides* 432, 438 f  
*Hypnum polypterum* 400, 641  
*Hyptis* 189  
 – *armillata* 325  
 – *cubensis* 308  
 – *minutifolia* 576  
 – *pedalipes* 498, 579, 615  
 – *rivularis* 331  
*Ichnanthus mayarensis* 585, 611  
 – *nemorosus* 626  
 – *pallens* 391  
*Ilex* 136, 140, 185, 222, 254, 267, 641, 643  
 – *berteroii* 396, 400, 643  
 – *cassine* 522  
 – *clementis* 419  
 – *cristaleensis* 356  
 – *eoae* 356  
 – *hypaneura* 400, 643  
 – *macfadyenii* 135, 395, 440, 623  
 – *moana* 135  
 – *nannophylla* 404, 641  
 – *nunezii* 372, 402, 641  
 – *paucinervis* 356  
 – *shaferi* 400, 643  
 – *subavenia* 356, 643  
 – *ternatifolia* 318  
 – *turquiniensis* 372 f, 402, 404, 641  
 – *victorini* 400  
 – *wrightii* 364  
*Illicium cubense* 222, 350, 401  
*Imperata brasiliensis* 444  
 – *contracta* 612  
*Indigofera tinctoria* 234  
*Inga* 265  
*Ionopsis turricularioides* 190  
*Ipomoea* 190, 217, 232, 511  
 – *alba* 506  
 – *batatas* 233  
 – *carolina* ssp. *ophiticola* 425  
 – *flavo-purpurea* 331  
 – *pes-caprae* 231, 332 f, 453, 506  
*stoloniferae* 507  
*Isachne leersioides* 398, 626  
*Iseertia haenkeana* 626  
*Isidorea* 266, 277 m, 287, 349  
 – *leonis* 385  
 – *polyneura* 367  
 – *rheedioidea* 385  
*Isocarpha glabrata* 347  
*Isoetes cubana* 194, 234  
*Isopterygium micans* 142  
*Ixora ferrea*  
*Jacaima* 290  
*Jacaranda* 254  
 – *arborea* 222, 401, 427, 587  
 – *coerulea* 225, 539  
 – *cowellii* 220, 249, 273, 275 m, 337, 605  
*Jacmaia* 290  
*Jacquemontia jamaicensis* 227, 420, 425, 511  
*Jacquinia* 268, 429

- *aculeata* 420
- *acunana* 358
- *berterii* 385, 424, 559
- *brevifolia* 545
- *brunnescens* 433, 583, 602
- *curtissii* 308
- *keyensis* 545
- *maisiana* 385, 424, 559
- *moana* 360 f, 597
- *robusta* 342
- *roigii* 437, 591
- *sessiliflora* 356
- *shaferi* 337, 605, 607 f
- *stenophylla* 420
- *verticillaris* 382
- *yunquensis* 364
- Jambosa vulgaris* 374 f
- Jatropha urupifolia* 337
- Juglans* 267
- Juniperus* 267
  - *lucayana* 225
  - *saxicola* 372, 404, 641
- Jussiaea (Ludwigia) decurrens* 631
- *peruviana* 233
- Justicia* 189
  - *agria* 378
  - *maestrensis* 369, 382
  - *stenophylla* 341
  - *tomentulosa* 343
- Kalmiella* 254, 267, 290, 578
  - *aggregata* 218, 303, 431
  - *ericoides* 295, 299
  - *simulata* 218, 303, 431
- Karwinskyia* 267
  - *bicolor* 369
  - *oblongifolia* 219, 338, 605
  - *orbiculata* 337, 605
  - *potrerilloana* 334, 419
  - *rocana* 333
- Kodalyvodendron* 128, 268, 288, 358
  - *cubense* 185, 359, 426–427
- Koehneola* 128, 135, 266, 288, 354
- Krokia* 128, 288
- Krugiodendron* 254
  - *ferreum* 228, 412, 420, 545, 547, 548
- Lachnanthes tinctoria* 498
- Lachnocalon* 267
  - *anceps* 295
  - *cubense* 330
  - *ekmanii* 295, 498
- Lachnorhiza* 288, 498
- Laelia lyonsii* 391
- Lagenocarpus cubensis* 356
- Lagerstroemia indica* 234
- Lagezia* 265, 286
  - *wrightiana* 319, 530
- Laguncularia* 451
  - *racemosa* 332 f, 451 f, 516, 518
- Lanium hiorami* 398
- Lantana* 429, 559, 566
  - *arida* 568
  - *involuta* 420, 424, 545, 560, 568
  - *parvifolia* 375, 568
  - *strigosa* 553
- Lantanopsis* 287
- Laplacea* 254, 265, 621, 643
  - *angustifolia* 184, 394 f, 621, 622
- *benitoënsis* 400, 644
- *cristalensis* 644
- *ekmanii* 350
- *modënsis* 358, 400, 626, 644
- *urbanii* 369, 621, 622
- *wrightii* 350
- Lasiaciis divaricata* 230
- Lasianthus lanceolatus* 395
- Lasiocroton* 286
  - *bahamensis* 225, 602, 620
  - *gracilis* 382
- Lemairocereus (Ritterocereus) hystrix* 348, 375, 424–425
- Lemna* 469
  - *minima* 469
  - *perpusilla* 469, 470
  - *trisulca* 233
- Lepanthes* 190, 222, 350, 419, 641
  - *acunae* 372, 400
  - *blepharantha* 400, 404
  - *ekmanii* 372, 400, 640 f
  - *fractiflexa* 400
  - *fulva* 400
  - *pergracilis* 372, 404
  - *turquinoensis* 372, 404
- Lepanthopsis microlepanthes* 404
- Leptocereus* 286, 416, 552
  - *assurgens* 314, 417, 553
  - *ekmanii* 314, 553
  - *leoni* 219, 554, 555
  - *prostratus* 314, 416 f
  - *sylvestris* 331, 378
  - *wrightii* 219, 329
- Leptocoryphium lanatum* 432, 444, 576, 614
- Leptogonium* 289
- Leptoscyphus cuneifolius* 142
- Lepturidium* 217, 288, 295, 302
- Lescaillea* 128, 187, 249, 268, 273, 288, 310, 316, 317
  - *equisetiformis* 249 m, 433, 583, 599
- Leskeodon andicola* 142
- Leucaena leucocephala* 231, 332 f, 571, 576
- Leucobryum giganteum* 142

- *polakowskyi* 142
- Leucocroton* 128, 136, 140, 222, 268, 287, 350, 354, 595, 634
- *acunae* 636
- *ekmani* 361
- *havanensis* 125, 325, 425, 602
- *longibracteatus* 623
- *microphyllus* 418, 423, 559
- *moaënsis* 595 f, 644
- *moncadae* 223, 325, 602
- *obovatus* 356, 636
- *revolutus* 317, 599
- *saxicola* 342–343
- *stenophyllus* 636
- *wrightii* 397 f, 623
- Licaria jamaicensis* 409 f
- Limnobium laevigatum* 505
- Limnocharis flava* 230, 448, 484, 494
- Lindernia alternifolia* 300
- Lindsaya* 140, 194
  - *cubensis* 234
  - *stricta* 237
- Linociera (Chionanthus)* 297
  - *buneliooides* 413 f
  - *cubensis* 360 f
  - *domingensis* 440
  - *ligustrina* 420
- Linodendron* 265, 268, 288
  - *aronifolium* 439, 588
  - *venosum* 248, 317
- Lisianthus glandulosus* 440
  - *silenefolius* 613
- Litachne pauciflora* 626
- Lithophila muscoides* 455, 510
- Lobelia* 254
  - *cacuminis* 372, 402, 641
  - *cubana* 553
- Lolium temulentum* 234
- Lonchocarpus domingensis* 225, 405, 446, 534
  - *latifolius* 405, 534
- Lophosoria quadripinnata* 194, 394 f, 399, 621
- Luehea* 269
  - *speciosa* 405, 534
- Lunania cubensis* 419
  - *elongata* 332
- Lycium carolinianum* 451
  - *tweedianum* 516
- Lycopodium* 194, 195
  - *cernuum* 237
  - *clavatum* 237, 400
  - *funiforme* 195, 391
  - *montanum* 235, 400
  - *serratum* 400
  - *taxifolium* 237, 400
- Lygodium cubense* 195, 234, 432
  - *volubile* 398
- Lyonia* 185, 222, 248, 254, 267, 350, 396, 623, 641
  - *affinis* 623
  - *calycosa* 400, 402, 623, 641
  - *ekmani* 295, 308
  - *elliptica* 248
  - *glandulosa* 401, 440, 587
  - *leonis* 372 f
  - *longipes* 593
  - *lucida* 226
  - *macrophylla* 435, 440, 587, 593
  - *maëstrensis* 623
  - *myrsinifolia* 587
  - *myrtilloides* 295, 305, 432, 578
  - *nipensis* 435
  - *obovata* 593
  - *turquini* 641
  - *vaccinioides* 295, 431, 577
- Lysiloma* 296, 297, 450 f
  - *bahamense* 227, 332 f, 420, 421 f, 446, 541, 551 f
  - *latisiliqua* 225
- Machaerina cubensis* 427
- Machaonia* 254, 325 m
  - *acunae* 308
  - *dumosa* 317, 433, 583, 599, 602
  - *havanensis* 249, 325 m, 326
  - - *ssp. orientalis* 325 m
  - *microphylla* 220, 325 m
  - *minutifolia* 325 m, 605, 610
  - *nipensis* 325 m, 596
  - - *ssp. moaensis* 325 m
  - *pauciflora* 308
  - *pubescens* 304, 325 m
  - *subinermis* 217 m, 325 m, 337, 605
  - - *ssp. armata* 325 m
  - *urbaniana* 325 m, 382
  - *urbinoi* 325 m, 343, 605
- Macrocarpaea* 287
- Macromitrium* 641
  - *harrisii* 400
  - *jamaicense* 400
  - *schwanckeanum* 400
- Macroptilium lathyroides* 442
- Magnolia* 265, 621
  - *cacuminicola* 621
  - *cubensis* 184, 220, 248, 333 m, 350, 392, 394 f, 396, 621, 622
  - - *ssp. acunae* 333 m, 622
  - - *ssp. cubensis* 333 m
  - - *ssp. baracoënsis* 333 m
- Malaxis* 395
- Mallotonia* 454 f
  - (*Tournefortia*) *gnaphaloides* 332 f, 453, 508
- Malpighia* 254, 415, 545
  - *acunana* 374, 419, 557

- *cniodes* 439, 595
- *cubensis* 420
- *horrida* 433
- *nummulariifolia* 325, 425, 595
- *pallens* 560
- *roigiana* 313, 417, 553
- *wrightiana* 319, 530
- Mamillaria* 314
- *prolifera* 314
- Manekia* 289
- Manilkara* 161
  - *albescens* 529
  - *grisebachii* 530
  - *jaimiqui* 420, 522
  - *ssp. wrightiana* 415, 523
- Marattia* 194
  - *alata* 237, 640
- Marcgravia* 187
  - *evenia* 391, 398
  - *reciflora* 419
- Margaritaria nobilis* 407 f., 409 f., 530
  - *scandens* 223
- Margaritopsis* 287, 349
- Marsilia* 482
  - *polycarpa* 194, 483
  - *quadrifolia* 483
- Mastichodendron foetidissimum* 184, 226, 405, 407 f., 412, 534
- Matayba apetala* (= *oppositifolia*) 407 f., 530
  - *domingensis* 394 f., 396, 397 f., 621
  - *oppositifolia* 184, 530
- Matelea bayatensis* 367
  - *nipensis* 354
- Mattfeldia* 289
- Maxillaria* 190, 350, 391
- Maxonia apifolia* 235
- Mayaca fluviatilis* 475, 476
  - *wrightii* 448, 492
- Maytenus* 267
  - *buxifolia* 185, 225, 252 m., 415, 424
    - *ssp. buxifolia* 252 m.
    - *ssp. cajalbanica* 252 m.
    - *ssp. cochlearifolia* 252 m., 380, 559, 620
    - *ssp. monticola* 252 m.
    - *ssp. serpentini* 252 m., 559, 360 f.
  - *lineata* 318
  - *saxicola* 369
- Mecranium* 440
  - *amygdalinum* 621
  - *haemanthum* 223
- Medicago lupulina* 234
- Megalopanax* 288, 324
  - *rex* 187, 344
- Meliosma oppositifolia* 220, 330
- Melocactus* 375, 381, 425, 559, 617
  - *actinacanthus* 338, 605
- *acunai* 380, 381 fm., 429, 564, 620
- *ssp. lagunaensis* 385
- *borbidi* 381 m., 424, 564, 620
- *evae* 381 m., 385
- *harlowii* 381 m., 424, 564, 620
- *holguinensis* 343, 605
- *matanzanus* 219, 327 f., 604
- *nagyii* 379, 381 m.
- *radozii* 361
- Melochia manducata* 308
  - *savannarum* 577, 615
- Melothria* 190
- Meriania leucantha* ssp. *nana* 335, 394 f., 621
- Merremia cissoides* 407 f.
- Mesechites minima* 325
  - *rosea* 425
- Mesosetum loliiforme* 576, 615
- Metacoriopsis paula* 400, 641
- Metopium* 285, 350
  - *brownei* 420, 519, 545, 547
  - *toxiferum* 226, 545, 547
- Mettenia (Chaetocarpus) oblongata* 440
- Mettenothamnium reptans* 142, 400, 641
- Miconia* 222, 265, 268, 350, 358, 369, 440, 643
  - *acunae* 623
  - *albescens* 622
  - *ancistrophora* 332
  - *androsaemifolia* 295, 577
  - *cerasiflora* 440, 593
  - *cubensis* 248, 333, 622
  - *dodecandra* 390, 391
  - *elata* 390 f., 391, 529
  - *ibaguensis* 305, 576, 583, 585
  - *javorkaeana* 360, 427
  - *lanata* 229
  - *lenticellata* 598
  - *nystroemii* 400
  - *obovata* 593
  - *perelegans* 308
  - *prasina* 583
  - *punctata* 392, 394 f.
  - *remotiflora* 623
  - *splendens* 305, 576, 585
  - *tetrandra* 397 f.
  - *turquinensis* 372 f., 400, 640
  - *wilsonii* 332
- Micranthemum rotundatum* 300
- Microcycus* 256, 261, 288, 295, 308, 313
  - *calocoma* 187, 188, 217, 270, 306 m., 307 f., 417, 553
- Microlepanthes* 641
- Microlepis* 639
- Micromeria bucheri* 404, 641
  - *suborbicularis* 385
- Micropholis* 161
  - *polit* 397 f.

- Mikania hastata* 227  
*Mimosa apuleia* 318  
 – *catalinae* 318  
 – *fagaracantha* 337  
 – *pigra* 232, 496  
 – *pubica* 232  
*Mitracarpus acunae* 372 f, 404, 641, 642  
 – *delpauperatus* 499  
 – *glabrescens* 433, 582  
 – *squarrosus* 614  
*Miranthes ottonis* 630  
*Mniachloa* 288  
*Moacroton* 128, 130, 249, 268, 273, 274 m, 288, 337, 351, 595  
 – *ekmanii* 359, 396, 626, 274 m  
 – *grandifolius* 274 m  
 – *lanceolatus* 274 m, 317, 342, 353, 596  
 – – *ssp. lanceolatus* 274 m  
 – – *ssp. revolutus* 274 m  
 – *leonis* 274 m, 358, 427  
 – *revolutus* 219, 327, 604  
 – *tetramerus* 274 m, 358  
 – *trigonocarpus* 274 m, f, 316, 317, 599  
*Mollugo* 190  
 – *brevipes* 301  
 – *cuneifolia* 568  
 – *enneandra* 301  
 – *pinosia* 218, 302  
*Mommsenia* 289  
*Morinda moenaensis* 398, 401  
 – *royoc* 420, 425, 511, 572  
*Mouriri emarginata* 588  
 – *valenzuelana* 319–321  
*Mozartia* 128, 287, 626, 634  
 – *gundlachii* 397 f, 623  
 – *maëstrensis* 631  
 – *manacalensis* 631  
*Muntingia* 160  
 – *calabura* 161  
*Musa paradisiaca* 234  
*Myrcia* 254, 626  
 – *gundlachii* 396  
 – *retivenia* 401  
 – *valenzuelana* 319, 530  
*Myrica* 160, 254, 265, 267, 296, 297, 358, 623, 641  
 – *cacuminis* 370, 402, 641  
 – *cerifera* 521, 576  
 – *punctata* 350, 441 f, 623  
 – *shaferi* 143, 350, 401, 593, 644  
*Myriophyllum pinnatum* 477  
 – *sparsiflorum* 448, 476, 477  
 – *verticillatum* 476  
*Myroxylon balsamum* 233  
*Myrsine coriacea* 184, 392, 397 f, 441 f, 622, 623  
 – *cubana* 446, 450 f, 519, 522  
 – *guianensis* 269, 519  
 – *microphylla* 370, 399, 640  
*Myrtus* 267  
 – *acunae* 597  
 – *anomala* 220, 337, 595  
 – *cabanasensis* 337, 568  
 – *crenulata* 331  
 – *del-riscoi* 356  
 – *elliptica* 595  
 – *matanzasia* 125, 327  
 – *micarensis* 356, 587  
 – *nummularioides* 385  
 – *oonophyla* 375, 559, 564  
 – *ophiticola* 437, 587  
 – *sagraei* 319  
*Najas* 448, 479  
 – *guadalupensis (microdon)* 477, 503  
 – *marina* 233, 477, 503  
 – *minor* 450  
 – *wrightiana* 503  
*Narvalina* 289  
*Nashia* 268, 286  
 – *armata* 382  
 – *cayensis* 220, 347  
 – *variifolia* 341  
*Nasturtium portoricense* 490  
*Nectandra antillana* 414 f  
 – *coriacea* 227, 408 f, 409 f, 420, 421 f  
 – *reticularis* 370, 399, 640  
*Neea shaferi* 559, 568  
*Nelumbo lutea* 189, 226, 448, 478, 481  
*Neo-abrottia* 289  
*Neobesseyea cubensis* 190, 343  
*Neobracea* 251 m, 273, 286, 350  
 – *acunaiana* 251 m  
 – *angustifolia* 249, 251 m, 293  
 – *ekmanii* 251 m, 359  
 – *howardii* 219, 251 m, 332, 557  
 – *mariiana* 251 m, 382  
 – *susannina* 251 m, 374, 419, 557  
 – *valenzuelana* 125, 130 f, 223, 249, 251 m, 272, 323, 425, 427, 439, 595, 599  
*Neobuchia* 289  
*Neocogniauxia*  
*Neolaugeria* 285  
*Neomacfadya* 269  
 – *podopogon* 227  
*Neomazaea* 128, 268, 289, 310, 316  
 – *phialanthoides* 143, 217, 433, 582, 599  
*Neoregnellia* 287  
 – *cubensis* 248  
*Neorudolphia* 290  
*Neothymopsis* 286  
*Neo-Urbania* 287  
*Nephelea* 621

- Nephrolepis* 194  
 – *biserrata* 446  
 – *rivularis* 237  
*Nodocarpaea* 289, 295, 304  
 – *radicans* 499  
*Nostoc commune* 451 f  
*Notodon* 268, 289, 595  
 – *cayensis* 347  
 – *savannarum* 337, 605, 610  
*Nuphar* 481  
 – *advena* 448  
 – *luteum* ssp., *macrophyllum* 478  
*Nymphaea* 297, 476, 481, 482  
 – *amazonum* 229  
 – *ampla* 189, 448, 478  
 – *blanda* 230, 298, 478  
 – *odorata* 233, 448  
*Nymphoides* 476  
 – *aurea* 483  
 – *grayanum* 189, 448, 478  
 – *humboldtianum* 478
- Ochroma* 269  
 – *pyramidalis* 389, 390 f, 528  
*Ocotea* 161, 184, 262, 265, 621  
 – *acunaiana* 621, 622  
 – *bucherii* 396  
 – *cuneata* 224, 333, 392, 394 f, 406, 621, 622  
 – *ekmanii* 220, 248, 333, 621, 622, 640  
 – *floribunda* 392, 419, 622  
 – *leucoxylon* 225, 389, 392, 396, 621  
 – *moaënsis* 623  
 – *wrightii* 333  
*Odontocline* 290  
*Odontosoria* 140, 195, 395, 398, 399  
 – *aculeata* 235  
 – *uncinella* 235  
 – *wrightiana* 318, 432, 583  
*Oleandra articulata* 235  
*Oldenlandia capillipes* 631  
 – *polyphylla* 634  
*Oliganthes* 262  
*Olyra latifolia* 161, 230  
*Omphalea commutata* 270, 277 m, 385  
 – *diandra* 229, 391  
 – *hypoleuca* 313, 416 f, 417, 553  
 – *trichotoma* 545, 547 f  
*Oncidium* 405  
 – *floridanum* 226  
*Ophioglossum* 194  
 – *vulgatum* 237  
*Oplonia* 251, 262, 263 m, 285, 415  
 – *acunae* 251 m  
 – *cubensis* 251 m, 437, 596, 597  
 – *moana* 251 m  
 – *multigemma* 251, 343
- *nannophylla* 218, 251 m, 326, 599, 602  
 – *polyce*, 251 m, 348, 424, 559  
 – *purpurascens* 251 m, 417, 552  
 – *spinosa* 225  
 – *tetrastricha* 223, 251 m, 323, 560  
*Opuntia* 511  
 – *dillenii* 227, 329 f, 383 f, 429, 455, 511, 548, 551 f, 572, 617  
 – *militaris* 385, 429, 620  
*Ormosia krugii* 391  
*Osmunda* 194  
 – *cinnamomea* 519  
 – *regalis* 237, 519  
*Ossaea* 140, 185, 222, 265, 268, 350, 359, 368, 369, 396, 643  
 – *acunae* 437  
 – *cristalensis* 356  
 – *elliptica* 361  
 – *ferruginea* 440  
 – *heterotricha* 364  
 – *maestrensis* 640  
 – *micarensis* 356  
 – *munizii* 587  
 – *navasensis* 440, 587  
 – *ottoschmidtii* 220, 333, 392, 621  
 – *pauciflora* 437, 583, 587  
 – *pinetorum* 356, 593  
 – *pseudopinetorum* 593, 594  
 – *rufescens* 593, 594  
 – *turquinensis* 370, 621  
 – *vazquezii* 594  
 – *yumuriensis* 364  
*Ottoschmidtia* 254, 287.  
 – *dorsiventralis* 218, 317, 327, 425, 604  
*Ottoschulzia* 286  
*Ouratea ilicifolia* 185  
 – *striata* 435, 587  
*Oxandra* 265, 285  
 – *lanceolata* 184, 224, 390, 391, 405, 406, 407 f, 409 f  
 – *laurifolia* 225, 333, 390, 391, 394 f, 405, 406, 529
- Pachyanthus* 268, 287, 643  
 – *angustifolius* 295, 305, 630  
 – *clementis* 335  
 – *cubensis* 295, 576, 577, 583  
 – *longifolius* 218, 303, 431, 578  
 – *lunanus* 332  
 – *mantuensis* 301, 577  
 – *moaënsis* 427  
 – *monocephalus* 356  
 – *poiretii* 217, 305, 432, 576, 579, 583  
 – *reticulatus* 435, 626  
 – *tetramerus* 319, 530  
 – *wrightii* 295, 576, 577, 579  
*Paepalanthus* 218, 267

- *alsinoides* 295, 499, 578
- *brittonii* 360 f, 427, 435
- *lamarckii* 499
- *pungens* 401
- *riparius* 401, 626
- *seslerioides* 295, 499, 578
- Palicourea domingensis* 414 f
- Paltonium lanceolatum* 237
- Panicum* 189, 297
  - *aciculare* 432, 440, 613, 614
  - *adspersum* 442, 612
  - *aquaticum* 487
  - *boliviense* 442
  - *caespitosum* 442, 612
  - *chrysopsidifolium* 443, 614
  - *condensum* 225, 496
  - *dichotomiflorum* 442
  - *fusiforme* 440
  - *geminatum* 612
  - *ghisbreghtii* 612
  - *glutinosum* 161
  - *lacustre* 487, 492
  - *lancearium* 496
  - *laxum* 501
  - *longiligulatum* 431
  - *maximum* 161, 442
  - *parvifolium* 492
  - *pilosum* 612
  - *reptans* 442
  - *scoparium* 225, 440
  - *stenodes* 613
  - *tenerum* 492
  - *virgatum* 233, 496
  - – var. *cubense* 494
- Panicum wrightianum* 498
- Paratheria* 269
- Parathesis cubana* 576
- Paspalidium paludivagum* 487
- Paspalum* 189, 190, 488
  - *alainii* 440
  - *breve* 437
  - *distactum* 507
  - *distichum* 442, 612
  - *distortum* 613
  - *divaricatum* 612
  - *edmondi* 338
  - *fimbriatum* 442, 612
  - *giganteum* 496
  - *motemboënse* 219, 338
  - *notatum* 442, 612
  - *plicatum* 612
  - *vaginatum* 231, 446, 506
- Passiflora* 190
  - *cubensis* 425, 595
  - *cuprea* 425
  - *multiflora* 412, 425
- *santiagana* 380, 425
- *suberosa* 412, 425
- Pavonia calcicola* 331
- *fruticosa* 161
- *intermixta* 577
- *spicata* 516
- Pectis* 455, 511
- *cubensis* 325
- *floribunda* 612
- *havanensis* 219, 327
- *leptocephala* 190
- *pinosia* 302
- *ritlandi* 331
- Pedinopetalum* 289
- Peltapteris peltata* 195, 237
- Peltophorum adnatum* 225, 405, 420, 442, 551 f, 612
- Peltostigma* 290
- Penelopeia* 289
- Pennisetum purpureum* 446, 486
- Peperomia* 227, 350, 395, 418, 419
  - *galioides* 404
  - *grisebachiana* 404
  - *guanensis* 314
  - *hirta* 223
  - *nummularia* 308
  - *verticillata* 417
- Pera* 184
  - *ekmanii* 396, 397 f, 626
  - *microcarpa* 369
- Peratanthe* 287, 349
  - *cubensis* 369, 402, 642
- Persea* 184, 265, 621
  - *anomala* 222, 369, 399, 640
  - *similis* 394 f, 402, 622, 640
- Petitia* 285
  - *urbanii* 277 m, 385
- Phaius* 395
  - *tankervilliae* 189, 231, 529, 631
- Phania* 268, 289, 310, 326
  - *cajalbanica* 189, 316, 433, 582
  - *matricarioides* 218, 275, 585
- Pharus glaber* 161, 230
- Pheidonocarpa* 287
  - *cubensis* 385
- Phenax* 262
- Phialanthus* 266, 268, 286, 595
  - *bissei* 293
  - *glaberrimus* 644
  - *macrostemon* 362, 644
  - *myrtilloides* 347
  - *parvifolius* 362
  - *rigidus* 317, 599
- Phidiasia* 128, 268, 289, 351, 352 m, 643
  - *lindavii* 352 m, 354
- Philodendron clementis* 405

- *krebsii* 405, 419, 446
- *lacerum* 339 f, 374 f, 405, 416 f, 419
- *urbanianum* 313, 416 f, 417
- Philonotis sphaericarpa* 142
- Philoxerus vermicularis* 231, 506, 515
- Phinnea* 269
- Phoradendron* 187
- *zapatanum* 294
- Phragmites australis* 492
- Phyla stoechadifolia* 490
- Phyllacanthus* 268, 289, 310, 316, 318
  - *grisebachianus* 217 m
- Phyllanthus* 128, 136, 185, 189, 222, 254, 265, 268, 272, 273, 295, 351, 354, 631
  - *chamaecristoides* 596, 636
  - *chrysaeus* 358, 360 f, 426, 596, 597
  - *comosus* 360, 426, 596, 597
  - *discolor* 326, 599
  - *epiphyllanthus* ssp. *dilatatus* 557
  - *erythrinus* 253
  - *estrellensis* 354
  - *heliotropus* 499, 501
  - *juncceus* 295, 432, 578
  - *myrtilloides* 352, 353, 596
  - – ssp. *erythrinus* 437
  - *orbicularis* 223, 249, 272, 273 m, 323, 424, 595, 599
  - *pachystylus* 593
  - *phlebocarpus* 354
  - *sagraeanus* 316
  - *selbyi* 303
  - *shaferti* 590
- Phyllogonium fulgens* 400, 641
- Phyllocladia* 128, 289, 310
  - *coronata* 319, 599
- Phyllostylon brasiliense* 343, 415, 617
- Picardaea* 287, 349
- Picramnia pentandra* 226, 412, 554
  - *reticulata* 530
- Picrodendron* 254, 286
  - *macrocarpum* 225, 227 m, 420, 424, 545, 547, 551 f
- Pictetia arborescens* 348
  - *marginata* 337, 605
- Pieris* 267, 290
  - *cubensis* 295
  - *Pilea* 190, 217, 219, 222, 265, 350, 364, 398, 418, 419, 556, 557, 626, 631, 634
  - *affinis* 314
  - *bullata* 319
  - *cacuminum* 333
  - *carnosa* 378
  - *cellulosa* 395
  - *clarana* 333, 395
  - *clementis* 335, 395
  - *cowelli* 378
- *güirana* 314
- *hemisphaerica* 314
- *micromerifolia* 404
- *pulchra* 557
- *sevillensis* 379
- *simplex* 314
- *sumideroensis* 314
- *uninervis* 374, 557
- Pilosocereus* 187, 297, 339 f, 416, 617
- *brooksianus* 221, 331, 418, 429, 551 f, 566 f, 617, 620
- *millspaughii* 225
- *robinii* 329, 545, 560, 617
- Pilotrichella flexilis* 400, 641
- Pinguicula albida* 300, 301 m
  - *benedicta* 301 m
  - *filifolia* 301 m, 497
  - *jackii* 301 m, 304
  - *lignicola* 301 m 636 f
- Pinillosia* 266, 287
- Pinosia* 289
- Pinus* 140, 246, 267, 269
  - *caribaea* 107, 162, 218, 293, 304, 305, 308, 310, 318, 319, 320 f, 431, 432, 433, 576, 579, 583, 616
  - *cubensis* 107, 352 m, 357, 359, 360–361, 368, 373, 396, 434 f, 435 f, 437, 439, 440, 587, 593, 594
  - *cubensis* × *maestrensis* 373
  - *elliottii* 576
  - *maestrensis* 107, 217 m, 369, 373, 440, 441 f, 622, 623
  - *occidentalis* 440, 622
  - *tropicalis* 107, 167, 168, 218, 295, 301, 302, 304, 308 m, 310, 431, 432, 576, 577, 615, 616
- Piper cojimaranum* 329
  - *guanahacabibense* 291
  - *holguinianum* 342
  - Piper perditum* 321
    - *sumideranum* 314
    - (*Potomorphe*) *umbellatum* 418
  - Piptocoma* 287
  - Piriqueta* 262
  - Piscidia* 541
    - *cubensis* 325, 595
    - *havanensis* 551 f
    - *piscipula* 415
  - Pisonia aculeata* 412, 414 f, 420
    - *petiolaris* 306, 599
    - *rotundata* 415
  - Pistia stratiotes* 469, 471
  - Pitcairnia cubensis* 351
  - Pithecellobium* 184, 265, 297, 613
    - *arboreum* 228
    - *cubense* 308, 405, 412, 418, 420, 442, 539, 585, 612
    - *hystrix* 225, 424
    - *lentiscifolium* 420, 541, 546 f

- *mäestrense* 369
- *millspaughii* 347
- *mucronatum* 617
- *nipense* 354
- *ovovale* 585
- *oppositifolium* 385, 617
- *savannarum* 343
- *trinitense* 333
- Pithecoctenium echinatum* 405, 412
- Pityrogramma* 194
  - *calomelanos* 237
  - *sulphurea* 235
  - *tartarea* 237
- Plagiochila steyermarkii* 140, 142
- Platygyne* 187, 250 *m*, 289
  - *dentata* 250 *m*, 369
  - *hexandra* 223, 250 *m*, 323, 412
  - *leonis* 250 *m*, 362
  - *obovata* 250 *m*, 359, 398
  - *parvifolia* 250 *m*, 273, 337, 605
  - *triandra* 250 *m*, 437, 590
  - *volubilis* 250 *m*
- Pleodendron* 287
- Plethadenia* 287, 349
- Pleurothallis* 190, 222, 350, 368, 392, 400, 419, 641
  - *gelida* 404
  - *obliquipetala* 404
  - *racemiflora* 227
  - *testifolia* 404
  - *velaticaulis* 404
  - *wrightii* 404
- Plinia* 254, 268, 553, 631
  - *dermatodes* 316, 433, 599
  - *ramosissima* 301, 366
  - *recurvata* 319
  - *ruberinervis* 314
- Pluchea* 488
  - rosea* 501
- Plumeria* 187, 188, 268, 424, 545
  - *cubensis* 325, 339 *f*
  - *clusioides* 435
  - *ekmanii* 418
  - *einmarginata (obtusa)* 225, 314 *f*, 417, 420, 547, 552
  - *filifolia* 418
  - *keyensis* 325, 547
  - *lanata* 350, 351 *f*, 564, 620
  - *nipensis* 587
  - *sericifolia* 547, 583
  - *stenophylla* 350
- Podocarpus* 140, 246, 254, 265, 351, 370, 621
  - *aristulatus* 623
  - *ekmanii* 222, 352, 396, 397 *f*, 623, 626
- Pogonatum tortile* 142
- Poiretia* 269
- Poitaea* 289
- Polygala* 190
  - *ambigua* 337
  - *carteri* 295
  - *cuneata* 184, 424, 559
  - *guantanama* 337, 424
  - *rhynchosperma* 370
  - *saginooides* 444, 614
  - *scabridula* 605
  - *squamifolia* 577
  - *wrightii* 577
- Polygonum portoricense* 448, 494
- Polypodium* 194, 195, 391, 395, 400, 639
  - *astrolepis* 194
  - *capillare* 194
  - *costatum* 237
  - *duale* 195, 237, 401
  - *flexuosum* 234
  - *lycopodioides* 194
  - *phyllitidis* 237, 406
  - *polypodioides* 194
- Polystachya* 190
  - *luteola* 398
- Polystichum* 194
  - *decoratum* 234
- Pontederia lanceolata* 233, 291, 487–488, 490, 492
- Ponthieva* 189, 395
- Portlandia* 266, 290
- Portulaca tuberculata* 378
- Potamogeton* 189, 478, 479, 481
  - *illinoensis* 448
  - *nodosus* 448
- Pouteria dityoneura* 409 *f*
  - *domingensis* 547, 548 *f*
- Prescottia* 395
- Prestoea montana* 185, 225, 389, 395, 621
- Priamosia* 289
- Prionodon densus* 400
- Pristimera coriacea* 553
- Pritchardia* 261
- Prockia crucis* 229
- Prosperpinaca palustris* 233
- Prosopis juliflora* 343, 617
- Protium maestrense* 406
- Proustia* 269
- Prunus* 267
  - *occidentalis* 407 *f*, 409 *f*
- Pseudocarpidium* 286, 350, 559
  - *avicennioides* 348, 375, 383 *f*, 424, 563
  - *ilicifolium* 222, 599, 602
  - *multidens* 375, 424, 563
  - *neglectum* 554
  - *pungens* 596
  - *shaferi* 385, 563
  - *wrightii* 225, 420, 424
- Pseudolmnia* 265
  - *spuria* 184, 228, 407, 408 *f*, 409 *f*, 530

- Psidium* 184, 267
  - *celastroides* 334
  - *cymosum* 316, 433, 582
  - *guava* 231, 442, 612
  - *havanense* 219, 327, 602–604
  - *montanum* 391
  - *munizianum* 219
  - *nummularium* 314, 321
  - *parvifolium* 437
  - *salutare* 432, 583, 585
  - *scopulorum* 553
  - *thomasianum* 553
  - *vicentinum* 314, 417, 553
- Psilotum nudum* 237
- Psychotria* 184, 222, 254, 265
  - *banaoana* 335
  - *ekmanii* 372, 623
  - *graminifolia* 590
  - *grandis* 407 f
  - *guadelupensis* 391, 529
  - *lopezii* 354
  - *martii* 219, 334, 395, 419
  - *moaënsis* 396
  - *nutans* 187
- Pteridium* 140
  - *caudatum* 439, 576, 613
- Pteris* 194
  - *longifolia* 237
- Pterocarpus officinalis* 519
- Pterocaulon alopecuroidaeum* 443
- Pterocissus* 289
- Purdiae* 246 m, 265, 350, 351, 634
  - *cubensis* 218, 432 f, 433, 576
  - *ekmanii* 636
  - *maëstrensis* 246 m, 372
  - *microphylla* 246 m, 644
  - *moaënsis* 246 m, 626, 636
  - *nipensis* 246 m, 350
  - *parvifolia* 246 m
- Purdiae shaferi* 636
  - *velutina* 246 m, 426, 636
- Quercus* 267, 269
  - *minima* 576
  - *myrtifolia* 576
  - *oleoides* ssp. *sagraeana* 217, 254, 301 m, 305, 308, 309 f, 319, 432, 576, 583, 585
  - *pumila* 576
  - *virginiana* 576
- Rachicallis* 297, 455, 511
  - *americana* 227, 455 f, 511
- Rajania* 190
  - *baracoënsis* 351, 398, 401
  - *howardii* 435
- Randia* 415
- *aculeata* 423
- *costata* 348
- *cubana* 318
- *spinifex* 223, 323, 424, 620
- Rapanea ferruginea* (= *Myrsine coriacea*) 394 f, 441 f
- Raphanus raphanistrum* 190
- Rauvolfia cubana* 218, 326
  - *salicifolia* 352, 396, 401
- Ravenala* 262
- Ravenia clementiana* 337
  - *simplicifolia* 354
- Reichenbachianthus emarginatus* 391
- Reimarocholoa brasiliensis* 230
- Reynaudia filiformis* 442
- Reynosia* 267, 285, 350
  - *canagüeyensis* 220, 341
  - *intermedia* 604
  - *microphylla* 327
  - *mucronata* 252 m
  - - ssp. *azulensis* 252 m, 380, 559
  - - ssp. *mucronata* 252 m
  - - ssp. *nipensis* 252 m
  - *retusa* 599, 602
- Rhabdadenia biflora* 451, 516
- Rhacoma* (*Crossopetalum*) *ekmanii* 331
- Rhamnidium potrerilloanum* 557
- *pruinatum* 361
- Rheedia* (*Garcinia*) 262
  - *fruticosa* 433, 599
  - *ophiticola* 439
  - *polyneura* 401
- Rhexia* 290
  - *cubensis* 498
- Rhipidopteris peltata* 195, 237, 391
- Rhipsalis* 416
  - *baccifer* 339 f
- Rhizogonium spiniforme* 400, 641
- Rhizophora* 161, 296
  - *mangle* 231, 332 f, 360 f, 383 f, 451 f, 516
- Rhizophora occidentalis* 515
- Rhodocactus cubensis* 221, 331, 429, 620
- Rhodogerion* 289, 324
  - *coronopifolius* 330
- Rhodopis* 289
- Rhoeo discolor* 228
- Rhus copallina* ssp. *leucantha* 226, 270, 305, 432, 583
- Rhynchoselytrum repens* (*roseum*) 134, 324, 442, 612
- Rhynchospora* 189, 226, 354, 427, 435, 443, 444, 484, 494, 577, 613
  - *brachychaeta* 498
  - *caphalotooides* 583
  - *cernua* 401
  - *corniculata* 487
  - *cubensis* 615

- *cyperoides* 443, 492, 498
- *diodon* 440, 443, 614, 615
- *divergens* 499
- *gigantea* 487, 492
- *leptorrhyncha* 498
- *lindeniana* 437, 440
- *marisculus* 398
- *nipensis* 440
- *plumosa* 615
- *pruinosa* 401
- *shaferi* 401, 440
- *stellata* 488, 496
- *tenuis* 433, 440, 443, 498, 613
- *traceri* 498
- Rhytidophyllum* 188, 285, 350, 418
- *acunae* 380
- *crenulatum* 329
- *exsertum* 327, 418, 555
- *intermedium* 351
- *lomense* 333
- *minus* 379
- *mogoticola* 374, 419
- *rupincola* 314, 416, 417
- *wrightianum* 218, 326, 333
- Richardia arenicola* 499
- *ciliata* 499
- *pectidifolia* 610
- Ricinus communis* 234
- Ritterocereus* 187
- *hystrix* 348, 383 f., 424, 428 f., 429, 617, 620
- Rivina humilis* 572
- Rochefortia* 285
- *oblongata* 382
- *spinosa* 553
- *stellata* 350
- Roigella* 268, 289, 295, 304
- *correifolia* 299 f., 305, 432, 576, 583
- Rondeletia* 184, 222, 254, 265, 268, 285, 350, 351, 369, 614, 631
- *alaternoides* 624, 636
- *apiculata* 380, 564
- *baracoensis* 351
- *bicolor* 335
- *calophylla* 631
- *camarioca* 249, 325, 425, 595, 599
- *chamaebuxifolia* 319, 582, 595, 599
- *convoluta* 335
- *gamboana* 345
- *ingrata* 385, 564
- *intermedia* 373
- *leonis* 332, 631
- *lomensis* 351
- *longibracteata* 433
- *microphylla* 631
- *myrtacea* 435 f., 590
- *naguensis* 370
- *norlindii* 380, 381, 382
- *odorata* 185, 218, 333, 583
- *pachyphylla* 398, 634, 636
- *pedicularis* 331, 332, 631
- *peduncularis* 631
- *peninsularis* 378
- *potrerilloana* 333–334, 557
- *rugelii* 329
- *savannarum* 343, 595, 599, 610
- *sevillensis* 379
- *shaferi* 343, 634
- (*Suberanthus*) *stellata* 359
- *susannae* 314
- *vacciniifolia* 636
- *vazquezii* 598
- *venosa* 316, 599
- Roystonea* 157, 160, 296, 297, 348 f., 350, 442, 446, 448
- *regia* 161, 185, 225, 324, 389, 390, 405, 408 f., 409 f., 414 f., 416 f., 442, 446, 530, 534, 612
- Rubus turquinensis* 370 f., 641
- Ruellia* 494
- Ruppia maritima* 233, 503
- Sabal* 107, 121, 157, 161, 162, 267, 296, 297, 348 f., 446, 448, 450 f.
- *parviflora* 185, 222 m., 223 f., 298 f., 420, 444, 446, 448, 522, 525, 613
- *japa* 222, 446
- Sacciolepis striata* 496
- Sachsia* 286, 498
- *polycephala* 444, 576
- Sagittaria* 494
- *intermedia* 487
- *lancifoia* 487, 488, 489 f.
- Salacia nipensis* 343
- *wrightii* 368
- Salicornia ambigua* 513
- *perennis* 233, 451, 513
- Salix caroliniana* (*longipes*) 448, 450 f., 521
- Salmea caleoides* 314
- *glaberrima* 314
- *umbratilis* 314
- Salvia speirematoides* 641
- Salvinia* 470, 482, 513
- *auriculata* 194, 469
- *natans* 194, 237
- *rotundifolia* 469
- Samanea* 157, 160, 442
- *saman* 231, 324, 405, 539, 612
- Samuelssonia* 289
- Samyda ramosissima* 348
- Sapium erythrococcum* 370
- *jamaicense* 405, 406, 407 f., 408 f.
- *leucogynum* 218, 248, 333, 417, 419, 554
- *maestrense* 369, 399, 640

- *pallens* 248
- Sapphoia* 128, 289, 352 m, 359
- *ekmanii* 352 m
- *rigidifolia* 352 m, 356, 631
- Sarcomphalus* 254, 267, 268, 285
- *acutifolius* 337
- *obovatus* 347
- Sarcopilea* 289
- Satureja (Micromeria) bucheri* 372
- Saugetia* 287
- Sauvagesia brownei* 440
- Sauvallea* 310
- Sauvallella* 128, 268, 289, 310, 316
- *immarginata* 433, 582
- Savia* 262, 297, 424
- *bahamensis* 560
- *cuneifolia* 397 f
- *erythroxyloides* 418, 557
- *sessiliflora* 225, 412, 419, 514
- Scaevela plumieri* 507, 571
- *wrightii* 437, 438 f
- Schaefferia ephedroides* 566, 569 f
- *frutescens* 420
- Schlotoheimia torquata* 142
- Schmidtiota* 128, 266, 268, 289, 351, 359, 634, 643
- *cubensis* 354, 636
- *monantha* 636
- *monticola* 644
- *multiflora* 361, 440
- *sessiliflora* 437, 644
- *shaferi* 587, 591
- *uliginosa* 636
- Schoenoplectus americanus* 487, 490
- Schoepfia* 350
- *chrysophylloides* 226
- *stenophylla* 404
- Schradera* 395
- *cubensis* 369, 398
- Schultesia guianensis* 443
- Schwenckia* 269
- Scirpus olneyi* 233, 448
- *validus* 233, 448, 487
- Scleria* 226, 442, 444, 498, 613
- *ciliata* 499, 613
- *cubensis* 585
- *georgiana* 616
- *havanensis* 325
- *interrupta* 499
- *motenboensis* 219
- *pauciflora* 499
- *pilosissima* 391
- *secans* 391, 398
- Scolosanthus* 254, 266, 268, 285
- *acunae* 317, 583, 599, 602
- *bahamensis* 225, 617
- *crucifer* 324, 595, 599
- *granulatus* 368
- *hirsutus* 385
- *hispidus* 385
- *lucidus* 440, 593
- *maestrensis* 369, 400, 640
- *maisianus* 385
- *strictus* 385
- *wrightianus* 362
- Scutachne* 287
- Sebastiana corniculata* 443
- Securidaca elliptica* 223, 412
- Selaginella* 194, 195
- *apus* 237
- *stolonifera* 235
- Selenicereus* 416
- *boeckelmannii* 410 f
- *brevispinus* 347
- *grandiflorus* 187, 417, 419, 545, 551 f
- *urbanianus* 419
- Selleola* 289
- Selleophytum* 289
- Senecio* 188, 222, 359, 626
- *ekmanii* 356
- *polyphlebius* 398, 626
- *rivalis* 398, 626
- *saugetii* 362
- *subsquarrosum* 354
- *trichotomus* 598, 439
- *vulgaris* 234
- Serjania* 187
- *crassinervis* 419
- *diversifolia* 229, 412, 414 f, 419
- *occidentalis* 291
- Sesuvium maritimum* 455, 510
- *portulacastrum* 231, 332 f, 455, 510
- Setaria* 444
- *geniculata* 442, 612, 613
- *tenax* 442, 612
- Shaferia* 128, 135, 268, 289, 358
- *platyphylla* 358, 396, 440, 594, 644
- Shaferocharis* 128, 266, 289, 352 m, 358
- *cubensis* 352 m, 358, 644
- *multiflora* 185, 352 m, 358, 360 f, 426, 597
- *villosa* 352 m, 358, 426
- Sida linifolia* 190
- *rhombifolia* 190
- Siemensia* 289, 310, 313
- *pendula* 311 f, 313, 319, 416 f, 417, 553
- Simaruba* 265
- *laevis* 397 f
- Siphocampylus* 351, 362, 365, 368, 557, 587
- *baracoensis* 361
- *glaber* 643 f, 644
- *impressus* 365 m
- *libanensis* 365 m
- *manettiiifolius* 364, 365 m

- *yumuriensis* 362, 364 f, 365 m
- Stoeane amygdalina* 223, 405, 406, 530
- *berteriana* 391
- *curatellifolia* 184, 389, 390, 397 f, 623
- Smilax domingensis* 407 f
- *havanensis* 226, 412, 420
- *laurifolia* 522
- Solandra longiflora* 412, 420, 554, 555 f
- Solanum aculeatum* 444
- *bahamense* 420
- *maëstrense* 369
- *nigrum* 233
- Solidago stricta* 496
- Solonia* 289, 369
- *reflexa* 395, 623
- Sorghastrum setosum* 612
- *stipooides* 230
- Spaniopappus* 266, 289
- *ekmanii* 364
- *iodostylus* 333
- Spartina juncea* 453, 507, 515
- Spathelia* 128, 140, 187, 222, 246 m, 256, 265, 286, 350, 351
- *brittonii* 186 f, 187, 217, 246 m, 313, 417, 553
- *cubensis* 246 m, 401
- *leonis* 246 m, 361
- *lobulata* 246 m, 366
- *pinetorum* 246 m, 396, 596
- *splendens* 246 m, 596
- *stipitata* 246 m
- *subintegra* 246 m
- *vernicosa* 246 m
- *wrightii* 246 m
- *yumuriensis* 246 m, 362
- Spermatoce exasperata* 368
- *matanzasia* 327
- Spiranthes* 189
- *elata* 229
- Spirotecomia* 287, 349
- *apiculata* 427, 440, 593
- *holguinensis* 342, 610
- *spiralis* 222, 375, 424, 559, 620
- Spondias* 160, 297
- *mombin* 161, 184, 442, 539, 546 f, 612
- Sporobolus indicus* 332 f, 442, 612
- *virginicus* 453, 506, 515
- *ssp. littoralis* 515
- Stahlia* 287
- Stegnosperma scandens* 412
- Stelis* 190, 641
- *cubensis* 400
- Stemodia maritima* 506
- Stenandrium* 189, 262, 498
- *crenatum* 337
- *droseroides* 615
- *ovatum* 579, 616
- *pinetorum* 308
- Stenochlaena sorbifolia* 235
- Sterculia apetala* 233
- *caribaea* 519
- Stevensia* 289
- Stigmaphyllon* 425
- *coccolobaeifolium* 385
- *lineare* 350
- *sagraeanum* 620
- *sericeum* 350
- Strempheliopsis* 287
- *stremphelioides* 419
- Strumpfia* 285
- *maritima* 227, 455, 511, 512
- Styleanthes hamata* 443, 444, 612
- Suaeda fruticosa* 233, 513
- *linearis* 451, 513
- Suberanthus* 268, 287
- *brachycarpus* 595
- *canellifolius* 354, 587
- *neriifolius* 218, 219 m
- *stellatus* 427, 437, 587, 598, 599 f
- Suriana* 454 f, 509
- *maritima* 332 f, 453, 507
- Swietenia* 157, 161, 265, 296, 376 f, 446
- *mahagoni* 184, 226, 228, 229 f, 346, 418, 420, 446, 530, 534, 551 f
- Sympisia alainii* 401
- Symplocos leonis* 370, 399
- Synapsis* 289, 350
- *ilicifolia* 557
- Syngonanthus* 218, 267, 290
- *androsaceus* 499
- *insularis* 431, 499, 578
- *lagopodioides* 499
- *leoni* 301, 499
- *wilsonii* 295, 499
- Syringodium filiforme* 505
- Syrrhopodon elongatus* 142
- *tenuifolius* 142
- Syzygium jambos* 234, 335, 446
- Tabebuia* 136, 161, 184, 188, 222, 254, 295, 296, 297, 354, 359, 541, 558, 559
- *acunana* 379
- *anafensis* 218, 327
- *angustata* 414 f, 446, 448, 523
- *anisophylla* 415
- *arenicola* 350
- *arimaoensis* 332
- *bibracteolata* 374, 376 f, 418, 557
- *calcicola* 552
- *capotei* 291, 560, 561
- *clementis* 626
- *dubia* 396, 397 f, 593, 626
- *elongata* 370

- *geronensis* 308
- *glaucescens* 332
- *hypoleuca* 350, 369
- *jackiana* 552
- *jaucoensis* 385
- *leonis* 385
- *lepidophylla* 218, 305, 306 f, 431, 432, 576, 585
- *lepidota* 225, 325, 425, 582, 595, 599, 605 f, 614
- *leptopoda* 316, 433
- *libanensis* 350, 380, 382 f, 564
- *linearis* 360, 426, 597
- *mogotensis* 367
- *myrtifolia* 424, 545, 548, 551 f, 554, 560
- *oligolepis* 369, 394 f, 622
- *petrophila* 552, 554
- *picotensis* 367
- *pinetorum* 439, 587
- *polynympha* 424, 564
- *rivularis* 638
- *sauvallei* 218, 333, 419
- *shaferi* 407 f, 412, 413 f, 435, 530
- *simplicifolia* 351, 587
- *trachycarpa* 337, 383 f, 415, 599, 605, 607 f
- *truncata* 348
- *turquinensis* 400
- Tabernaemontana amblyocarpa* 184, 185 f, 323, 414 f, 420, 421 f, 446
- Talauma* 265, 621
  - *minor* 350 m, 394 f, 621
  - - ssp. *minor* 135
  - - ssp. *oblongifolia* 135, 358, 396, 397 f, 623
  - *orbiculata* 369
  - *truncata* 369
- Tamarindus indica* 234
- Tapura cubensis* 396, 401
- Taraxacum officinale* 234
- Tectaria* 194
- Teobroma cacao* 233
- Tephrosia clementis* 382
- Terminalia* 161
  - *catappa* 234
  - *intermedia* 530
  - *maestrensis* 528
  - *neglecta* 419, 420, 552
  - *nipensis* 397 f, 593, 623
  - *orientensis* 593, 623
  - *pachystyla* 623
- Ternstroemia baracoensis* 361
  - *flavescens* 440
  - *leonis* 400, 640
  - *monticola* 394 f, 400, 640, 641
  - *peduncularis* ssp. *obovalis* 125
- Tetralix* 128, 268, 289, 351
- Tetramicra simplex* 223
- Tetranthus* 286
- Tetraperone* 266, 289, 295
- Tetrapteris buxifolia* 621
- Tetrasiphon* 290
- Tetrazygia* 285
  - *aurea* 335
  - *bicolor* 226
  - *coriacea* 316, 433, 582
  - *cristalensis* 396
  - *delicatula* 432, 576, 577, 578
  - *elegans* 640
  - *lanceolata* 321, 553
  - *minor* 314
- Tetrazygiopsis* 285
  - *laxiflora* 248
- Teucrium canadense* 233
  - *cubense* 233
- Thalassia testudinum* 189, 230, 332 f, 505
- Thalia geniculata* 298, 492, 494
- Thelypteris* 194
  - *palustris* 496
  - *pinnata* 235
- Theophrasta* 289
- Thespesia populnea* 332 f, 571
- Thevetia peruviana* 233
- Thogsennia* 266, 287
- Thouinia* 267, 557
  - *canescens* 552
  - *clarensis* 419
  - *elliptica* 547
  - *leonis* 385, 564, 568
  - *pseudopunctata* 348, 424, 564
  - *nervosa* (*trifoliata*) 418, 552
  - *stricta* 564
- Thrinax* 249, 250 m, 285, 297, 351
  - (*Hemithrinax*) *compacta* 250 m, 366, 557
  - *drudei* 249, 250 m
    - *ekmaniana* 250 m, 344
    - *morrisii* (*punctulata*) 249, 250 m, 275, 313, 327, 416 f, 417, 418, 552, 553, 555
    - *radiata* (*wendlandiana*) 185, 249, 250 m, 332 f, 420, 550, 571, 572
    - *rivularis* 250 m, 358
      - - ssp. *savannarum* 358, 597
      - *savannarum* 250 m
  - *Thuinidium pulverulentum* 620
  - Thymopsis thymoides* 337, 444
  - Tillandsia* 190, 392, 412, 417, 446, 534
    - *balbisiana* 425
    - *caribaea* 227
    - *circinnata* 425
    - *fasciculata* 228, 405
    - *flexuosa* 425
    - *polystachya* 405
    - *pruinosa* 425
    - *pulchella* 227
    - *recurvata* 405, 425
    - *tenuifolia* 405