



- 99.Morisita, M. Measuring of the dispersion and analysis of distribution patterns. Memoires of the Faculty of Science, Kyushu University, Series E. Biology, 2: 215–235. 1959.
- 100.Mueller-Dombois, D., Ellenberg, H. Aims and Methods of Vegetation Ecology. John Wiley & Sons, New York, 1974, 547 p.
- 101.Muller, P. Introducción a la zoogeografía. Ed. Blume, Barcelona, 1974, 216 p.
- 102.Murray, S.N., Ambrose, R.F., Dethier, M.N. Methods for Performing Monitoring, Impact, and Ecological studies on Rocky Shores. MMS OCS Study 2001-070. Coastal Research Center, Marine Science Institute, University of California, Santa Barbara, California, 2002, 73 p.
- 103.Nava, A.F.D. Espécies sentinelas para a Mata Atlântica: as consequências epidemiológicas da fragmentação florestal no Pontal do Paranapanema, São Paulo. Tese (doutorado) - Universidade de São Paulo. Faculdade de Medicina Veterinária e Zootecnia. Departamento de Medicina Veterinária Preventiva e Saúde Animal, 2008, 147 p.
- 104.Olabarria, C., Chapman, M.G. Comparison of patterns of spatial variation of microgastropods between two contrasting intertidal habitats. Mar. Ecol. Progr. Ser., 220: 202-211. 2001.
- 105.Odum, E.P. The strategy of ecosystem development. Science, 164(3877): 262-270. 1969.
- 106.Olsgard, F., Somerfield, P.J., Carr, M.R. Relationships between taxonomic resolution and data transformations in analyses of a macrobenthic community along an established pollution gradient. Mar. Ecol. Prog. Ser., 149: 173-181. 1997.
- 107.Orfanidis, S., Panayotidis, P., Stamatis, N. Ecological evaluation of transitionnal and coastal waters: a marine benthic macrophytes-based model. Medit. Mar. Sci. 2(2): 45–65. 2001.
- 108.Orfanidis, S., Panayotidis, P., Stamatis, N. An insight to the ecological evaluation index (EEI). Ecological Indicators, 3: 27-33. 2003.
- 109.Orlando-Bonaca, M., Lipej, L., Orfanidis, S. Benthic macrophytes as a tool for delineating, monitoring and assessing ecological status: The case of Slovenian coastal waters. Mar. Poll. Bull., 56: 666-676. 2008.
- 110.Padilla, D.K., Allen, B.J. Paradigm lost: reconsidering functional form and group hypotheses in marine ecology. J. Exp. Mar. Biol. Ecol., 250: 207–221. 2000.
- 111.Panayotidis, P., Montesanto, B., Orfanidis, S. Use of low-budget monitoring of macroalgae to implement the European Water Framework Directive. J. Appl. Phycol., 16: 49-59. 2004.



- 112.Pearson, T.H., Rosenberg, R. Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. *Oceanogr. Mar. Biol. Ann. Rev.*, 16: 229-311. 1978.
- 113.Peterson, A.T. Predictability of the geography of species' invasions via ecological niche modeling. *Q. Rev. Biol.*, 78: 419-433. 2004.
- 114.Peterson, A.T., Ball, L.G., Cohoon, K.P. Predicting distributions of Mexican birds using ecological niche modelling methods. *Ibis* (2002), 144 (on-line), E27-E32.
- 115.Phillips, J.C., Kendrick, G.A., Lavery, P.S. A test of a functional group approach to detecting shifts in macroalgal communities along a disturbance gradient. *Mar. Ecol. Prog. Ser.*, 153: 125-138. 1997.
- 116.Pinedo, S. García, M., Satta, M.P., de Torres, M., Ballesteros, E. Rocky-shore communities as indicators of water quality: A case study in the Northwestern Mediterranean. *Mar. Poll. Bull.*, 55: 126-135. 2007.
- 117.Preston, F.W. The commonness and rarity of species. *Ecology*, 29: 254-283. 1948.
- 118.Salen-Picard, C. Schémas d'évolution d'une biocenose macrobenthique du substrat meuble. *Comptes Rendus de l'Academie des Sciences de Paris*, 296: 587-590. 1983.
- 119.Santos, A., Cocentino, A L. M., Reis, T. N. V. Macroalgas como indicadoras da qualidade ambiental da praia de Boa Viagem, Pernambuco, Brasil. *Bol. Téc. Cient. CEPENE, Tamandaré*, 14(2): 25-33. 2006.
- 120.Scheiner, S.M. Six types of species-area curves. *Global Ecol. Biogeogr.*, 12: 441-447. 2003.
- 121.Schoener, T.W. The species-area relation within archipelagos: models and evidence from island land birds. *Proceedings of the 16th International Ornithological Conference, Australian Academy of Science, Canberra*. 629-642 p. 1976.
- 122.Schramm, W., Nienhuis, P.H. Marine benthic vegetation: recent changes and the effects of eutrophication. Springer-Verlag, Berlin, Germany. 1996, 480 p.
- 123.Scott, J.M., Heglund, P.J., Morrison, M.L. Predicting species occurrences: issues of accuracy and scale. Washington, DC, Island Press. 2002, 840 p.
- 124.Sfriso, A., Facca, C., Ghetti, P.F. Rapid Quality Index (R-MaQI), based mainly on macrophyte associations, to assess the ecological status of Mediterranean transitional environments. *Chem. Ecol.*, 23(6): 493-503. 2007.
- 125.Silva, I.B. Diversidade de macroalgas marinhas bentônicas dos recifes de Maraçajá, área de preservação ambiental dos recifes de corais, Rio Grande do Norte, Brasil. Univ. Federal do Rio Grande do Norte. Centro de Biociências. Departamento de





Oceanografia e Limnologia. Programa de Pós Graduação em Bioecologia Aquática, Natal. 2006, 58 p.

126. Smith, E.P. BACI design. In: El-Shaarawi, A.H., Piegorsch, W.W. (eds.), Encyclopedia of Environmetrics. John Wiley & Sons, Chichester, England, 141-148 p. 2002.

127. Snedecor, G.W., Cochran, W.G. Statistical Methods. Eighth Edition. Iowa State University Press, Ames, Iowa, 1989, 503 p.

128. Soares, L.P., Reis, T.N.V., Barros, N.C.G., Araujo, P.G., Silva, I.B., Vasconcelos, E.R.T.P.P., Cocentino, A.L.M., Areces, A.J., Fujii, M.T. Índice de qualidade ambiental utilizando macroalgas como bioindicadoras nas praias de Boa Viagem, Suape e Tamandaré, Pernambuco, Brasil. IV Congresso Brasileiro de Oceanografia, 2010, Rio Grande. AOCEANO, 02986-02988. 2010a.

129. Soares, L.P., Reis, T.N.V., Bernardi, J., Barros, N.C.G., Minei, C.C., Areces, A.J., Guimarães, S.M.P.B., Fujii, M.T. Macroalgas marinhas bentônicas como indicadoras da qualidade ambiental nas praias do litoral norte do Estado de São Paulo, Brasil. IV Congresso Brasileiro de Oceanografia, 2010, Rio Grande. AOCEANO, 02351-02353. 2010b.

130. Soberón, J., Peterson, A.T. Biodiversity informatics: Managing and applying primary biodiversity data. *Philos. T. Roy. Soc. B.*, 359(1444): 689-698. 2004.

131. Soltan, D., Verlaque, M., Boudouresque, C.F., Francour, P. Changes in macroalgal communities in the vicinity of the Mediterranean sewage outfall after the setting up of a treatment plant. *Mar. Poll. Bull.*, 42, 59-70. 2001.

132. Sørensen, T. A method of stabilizing groups of equivalent amplitude in plant sociology based on the similarity of species content and its application to analysis of the vegetation of Danish commons. *Biol., Skr.*, 5(4): 1-34. 1948.

133. Souza, G., Cocentino, A. Macroalgas como indicadoras da qualidade ambiental da Praia de Piedade-PE. *Tropical Oceanography*, 32(1):1-22. 2004.

134. Steneck, R.S., Dethier, M.N. A functional group approach to the structure of algal-dominated communities. *Oikos*, 69: 476-498. 1994.

135. Taquil, A., Yoneshigue-Valentin, Y. Alterações na composição florística das algas da Praia de Boa Viagem (Niterói, RJ). *Braz. J. Bot.*, 25(4): 405-412. 2002.

136. Ulanowicz, R.E., Kay, J.J. A package for the analysis of ecosystems flows networks. *Environ. Softw.*, 6(3): 131-142. 1991.

137. Ulanowicz, R.E., Kemp, W.M. Toward canonical trophic aggregations. *Am. Nat.* 114 (6): 871-883. 1979.



- 138.Underwood, A.J. "Beyond BACI": experimental designs for detecting human environmental impacts on temporal variations in natural populations. *Aust. J. Mar. Fresh. Res.*, 42: 569-587. 1991.
- 139.Underwood, A.J., Peterson, C.H. Towards an ecological framework for investigating pollution. *Mar. Ecol. Prog. Ser.*, 46: 227-234. 1988.
- 140.Valle, R., Campos, I., Areces, A.J. Comunidades bentónicas del mesolitoral. In *Resultados, Proyecto PNUD-PNUMA-UNESCO CUB/80/001*, 2(2): 624-640. 1985.
- 141.Vasconcelos, E.R.T.P.P. Índice de Distúrbio Ambiental (IDA) através das Macroalgas Marinhas Bentônicas dos Recifes Areníticos de Pernambuco. Dissertação ao grau de Mestre em Oceanografia Biológica, Programa de Pós-graduação em Oceanografia, Universidade Federal de Pernambuco, Recife, 2012, 60 p.
- 142.Warwick, R.M. Analysis of community attributes of the macrobenthos of Frierfjord/Langesundsfjord at taxonomic levels higher than species. *Mar. Ecol. Progr. Ser.*, 46: 167-170. 1988a.
- 143.Warwick, R.M. The level of taxonomic discrimination required to detect pollution effects on marine benthic communities. *Mar. Poll. Bull.*, 19: 259-268. 1988b.
- 144.Wells, E., Wilkinson, M., Wood, P., Scanlan, C. The use of macroalgal species richness and composition on intertidal rocky seashores in the assessment of ecological quality under the European Water Framework Directive. *Mar. Poll. Bull.*, 55, 151-161. 2007.
- 145.Water Framework Directive (WFD). Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for community action in the field of water policy. OJ L 327, 22.12.2000, 2000, 82 p.
- 146.Wolda, H. Similarity indices, simple size and diversity. *Oecologia*, 50: 296-302. 1981.

